

Runs of Homozygosity and Testicular Cancer Risk

Chey Loveday¹, Amit Sud¹, Kevin Litchfield², Max Levy¹, Amy Holroyd¹, Peter Broderick¹, Zsofia Kote-Jarai¹, Alison M Dunning³, Kenneth Muir^{4,5}, Julian Peto⁶, Rosalind Eeles^{1,7}, Douglas F Easton^{3,8}, Darshna Dudakia¹, Nick Orr⁹, Nora Pashayan¹⁰, UK Testicular Cancer Collaboration[†], The PRACTICAL Consortium[†], Alison Reid¹¹, Robert A Huddart¹², Richard S Houlston¹ and Clare Turnbull^{1,13,14,15*}

¹Division of Genetics & Epidemiology, The Institute of Cancer Research, London, UK.

²Translational Cancer Therapeutics Laboratory, The Francis Crick Institute, 1 Midland Rd, London, UK.

³Centre for Cancer Genetic Epidemiology, Department of Oncology, University of Cambridge, Cambridge, UK.

⁴Division of Health Sciences, Warwick Medical School, Warwick University, Warwick, UK.

⁵Institute of Population Health, University of Manchester, Manchester, UK.

⁶Department of Non-communicable Disease Epidemiology, London School of Hygiene and Tropical Medicine, London, UK.

⁷Royal Marsden NHS Foundation Trust, London, UK.

⁸Centre for Cancer Genetic Epidemiology, Department of Public Health and Primary Care, University of Cambridge, Cambridge, UK.

⁹The Breast Cancer Now Toby Robins Research Centre, The Institute of Cancer Research, London, UK.

¹⁰Department of Applied Health Research, University College London, London, UK.

¹¹Academic Uro-oncology Unit, The Royal Marsden NHS Foundation Trust, Sutton, Surrey, UK.

¹²Academic Radiotherapy Unit, Institute of Cancer Research, Sutton, Surrey, UK.

¹³William Harvey Research Institute, Queen Mary University, London, UK.

¹⁴Guys and St Thomas' NHS Foundation Trust, Great Maze Pond, London, UK.

¹⁵Public Health England, National Cancer Registration and Analysis Service, London, UK.

[†]A comprehensive list of consortium members appears at the end of the paper.

*Corresponding author. Division of Genetics and Epidemiology, The Institute of Cancer Research, London, SM2 5NG, UK; Tel: +44 (0) 208 722 4485; E-mail: clare.turnbull@icr.ac.uk (C. Turnbull).

Keywords: Testicular germ cell tumour, cancer, GWAS, Runs of homozygosity, homozygosity mapping, recessive, genetics.

Abstract

Background

Testicular germ cell tumour (TGCT) is highly heritable but >50% of the genetic risk remains unexplained. Epidemiological observation of greater relative risk to brothers of men with TGCT compared to sons has long alluded to recessively acting TGCT genetic susceptibility factors, but to date none have been reported. Runs of homozygosity (RoH) are a signature indicating underlying recessively acting alleles and have been associated with increased risk of other cancer types.

Objective

To examine if RoH are associated with TGCT risk.

Methods

We performed a genome-wide RoH analysis using GWAS data from 3,206 TGCT cases and 7,422 controls uniformly genotyped using the OncoArray platform.

Results

Global measures of homozygosity were not significantly different between cases and controls, and the frequency of individual consensus RoH were not significantly different between cases and controls, after correction for multiple testing. RoH at three regions, 11p13-11p14.3, 5q14.1-5q22.3 and 13q14.11-13q.14.13, were however nominally statistically significant at $P < 0.01$. Intriguingly, RoH200 at 11p13-11p14.3 encompasses *Wilms tumor 1 (WT1)*, a recognized cancer susceptibility gene with roles in sex determination and developmental transcriptional regulation, processes repeatedly implicated in TGCT etiology.

Discussion and Conclusion

Overall, our data does not support a major role in the risk of TGCT for recessively acting alleles acting through homozygosity, as measured by RoH in outbred populations of cases and controls.

Introduction

Testicular germ cell tumor (TGCT) is the most common cancer in young men with over 52,000 new cases diagnosed annually worldwide (Le Cornet et al. 2014). TGCT has a strong heritable basis, as evidenced by the 4 to 8-fold increased risk of TGCT seen in first-degree relatives of TGCT patients (Litchfield, Thomsen, et al. 2015; Hemminki and Li 2004; Swerdlow et al. 1997; McGlynn et al. 2005; Kharazmi et al. 2015). Statistical analyses of heritability estimate that genetic factors may contribute to approximately half of all TGCT disease risk (Litchfield, Thomsen, et al. 2015).

Early linkage analyses in familial TGCT did not support existence of a major Mendelian TGCT susceptibility locus, but these studies were limited in power on account of modest sample sizes and the low prevalence of multiplex TGCT pedigrees (Crockford et al. 2006; Rapley et al. 2003; Rapley et al. 2000). More recently large-scale exome sequencing studies have also failed to identify rare high-penetrance susceptibility alleles, despite improved power compared to previous linkage analyses (Litchfield, Loveday et al. 2018). Nevertheless, neither analysis excludes the possibility that susceptibility genes/alleles for TGCT of lower frequency and/or more moderate effect size may exist. Indeed, very rare alleles in ciliary microtubule genes have recently been implicated through functional analyses in TGCT susceptibility in a minority of familial cases (Litchfield, Levy, Dudakia, et al. 2016).

However, collectively findings are consistent with advanced analyses of TGCT heritability, which have indicated that the genetic component of TGCT heritability is largely constituted by common variants. Recent genome-wide association studies (GWAS) have made substantial progress in exposition of this partition of heritability with 49 independent TGCT risk loci identified, together accounting for ~37% of the excess genetic risk of disease (Loveday et al. 2018; Wang et al. 2017; Litchfield et al. 2017; Litchfield, Levy, Orlando, et al. 2016; Litchfield, Shipley, and Turnbull 2015; Litchfield, Sultana, et al. 2015; Litchfield, Holroyd, et al. 2015; Koster et al. 2014; Ruark et al. 2013; Turnbull et al. 2010; Rapley et al. 2009; Kanetsky et al. 2009; Kristiansen et al. 2015; Schumacher et al. 2013; Turnbull and

Rahman 2011). These TGCT susceptibility loci have provided invaluable insight into the biology of TGCT susceptibility, implicating as underlying mechanisms, widespread transcriptional dysregulation linked to developmental arrest of primordial germ cells, aberrant KIT-MAPK signaling and defective microtubule function (Litchfield et al. 2017). From these GWAS loci approximately half of the genetic component of TGCT heritability has been accounted for, with heritability analysis indicating that the outstanding ‘missing heritability’ of TGCT is likely polygenic, with substantial contribution from common variation (Litchfield et al. 2017; Litchfield, Mitchell, et al. 2016; Litchfield, Thomsen, et al. 2015).

GWAS analysis has likewise made substantial impact in delineating the genetic architecture of many other common cancers but almost uniformly the reported susceptibility loci have been identified through analyses based on a log additive (multiplicative) model of inheritance, with little evidence generated for alleles acting recessively (Sud, Kinnersley, and Houlston 2017). This observation may be a reflection that GWAS is suboptimal in its ability to detect these alleles rather than an observation truly reflective of the underlying biology. In principle, it is entirely plausible that there may be an association between recessively acting disease alleles and susceptibility to cancer. Such a hypothesis is supported by observations reporting an increased burden of cancer in the offspring of consanguineous unions and in populations with a high degree of inbreeding (Bener et al. 2009; Lebel and Gallagher 1989; Shami, Qaisar, and Bittles 1991; Simpson et al. 1981; Assie et al. 2008). Furthermore, experimental inbreeding (e.g. backcrossing mice) has also been shown to increase tumor burden in mice (Demant 2003). In addition, uniparental disomy through dysregulated imprinting is a specific situation in which homozygosity can be directly associated with cancer (Henry et al. 1991). Of note, for TGCT, there has been a long-standing hypothesis that recessive (or X-linked) susceptibility factors are highly likely to be important, based on epidemiological data that siblings’ relative disease risks are higher than parent-offspring risks (Hemminki and Li 2004; Kharazmi et al. 2015).

Homozygosity mapping provides a means of identifying recessive components of inheritance. It has been demonstrated that, on account of selective pressure, runs of homozygosity (RoH) occur at high frequency in outbred populations, the result of autozygosity (i.e. the co-location of two alleles at a given locus originating from a common ancestor by way of non-random mating) (McQuillan et al. 2008; Ku et al. 2011). These RoH can be enriched for rare deleterious variants in homozygous form (Szpiech et al. 2013); multiple susceptibility loci have been reported for different diseases, identified through genome-wide analyses for RoH of SNP array data (reviewed in [(Ceballos et al. 2018)]).

Here, we sought to identify associations between homozygosity and TGCT risk through the characterization and comparison of genome-wide homozygosity measures and specific loci identified through consensus mapping of recurrent RoH in 3,206 TGCT cases vs 7,422 controls directly genotyped for 371,504 SNPs.

Methods

Sample description

TGCT cases (n=3,206) were ascertained via two UK studies: (1) a UK study of familial testicular cancer and (2) a systematic collection of UK TGCT cases. Case recruitment was via the UK Testicular Cancer Collaboration, a group of oncologists and surgeons treating TGCT in the UK. The studies were coordinated at the Institute of Cancer Research (ICR). Samples and information were obtained with full informed consent and Medical Research and Ethics Committee approval (MREC02/06/66 and 06/MRE06/41). All experiments were performed in accordance with relevant guidelines and regulations.

Control samples for the primary GWAS were all taken from within the UK. Specifically 2,976 cancer-free, male controls were recruited through two studies within the PRACTICAL Consortium: (1) the UK Genetic Prostate Cancer Study (UKGPCS) (age <65), a study conducted through the Royal Marsden NHS Foundation Trust and (2) SEARCH (Study of Epidemiology & Risk Factors in Cancer), recruited via GP practices in East Anglia (2003-2009). 4,446 cancer-free female controls from across the UK were recruited via the Breast Cancer Association Consortium (BCAC).

GWAS

Genotyping was conducted using a custom Infinium OncoArray-500K BeadChip (OncoArray) from Illumina (Illumina, San Diego, CA, USA), comprising a 250K SNP genome-wide backbone and 250K SNP custom content selected across multiple consortia within COGS (Collaborative Oncological Gene-environment Study). OncoArray genotyping was conducted in accordance with the manufacturer's recommendations by the Edinburgh Clinical Research Facility, Wellcome Trust CRF, Western General Hospital, Edinburgh EH4 2XU.

The UK TGCT OncoArray dataset was filtered as follows: we excluded individuals with low call rate (<95%), with abnormal autosomal heterozygosity (>3 SD above the mean) or with >10% non-European ancestry (based on multi-dimensional scaling); we excluded SNPs with minor allele frequency <1%, a call rate of <95% in cases or controls or with a minor allele frequency of 1–5% and a call rate of <99%; and those deviating from Hardy-Weinberg equilibrium ($P > 10^{-12}$ in controls and 10^{-5} in cases). The final number of SNPs passing quality control filters was 371,504.

Bioinformatic and statistical analysis

Bioinformatic and statistical analyses were performed as previously described (Sud et al. 2015). Briefly, we detected RoH using PLINK v1.90 (Purcell et al. 2007), which moves a sliding window of SNPs across the entire genome. To allow for genotyping error or other sources of artificial

heterozygosity (such as paralogous sequences) within a stretch of truly homozygous SNPs, 2% heterozygous SNPs were allowed in each window. This measure was implemented to prevent underestimation of the number and size of RoH. Default parameter values were employed (including allowing 5 missing calls per window), with the exception that we varied the parameter homozyg-snp according to our heuristic preferences for defining the RoH as detailed below. Subsequent statistical analyses for comparison of frequencies of ROH were performed using packages available in R (version 3.4.1) with integration of results against genomic references executed using and custom written Perl code. Comparisons of global homozygosity measures between cases and controls were made using the Student t-test. Adjustment for multiple testing was based on the Bonferroni correction.

We used three metrics to investigate the selection pressure on each RoH. Integrated Haplotype Score (iHS) is based on Linkage disequilibrium (LD) surrounding a positively selected allele compared to background, providing evidence of recent positive selection at a locus (Voight et al. 2006). An iHS score >2.0 reflects that haplotypes on the ancestral background are longer compared to the derived allelic background. Episodes of selection tend to skew SNP frequencies in different directions and Tajima's D is based on the frequencies of SNPs segregating in the region of interest (Tajima 1989). Fixation index (Fst) measures the degree of population differentiation at a locus, taking values from 0 to 1.0 (Holsinger and Weir 2009). iHS, D and Fst metrics were obtained from dbPSHP (Li et al. 2014).

Identification of Consensus RoH

In order to focus on more commonly occurring RoH and to empower our analysis to identify meaningful associations, only RoH in which 10 or more individuals shared the same RoH were retained for these analyses. The initial search for RoH was performed using PLINK (Purcell et al. 2007) with a specified length of 68 consecutive SNPs (homozyg-snp parameter). This RoH length was chosen (i) to be more than an order of magnitude larger than the mean haploblock size in the human

genome (ii) without being so large as to be very rare. The likelihood of observing 68 consecutive chance events can be calculated as follows (Lencz et al. 2007). Mean heterozygosity in the samples was calculated to be 42%. Thus, given 371,504 SNPs and 10,628 individuals, a minimum length of 47 would be required to produce <5% randomly generated RoH across all subjects ($[1 - 0.42]^{47} \times 371,504 \times 10,628 < 0.05$). A consequence of LD is that the SNP genotypes are not always independent, thereby inflating the probability of chance occurrences of biologically meaningless ROH. Analysis based on PLINK's pairwise LD SNP pruning function showed an approximate reduction of information compared to the original number of SNPs of 25%. Thus RoH of length 68 SNPs were used to approximate the degrees of freedom of 47 independent SNP calls.

Once all RoH of at least 68 SNPs in length were identified, these were pruned to only those RoH that occurred in more than 10 individuals. To ensure that a minimum length and minimum number of SNPs in each RoH was maintained, each individual's SNP data were recoded as one if the SNP was in an RoH for that individual and zero otherwise. Then, for each SNP, those SNPs with fewer than 10 individuals coded as one were recoded to zero before removing any ROH that due to this recoding were now less than the required number of SNPs in length. This process therefore resulted in a list of "consensus" ROH having a minimum of 68 consecutive homozygous SNP calls across 10 or more samples.

Data availability

Case GWAS data (PLINK binaries) are deposited at European Genome-phenome Archive [EGA] under accession code EGAS00001001836.

Results

We have previously implemented rigorous quality control measures to the UK TGCT OncoArray GWAS dataset²², excluding samples and SNPs with poor call rates, SNPs with significant departure

from HWE, and samples of non-European ancestry or with a sex discrepancy as inferred from the data. The final dataset included 10,628 individuals from the UK and of European ancestry, comprising 3,206 TGCT cases and 7,422 controls, all genotyped on the same platform. The final number of SNPs passing quality control filters was 371,504.

Across all samples ($n=10,628$), the total number of discrete autosomal RoH >1000 kb and comprising at least 68 consecutive SNPs as identified by PLINK was 137,833, with an average number of 12.97 RoH per individual, an average size of 1630.17 Kb per RoH per individual, and an average total length of the genome covered by RoH of 21,216.01 Kb per individual. These results are broadly similar to other studies using similar methodologies (Sud et al. 2015; Hosking et al. 2010; Thomsen et al. 2016; Thomsen et al. 2015). There was no significant difference in the average number, length per RoH, or total length of RoH per individual between TGCT cases and controls when compared using Student's T test (Table 1). Likewise, the cumulative distribution of RoH was broadly similar for TGCT cases and controls (Fig. 1).

Data indicate two different types of RoH shaped by different selective pressures, with the different types characterised by different run length (Pemberton et al. 2012). Small/intermediate sized RoH (<1.6 Mb) are shaped via serial migration as a result of decreasing population size, generating LD, reducing haplotype diversity and increasing chance pairing of identical haplotypes. Conversely, long RoH (>1.6 Mb) are generated through inbreeding. There was no difference in global homozygosity measures between TGCT cases and controls when performing these analyses on RoH separated into these size categories (Table 1).

We next identified a set of 319 consensus RoH (Supplementary Table S1), that is RoH that are present in at least 10 individuals. Eight of these consensus RoH had a frequency of greater than 25% across all individuals (Table 2). The vast majority of these common consensus RoH has been

previously reported in other studies of RoH. For these RoH, selective pressure metrics are indicative of positive selection in Caucasian populations, and their locations are within genomic regions characterised by reduced numbers of structural variants and low recombination rates. The most frequently occurring RoH in our dataset (RoH164) has previously been identified as a site of selective sweep in multiple studies (Voight et al. 2006; Wang et al. 2006; Williamson et al. 2007) and is frequently identified in studies of common consensus RoH. Importantly, previous reports of these RoH provide further validity of our approach.

Fig. 2 shows the correlation between the frequency of consensus RoH in TGCT cases and controls. No consensus RoH was exclusive to either group nor significantly associated with TGCT risk after correcting for multiple testing (i.e. $P < 0.0001$). Three consensus RoH demonstrated nominal associations with TGCT at a suggestive significance level ($P < 0.01$) (Table 3). Each of these regions showed highly significant values for three estimates of selective pressure (iHS_{max}, Tajimas' D_{max}, and Fay Wu's H), indicating that these regions may have been generated as the result of a selective sweep.

The RoH with the strongest evidence of association, RoH200, was identified in 5% of TGCT cases (n=148) and 3% of controls (n=243) ($P = 0.0009$; Table 3). It comprises 866 SNPs spanning 9 Mb of chromosome 11 and encompasses 52 genes/predicted transcripts, including *Wilms Tumor 1 (WT1)*, a developmental transcription factor involved in sex determination and establishment of the urogenital system, and with established oncogenic and proto-oncogenic roles in tumor formation. To further investigate a potential link between *WT1* and TGCT risk, we performed an association analysis of individual SNPs within 25 kb of *WT1*, considering only those with an info score > 0.8 and MAF > 0.01 (n=432). The strongest putative association was for a directly genotyped SNP, rs11031783, which maps to the non-coding *WT1 antisense RNA (WT1-AS)*, OR = 1.18, $P = 0.0003$. This putative association warrants additional validation. Of note this region also contains two

additional genes related to TGC tumorigenesis: *LGR4* and *FSHB*. *LGR4* is involved in Wnt signalling and whilst variation in the FSH receptor has been implicated in TGCT susceptibility (Bang et al. 2018).

Discussion and Conclusion

In conclusion, our analyses demonstrate that levels of homozygosity are unlikely to play a substantial role in defining the risk of TGCT. Moreover, our findings suggest that existence of large numbers of recessive alleles that predispose to TGCT when unmasked by autozygosity is unlikely in outbred populations such as that of the UK. Therefore, from these analyses we are unable to provide explanation for epidemiological observation of the higher risks to siblings of cases than to other male family members. However, due to genome-wide testing and requisite correction for multiple-testing, this analysis certainly does not preclude existence of recessively acting disease alleles in TGCT risk; alternative analytic strategies will be needed to identify such alleles if they do exist. Though not statistically significant, the possible link between TGCT and an RoH hotspot that encompasses 11p.13 and *WT1* is an interesting observation that warrants further investigation.

Acknowledgements

We thank the patients with TGCT and the clinicians involved in their care for participation in this study. We thank the patients and all clinicians forming part of the UK Testicular Cancer Collaboration (UKTCC) for their participation in this study. This study would not have been possible without the contributions of M.K. Bolla (Breast Cancer Association Consortium (BCAC)), Q. Wang (BCAC), K. Michailido (BCAC), J. Dennis (BCAC), P. Hall (Collaborative Oncological Gene-environment Study (COGS)); D.F. Easton (BCAC), A. Berchuck (Ovarian Cancer Association Consortium), R. Eeles (PRACTICAL), G. Chenevix-Trench (The Consortium of Investigators of Modifiers of BRCA1/2), J. Dennis, P. Pharoah, A. Dunning, K. Muir, J. Peto, A. Lee and E. Dicks. We also thank the following for their contributions to this project: J. Simard, P. Kraft, C. Luccarini and the staff of the Centre for Genetic Epidemiology Laboratory; and K.F. Doheny and the staff of the Center for Inherited Disease

Research (CIDR) genotyping facility. The results published here are based in part on data generated by the TCGA Research Network. This study makes use of data generated by the Wellcome Trust Case Control Consortium 2 (WTCCC2).

We would also like to thank the following for funding support: The Institute of Cancer Research and The Everyman Campaign, The Prostate Cancer Research Foundation, Prostate Research Campaign UK (now Prostate Action), The Orchid Cancer Appeal, The National Cancer Research Network UK, The National Cancer Research Institute (NCRI) UK. We are grateful for support of NIHR funding to the NIHR Biomedical Research Centre at The Institute of Cancer Research and The Royal Marsden NHS Foundation Trust.

We thank all the individuals who took part in these studies and all the researchers, clinicians, technicians and administrative staff who have enabled this work to be carried out.

Funding Information

This study was supported by the Movember Foundation and the Institute of Cancer Research. A.S. is supported by a clinical fellowship from Cancer Research UK. K.L. was supported by a PhD Studentship from Cancer Research UK. R.S.H. and P.B. are supported by Cancer Research UK.

Genotyping of the OncoArray was funded by the US National Institutes of Health (NIH) [U19 CA 148537 for ELucidating Loci Involved in Prostate cancer SuscEptibility (ELLIPSE) project and X01HG007492 to the Center for Inherited Disease Research (CIDR) under contract number HHSN268201200008I]. Additional analytic support was provided by NIH NCI U01 CA188392 (PI: Schumacher).

The PRACTICAL consortium was supported by Cancer Research UK Grants C5047/A7357, C1287/A10118, C1287/A16563, C5047/A3354, C5047/A10692, C16913/A6135, European Commission's Seventh Framework Programme grant agreement n° 223175 (HEALTH-F2-2009-223175), and The National Institute of Health (NIH) Cancer Post-Cancer GWAS initiative grant: No. 1 U19 CA 148537-01 (the GAME-ON initiative).

Authors' Contributions

C.T., C.L., and A.S. designed the study. Case samples were recruited by A.R., R.A.H. and through UKTCC. R.E., A.M.D., K.M., J.P., Z.K.-J., N.P. and D.F.E. supplied OncoArray control data, via the PRACTICAL Consortium. N.O. administrated genotyping of OncoArray case samples. D.D. coordinated all case sample administration and tracking. K.L., M.L., A.H. and P.B. prepared samples for genotyping experiments. C.T., R.S.H., A.S. and C.L. designed bioinformatics and statistical analyses. C.L., K.L. and M.L. performed bioinformatics and statistical analyses. C.L. drafted the manuscript with assistance from C.T., A.S., and R.S.H. All authors reviewed and contributed to the manuscript.

Disclosure of Interest

The authors have no conflicts of interest to disclose.

References

- Assie, G., T. LaFramboise, P. Platzer, and C. Eng. 2008. 'Frequency of germline genomic homozygosity associated with cancer cases', *Jama*, 299: 1437-45.
- Bang, A. K., A. S. Busch, K. Almstrup, J. Gromoll, S. Kliesch, E. Rajpert-De Meyts, N. E. Skakkebaek, A. Juul, F. Tuttelmann, and N. Jorgensen. 2018. 'Is the FSHR 2039A>G variant associated with susceptibility to testicular germ cell cancer?', *Andrology*, 6: 176-83.
- Bener, A., H. R. El Ayoubi, L. Chouchane, A. I. Ali, A. Al-Kubaisi, H. Al-Sulaiti, and A. S. Teebi. 2009. 'Impact of consanguinity on cancer in a highly endogamous population', *Asian Pac J Cancer Prev*, 10: 35-40.
- Ceballos, F. C., P. K. Joshi, D. W. Clark, M. Ramsay, and J. F. Wilson. 2018. 'Runs of homozygosity: windows into population history and trait architecture', *Nat Rev Genet*.

- Crockford, G. P., R. Linger, S. Hockley, D. Dudakia, L. Johnson, R. Huddart, K. Tucker, M. Friedlander, K. A. Phillips, D. Hogg, M. A. Jewett, R. Lohynska, G. Daugaard, S. Richard, A. Chompret, C. Bonaiti-Pellie, A. Heidenreich, P. Albers, E. Olah, L. Geczi, I. Bodrogi, W. J. Ormiston, P. A. Daly, P. Guilford, S. D. Fossa, K. Heimdal, S. A. Tjulandin, L. Liubchenko, H. Stoll, W. Weber, D. Forman, T. Oliver, L. Einhorn, M. McMaster, J. Kramer, M. H. Greene, B. L. Weber, K. L. Nathanson, V. Cortessis, D. F. Easton, D. T. Bishop, M. R. Stratton, and E. A. Rapley. 2006. 'Genome-wide linkage screen for testicular germ cell tumour susceptibility loci', *Hum Mol Genet*, 15: 443-51.
- Demant, P. 2003. 'Cancer susceptibility in the mouse: genetics, biology and implications for human cancer', *Nat Rev Genet*, 4: 721-34.
- Hemminki, K., and X. Li. 2004. 'Familial risk in testicular cancer as a clue to a heritable and environmental aetiology', *Br J Cancer*, 90: 1765-70.
- Henry, I., C. Bonaiti-Pellie, V. Chehensse, C. Beldjord, C. Schwartz, G. Utermann, and C. Junien. 1991. 'Uniparental paternal disomy in a genetic cancer-predisposing syndrome', *Nature*, 351: 665-7.
- Holsinger, K. E., and B. S. Weir. 2009. 'Genetics in geographically structured populations: defining, estimating and interpreting F_{ST} ', *Nat Rev Genet*, 10: 639-50.
- Hosking, F. J., E. Papaemmanuil, E. Sheridan, S. E. Kinsey, T. Lightfoot, E. Roman, J. A. Irving, J. M. Allan, M. Taylor, I. P. Tomlinson, M. Greaves, and R. S. Houlston. 2010. 'Genome-wide homozygosity signatures and childhood acute lymphoblastic leukemia risk', *Blood*, 115: 4472-7.
- Kanetsky, P. A., N. Mitra, S. Vardhanabhuti, M. Li, D. J. Vaughn, R. Letrero, S. L. Ciosek, D. R. Doody, L. M. Smith, J. Weaver, A. Albano, C. Chen, J. R. Starr, D. J. Rader, A. K. Godwin, M. P. Reilly, H. Hakonarson, S. M. Schwartz, and K. L. Nathanson. 2009. 'Common variation in KITLG and at 5q31.3 predisposes to testicular germ cell cancer', *Nat Genet*, 41: 811-5.
- Kharazmi, E., K. Hemminki, E. Pukkala, K. Sundquist, L. Tryggvadottir, S. Tretli, J. H. Olsen, and M. Fallah. 2015. 'Cancer Risk in Relatives of Testicular Cancer Patients by Histology Type and Age at Diagnosis: A Joint Study from Five Nordic Countries', *Eur Urol*, 68: 283-9.
- Koster, R., N. Mitra, K. D'Andrea, S. Vardhanabhuti, C. C. Chung, Z. Wang, R. Loren Erickson, D. J. Vaughn, K. Litchfield, N. Rahman, M. H. Greene, K. A. McGlynn, C. Turnbull, S. J. Chanock, K. L. Nathanson, and P. A. Kanetsky. 2014. 'Pathway-based analysis of GWAs data identifies association of sex determination genes with susceptibility to testicular germ cell tumors', *Hum Mol Genet*, 23: 6061-8.
- Kristiansen, W., R. Karlsson, T. B. Rounge, T. Whittington, B. K. Andreassen, P. K. Magnusson, S. D. Fossa, H. O. Adami, C. Turnbull, T. B. Haugen, T. Grotmol, and F. Wiklund. 2015. 'Two new loci and gene sets related to sex determination and cancer progression are associated with susceptibility to testicular germ cell tumor', *Hum Mol Genet*, 24: 4138-46.
- Ku, C. S., N. Naidoo, S. M. Teo, and Y. Pawitan. 2011. 'Regions of homozygosity and their impact on complex diseases and traits', *Hum Genet*, 129: 1-15.
- Le Cornet, C., J. Lortet-Tieulent, D. Forman, R. Beranger, A. Flechon, B. Fervers, J. Schuz, and F. Bray. 2014. 'Testicular cancer incidence to rise by 25% by 2025 in Europe? Model-based predictions in 40 countries using population-based registry data', *Eur J Cancer*, 50: 831-9.
- Lebel, R. R., and W. B. Gallagher. 1989. 'Wisconsin consanguinity studies. II: Familial adenocarcinomatosis', *Am J Med Genet*, 33: 1-6.
- Lencz, T., C. Lambert, P. DeRosse, K. E. Burdick, T. V. Morgan, J. M. Kane, R. Kucherlapati, and A. K. Malhotra. 2007. 'Runs of homozygosity reveal highly penetrant recessive loci in schizophrenia', *Proc Natl Acad Sci U S A*, 104: 19942-7.
- Li, M. J., L. Y. Wang, Z. Xia, M. P. Wong, P. C. Sham, and J. Wang. 2014. 'dbPSHP: a database of recent positive selection across human populations', *Nucleic Acids Res*, 42: D910-6.
- Litchfield, K, M. Levy, G. Orlando, P. Law, G. Migliorini, A. Holroyd, P. Broderick, J. Nsengimana, R. Eeles, D.F. Easton, D. Dudakia, D.T. Bishop, A. Reid, R.A. Huddart, J. Shipley, T. Grotmol, F.

- Wiklund, R.S. Houlston, and C. Turnbull. 2016. 'Identification and functional annotation of 19 novel loci reveals gene regulatory mechanisms determining susceptibility to testicular germ cell tumour', *Nat Genet*, (in press).
- Litchfield, K., A. Holroyd, A. Lloyd, P. Broderick, J. Nsengimana, R. Eeles, D. F. Easton, D. Dudakia, D. T. Bishop, A. Reid, R. A. Huddart, T. Grotmol, F. Wiklund, J. Shipley, R. S. Houlston, and C. Turnbull. 2015. 'Identification of four new susceptibility loci for testicular germ cell tumour', *Nat Commun*, 6: 8690.
- Litchfield, K., M. Levy, D. Dudakia, P. Proszek, C. Shipley, S. Basten, E. Rapley, D. T. Bishop, A. Reid, R. Huddart, P. Broderick, D. G. Castro, S. O'Connor, R. H. Giles, R. S. Houlston, and C. Turnbull. 2016. 'Rare disruptive mutations in ciliary function genes contribute to testicular cancer susceptibility', *Nat Commun*, 7: 13840.
- Litchfield, K., M. Levy, G. Orlando, C. Loveday, P. J. Law, G. Migliorini, A. Holroyd, P. Broderick, R. Karlsson, T. B. Haugen, W. Kristiansen, J. Nsengimana, K. Fenwick, I. Assiotis, Z. Kote-Jarai, A. M. Dunning, K. Muir, J. Peto, R. Eeles, D. F. Easton, D. Dudakia, N. Orr, N. Pashayan, D. T. Bishop, A. Reid, R. A. Huddart, J. Shipley, T. Grotmol, F. Wiklund, R. S. Houlston, and C. Turnbull. 2017. 'Identification of 19 new risk loci and potential regulatory mechanisms influencing susceptibility to testicular germ cell tumor', *Nat Genet*, 49: 1133-40.
- Litchfield, K., C. Loveday, M. Levy, D. Dudakia, E. Rapley, J. Nsengimana, D. T. Bishop, A. Reid, R. Huddart, P. Broderick, R. S. Houlston, and C. Turnbull. 2018. 'Large-scale Sequencing of Testicular Germ Cell Tumour (TGCT) Cases Excludes Major TGCT Predisposition Gene', *Eur Urol*, 73: 828-31.
- Litchfield, K., J. S. Mitchell, J. Shipley, R. Huddart, E. Rajpert-De Meyts, N. E. Skakkebaek, R. S. Houlston, and C. Turnbull. 2016. 'Polygenic susceptibility to testicular cancer: implications for personalised health care', *Br J Cancer*, 114: e22.
- Litchfield, K., J. Shipley, and C. Turnbull. 2015. 'Common variants identified in genome-wide association studies of testicular germ cell tumour: an update, biological insights and clinical application', *Andrology*, 3: 34-46.
- Litchfield, K., R. Sultana, A. Renwick, D. Dudakia, S. Seal, E. Ramsay, S. Powell, A. Elliott, M. Warren-Perry, R. Eeles, J. Peto, Z. Kote-Jarai, K. Muir, J. Nsengimana, M. R. Stratton, D. F. Easton, D. T. Bishop, R. A. Huddart, N. Rahman, and C. Turnbull. 2015. 'Multi-stage genome-wide association study identifies new susceptibility locus for testicular germ cell tumour on chromosome 3q25', *Hum Mol Genet*, 24: 1169-76.
- Litchfield, K., H. Thomsen, J. S. Mitchell, J. Sundquist, R. S. Houlston, K. Hemminki, and C. Turnbull. 2015. 'Quantifying the heritability of testicular germ cell tumour using both population-based and genomic approaches', *Sci Rep*, 5: 13889.
- Loveday, C., K. Litchfield, M. Levy, A. Holroyd, P. Broderick, Z. Kote-Jarai, A. M. Dunning, K. Muir, J. Peto, R. Eeles, D. F. Easton, D. Dudakia, N. Orr, N. Pashayan, A. Reid, R. A. Huddart, R. S. Houlston, and C. Turnbull. 2018. 'Validation of loci at 2q14.2 and 15q21.3 as risk factors for testicular cancer', *Oncotarget*, 9: 12630-38.
- McGlynn, K. A., S. S. Devesa, B. I. Graubard, and P. E. Castle. 2005. 'Increasing incidence of testicular germ cell tumors among black men in the United States', *J Clin Oncol*, 23: 5757-61.
- McQuillan, R., A. L. Leutenegger, R. Abdel-Rahman, C. S. Franklin, M. Pericic, L. Barac-Lauc, N. Smolej-Narancic, B. Janicijevic, O. Polasek, A. Tenesa, A. K. Macleod, S. M. Farrington, P. Rudan, C. Hayward, V. Vitart, I. Rudan, S. H. Wild, M. G. Dunlop, A. F. Wright, H. Campbell, and J. F. Wilson. 2008. 'Runs of homozygosity in European populations', *Am J Hum Genet*, 83: 359-72.
- Pemberton, T. J., D. Absher, M. W. Feldman, R. M. Myers, N. A. Rosenberg, and J. Z. Li. 2012. 'Genomic patterns of homozygosity in worldwide human populations', *Am J Hum Genet*, 91: 275-92.

- Purcell, S., B. Neale, K. Todd-Brown, L. Thomas, M. A. Ferreira, D. Bender, J. Maller, P. Sklar, P. I. de Bakker, M. J. Daly, and P. C. Sham. 2007. 'PLINK: a tool set for whole-genome association and population-based linkage analyses', *Am J Hum Genet*, 81: 559-75.
- Rapley, E. A., G. P. Crockford, D. F. Easton, M. R. Stratton, and D. T. Bishop. 2003. 'Localisation of susceptibility genes for familial testicular germ cell tumour', *Apmis*, 111: 128-33; discussion 33-5.
- Rapley, E. A., G. P. Crockford, D. Teare, P. Biggs, S. Seal, R. Barfoot, S. Edwards, R. Hamoudi, K. Heimdal, S. D. Fossa, K. Tucker, J. Donald, F. Collins, M. Friedlander, D. Hogg, P. Goss, A. Heidenreich, W. Ormiston, P. A. Daly, D. Forman, T. D. Oliver, M. Leahy, R. Huddart, C. S. Cooper, J. G. Bodmer, D. F. Easton, M. R. Stratton, and D. T. Bishop. 2000. 'Localization to Xq27 of a susceptibility gene for testicular germ-cell tumours', *Nat Genet*, 24: 197-200.
- Rapley, E. A., C. Turnbull, A. A. Al Olama, E. T. Dermitzakis, R. Linger, R. A. Huddart, A. Renwick, D. Hughes, S. Hines, S. Seal, J. Morrison, J. Nsengimana, P. Deloukas, U. K. Testicular Cancer Collaboration, N. Rahman, D. T. Bishop, D. F. Easton, and M. R. Stratton. 2009. 'A genome-wide association study of testicular germ cell tumor', *Nat Genet*, 41: 807-10.
- Ruark, E., S. Seal, H. McDonald, F. Zhang, A. Elliot, K. Lau, E. Perdeaux, E. Rapley, R. Eeles, J. Peto, Z. Kote-Jarai, K. Muir, J. Nsengimana, J. Shipley, U. K. Testicular Cancer Collaboration, D. T. Bishop, M. R. Stratton, D. F. Easton, R. A. Huddart, N. Rahman, and C. Turnbull. 2013. 'Identification of nine new susceptibility loci for testicular cancer, including variants near DAZL and PRDM14', *Nat Genet*, 45: 686-9.
- Schumacher, F. R., Z. Wang, R. I. Skotheim, R. Koster, C. C. Chung, M. A. Hildebrandt, C. P. Kratz, A. C. Bakken, D. T. Bishop, M. B. Cook, R. L. Erickson, S. D. Fossa, M. H. Greene, K. B. Jacobs, P. A. Kanetsky, L. N. Kolonel, J. T. Loud, L. A. Korde, L. Le Marchand, J. P. Lewinger, R. A. Lothe, M. C. Pike, N. Rahman, M. V. Rubertone, S. M. Schwartz, K. D. Siegmund, E. C. Skinner, C. Turnbull, D. J. Van Den Berg, X. Wu, M. Yeager, K. L. Nathanson, S. J. Chanock, V. K. Cortessis, and K. A. McGlynn. 2013. 'Testicular germ cell tumor susceptibility associated with the UCK2 locus on chromosome 1q23', *Hum Mol Genet*, 22: 2748-53.
- Shami, S. A., R. Qaisar, and A. H. Bittles. 1991. 'Consanguinity and adult morbidity in Pakistan', *Lancet*, 338: 954.
- Simpson, J. L., A. O. Martin, S. Elias, G. E. Sarto, and J. K. Dunn. 1981. 'Cancers of the breast and female genital system: search for recessive genetic factors through analysis of human isolate', *Am J Obstet Gynecol*, 141: 629-36.
- Sud, A., R. Cooke, A. J. Swerdlow, and R. S. Houlston. 2015. 'Genome-wide homozygosity signature and risk of Hodgkin lymphoma', *Sci Rep*, 5: 14315.
- Sud, A., B. Kinnersley, and R. S. Houlston. 2017. 'Genome-wide association studies of cancer: current insights and future perspectives', *Nat Rev Cancer*.
- Swerdlow, A. J., B. L. De Stavola, M. A. Swanwick, and N. E. Maconochie. 1997. 'Risks of breast and testicular cancers in young adult twins in England and Wales: evidence on prenatal and genetic aetiology', *Lancet*, 350: 1723-8.
- Szpiech, Z. A., J. Xu, T. J. Pemberton, W. Peng, S. Zollner, N. A. Rosenberg, and J. Z. Li. 2013. 'Long runs of homozygosity are enriched for deleterious variation', *Am J Hum Genet*, 93: 90-102.
- Tajima, F. 1989. 'Statistical method for testing the neutral mutation hypothesis by DNA polymorphism', *Genetics*, 123: 585-95.
- Thomsen, H., B. Chen, G. Figlioli, R. Elisei, C. Romei, M. Cipollini, A. Cristaudo, F. Bambi, P. Hoffmann, S. Herms, S. Landi, K. Hemminki, F. Gemignani, and A. Forsti. 2016. 'Runs of homozygosity and inbreeding in thyroid cancer', *BMC Cancer*, 16: 227.
- Thomsen, H., M. I. Filho, A. Woltmann, R. Johansson, J. E. Eyfjord, U. Hamann, J. Manjer, K. Enquist-Olsson, R. Henriksson, S. Herms, P. Hoffmann, B. Chen, S. Huhn, K. Hemminki, P. Lenner, and A. Forsti. 2015. 'Inbreeding and homozygosity in breast cancer survival', *Sci Rep*, 5: 16467.
- Turnbull, C., and N. Rahman. 2011. 'Genome-wide association studies provide new insights into the genetic basis of testicular germ-cell tumour', *Int J Androl*, 34: e86-96; discussion e96-7.

- Turnbull, C., E. A. Rapley, S. Seal, D. Pernet, A. Renwick, D. Hughes, M. Ricketts, R. Linger, J. Nsengimana, P. Deloukas, R. A. Huddart, D. T. Bishop, D. F. Easton, M. R. Stratton, N. Rahman, and U. K. Testicular Cancer Collaboration. 2010. 'Variants near DMRT1, TERT and ATF7IP are associated with testicular germ cell cancer', *Nat Genet*, 42: 604-7.
- Voight, B. F., S. Kudaravalli, X. Wen, and J. K. Pritchard. 2006. 'A map of recent positive selection in the human genome', *PLoS Biol*, 4: e72.
- Wang, E. T., G. Kodama, P. Baldi, and R. K. Moyzis. 2006. 'Global landscape of recent inferred Darwinian selection for Homo sapiens', *Proc Natl Acad Sci U S A*, 103: 135-40.
- Wang, Z., K. A. McGlynn, E. Rajpert-De Meyts, D. T. Bishop, C. C. Chung, M. D. Dalgaard, M. H. Greene, R. Gupta, T. Grotmol, T. B. Haugen, R. Karlsson, K. Litchfield, N. Mitra, K. Nielsen, L. C. Pyle, S. M. Schwartz, V. Thorsson, S. Vardhanabhuti, F. Wiklund, C. Turnbull, S. J. Chanock, P. A. Kanetsky, and K. L. Nathanson. 2017. 'Meta-analysis of five genome-wide association studies identifies multiple new loci associated with testicular germ cell tumor', *Nat Genet*, 49: 1141-47.
- Williamson, S. H., M. J. Hubisz, A. G. Clark, B. A. Payseur, C. D. Bustamante, and R. Nielsen. 2007. 'Localizing recent adaptive evolution in the human genome', *PLoS Genet*, 3: e90.

Figures




Fig. 1. Cumulative distribution of runs of homozygosity (RoH) in TGCT cases and controls. Data is presented in such a way that each data point represents the cumulative fraction (y-axis) of the samples with the corresponding minimum total length of the genome covered by RoH (x-axis), as determined from PLINK.




Fig. 2. Frequency of consensus runs of homozygosity (RoH) in TGCT cases versus controls. Consensus RoH were defined on the basis of being present in 10 or more individuals.

Tables

Table 1. Global Homozygosity Measures in TGCT Cases versus Controls

Measure	TGCT Cases (n=3206)	Controls (n=7442)	P
Any size			
Average number of RoH per individual	12.9	12.9	0.8
Average length per RoH per individual (Kb)	1,628.8	1,633.4	0.4
Average total length of RoH per individual (Kb)	21,346.9	21,159.5	0.2
< 1.6 Mb			
Average number of RoH per individual	8.4	8.3	0.7
Average length per RoH per individual (Kb)	1,256.5	1,255.9	0.8
Average total length of RoH per individual (Kb)	10,517.8	10,491.4	0.2
> 1.6 Mb			
Average number of RoH per individual	4.7	4.6	0.7
Average length per RoH per individual (Kb)	2,299.1	2,302.4	0.7
Average total length of RoH per individual (Kb)	10,945.2	10,781.3	0.7

RoH, runs of homozygosity. *P* calculated using Student's t-test.

Table 2. Consensus RoH with frequency > 25% in controls

RoH ID	Chr	Start (b37)	End	Length (Kb)	No. SNPs	Controls	Cases	P	Centromeric	iHS _{max}	Tajima D _{max}	Fst _{max}
ROH164	8	29737732	70143771	40,406.04	3286	34.6%	34.8%	0.9	Yes	3.16	4.80	0.30
ROH68	3	74113968	116444174	42,330.21	4272	34.0%	32.4%	0.1	Yes	3.27	5.02	0.25
ROH117	6	23594993	40225964	16,630.97	7636	32.4%	34.0%	0.1	No	3.17	4.80	0.27
ROH203	11	44966113	69074890	24,108.78	2174	30.6%	30.4%	0.9	Yes	3.91	4.48	0.24
ROH43	2	132755417	169516830	36,761.41	3352	27.4%	28.2%	0.4	No	3.24	4.99	0.46
ROH86	4	58139970	111262618	53,122.65	5794	26.0%	26.7%	0.5	No	3.25	5.10	0.33
ROH120	6	53377834	91107018	37,729.18	3378	25.8%	26.1%	0.8	Yes	3.31	4.67	0.26

RoH, runs of homozygosity. Chr, chromosome. iHS, D and Fst metrics were obtained from dbPSHP. P was calculated using Fisher's exact.

Table 3. Consensus RoH putatively associated with TGCT risk (P<0.01)

RoH ID	Chr	Start (b37)	End	Length (Kb)	No. SNPs	Controls	Cases	P	iHS _{max}	Tajima D _{max}	Fst _{max}
ROH200	11	24794324	33879547	9,085,223	866	3.3%	4.6%	0.001	4.05	4.91	0.25
ROH101	5	79191702	115939896	36,748,194	3520	22.2%	19.7%	0.005	3.46	4.62	0.31
ROH229	13	44234831	46712448	2,477,617	272	0.5%	0.1%	0.009	2.43	4.70	0.28

RoH, runs of homozygosity. Chr, chromosome. iHS, D and Fst metrics were obtained from dbPSHP. P was calculated using Fisher's exact.

Consortia

The UK Testicular Cancer Collaboration (UKTCC)

Gordon Rustin¹⁶, Narayanan N Srihari¹⁷, David Cole¹⁸, Colin Askill¹⁹, Gianfilippo Bertelli¹⁹, James Barber²⁰, Ed Gilby²¹, Jeff White²², Jeremy Baybrooke²³, Michael Leahy²⁴, Richard Welch²⁴, Prabir Chakraborti²⁵, Johnathan Joffe²⁶, Richard Brown²⁷, Guy Faust^{28,29}, Peter Simmonds³⁰, Danish Mazhar³¹, Andrew Stockdale³², David Hrounda³², Caroline Humber³², Wiebke Appel³³, Anne Hong³⁴, Grahame Howard³⁵, Fiona Douglas³⁶, David Bloomfield³⁷, Mohammad Butt³⁸, Kay Kelly³⁹, Rakesh Mehra⁴⁰, Richard Brown⁴¹, Paul Rogers⁴¹, Prabir Chakraborti⁴², Matthew Hatton⁴³, Ivo Hennig⁴⁴, John McAteer⁴⁵, Philip Savage⁴⁶, Michael Seckl⁴⁶, Joanna Gale⁴⁷, Gordon Rustin⁴⁸, Peter Clark⁴⁹, Steve Woby⁵⁰, Adrian Rathmell⁵¹, Alan Lamont⁵², Naveed Sarwar⁵³, Nick Stuart⁵⁴, Simon Chowdhury⁸, Sharon Beesley⁵⁵, Mathius Winkler⁵⁶, Abdel Hamid⁵⁷, Sanjeev Pathak⁵⁸, Krishnaswamy Madhavan⁵⁹, Martin Highley⁶⁰, Julian Money-Kryle⁶¹, Cathryn Brock⁶², Thiagarajan Sreenivasan^{63,64}.

¹⁶Mount Vernon Cancer Centre, Mount Vernon Hospital, Northwood, UK.

¹⁷Trials Unit, Royal Shrewsbury Hospital, Shrewsbury, UK.

¹⁸Osprey Unit, Great Western Hospital, Swindon, UK.

¹⁹South West Wales Cancer Institute, Singleton Hospital and Morrision Hospital, Sketty, Swansea, UK.

²⁰Clinical Trials Unit, Velindre Hospital, Velindre Cancer Centre, Whitchurch, Cardiff, UK.

²¹Department of Oncology and Haematology, Royal United Hospital, Combe Park, Bath, UK.

²²Beatson Oncology Centre, Beatson West of Scotland Cancer Centre, Glasgow, UK.

²³Bristol Haematology & Oncology Centre, United Bristol Healthcare NHS Foundation Trust, Bristol, UK.

²⁴Christie Hospital, Withington, Manchester, UK.

²⁵Derbyshire Royal Infirmary, Derby Teaching Hospitals NHS Foundation Trust, Derby, UK.

²⁶Department of Medical Oncology, St James Hospital Leeds, Leeds, UK.

²⁷Cancer Clinical Trials, Wexham Park Hospital, John Ulster Post Grad Centre, Slough, UK.

²⁸LNR Cancer Reseach Network, Leicester Royal Infirmary, Leicester, UK.

²⁹Northampton General, Cliftonville, Northampton, UK.

³⁰Cancer Care Directorate, Medical Oncology, Southampton General Hospital, Southampton, UK.

³¹Addenbrookes Hospital, Cambridge Clinical Trials Centre, Oncology Clinical Trials, Cambridge, UK.

³²Arden Cancer Centre, University Hospital Walsgrave, West Wing, Coventry, UK.

³³Department of Oncology, Royal Preston Hospital, Fulwood, Preston, UK.

³⁴Exeter Oncology Centre, Royal Devon and Exeter Hospital, Exeter, UK.

³⁵Scottish Cancer Research Network, Oncology Admin Corridor, Edinburgh Cancer Centre, Western General Hospital, Edinburgh, UK.

³⁶Clinical Trials Unit, Newcastle General Hospital, Newcastle-upon Tyne, UK.

³⁷The Sussex Cancer Centre, The Royal Sussex County Hospital, Brighton, UK.

³⁸Castle Hill Hospital, Cottingham, UK.

³⁹Raigmore Hospital, NHS Highland, Inverness, UK.

⁴⁰Greater Midlands Cancer Research Network, The Chestnuts, The Royal Wolverhampton Hospitals NHS Trust, Wolverhampton, UK.

⁴¹Berkshire Cancer Centre, Royal Berkshire Hospital, Reading, UK.

⁴²Queens Hospital, Burton upon Trent, Burton, UK.

⁴³Sheffield Teaching Hospitals NHS Foundation Trust, Sheffield, UK.

⁴⁴Nottingham University Hospitals NHS Trust, City Hospital Campus, Nottingham, UK.

⁴⁵Northern Ireland Cancer Centre, Belfast City Hospital, Belfast, UK.

⁴⁶Department of Medical Oncology, Charing Cross Hospital, London, UK.

⁴⁷Level B Queen Alexandra Hospital, Cosham, Portsmouth, UK.

⁴⁸R&D Office, Education Centre, Hillingdon Hospital, Hillingdon, UK.

⁴⁹Royal Liverpool & Broadgreen University Hospitals NHS Trust, Liverpool, UK.

⁵⁰Royal Oldham Hospital/Pennine Acute Hospital, Oldham, UK.

⁵¹James Cook University Hospital, South Tees Hospitals NHS Foundation Trust, Middlesbrough, UK.

⁵²Essex County Hospital, Colchester Hospital University NHS Foundation Trust, Colchester, UK.

⁵³Basildon University Hospital, Basildon and Thurrock Hospitals NHS Foundation Trust, Nethermayne, Basildon, UK.

⁵⁴North Wales Cancer Treatment Centre, Glan Clwyd Hospital, Rhyl, UK.

⁵⁵Maidstone Hospital, Maidstone and Tunbridge Wells NHS Trust, Barming, Maidstone, UK.

⁵⁶R&D Department, West Middlesex University Hospital NHS Trust, Isleworth, UK.

⁵⁷ West Wing 2, Broomfield Hospital, Broomfield, Chelmsford, UK.

⁵⁸ Joint Research Office of Doncaster and Bassetlaw Hospitals NHS Foundation Trust, Doncaster Royal Infirmary, Doncaster, UK.

⁵⁹ Southend University Hospital, Westcliff-On-Sea, UK.

⁶⁰ Derriford Hospital (Plymouth), Plymouth Hospitals NHS Trust, Derriford Hospital, Plymouth, UK.

⁶¹ St Lukes Cancer Centre, Royal Surrey County Hospital, Guildford, UK.

⁶² Unit 101, Chelsea & Westminster Hospital, Harbour Yard, Chelsea Harbour, London, UK.

⁶³ Lincoln County Hospital, United Lincolnshire Hospitals NHS Trust, Lincoln, UK.

⁶⁴ Pilgrim Hospital, United Lincolnshire Hospitals NHS Trust, Boston, UK.

The PRACTICAL Consortium

Brian E. Henderson⁶⁵, Christopher A. Haiman⁶⁵, Fredrick R. Schumacher^{66,67}, Ali Amin Al Olama^{8,68}, Sara Benlloch^{8,1}, Sonja I. Berndt⁶⁹, David V. Conti⁶⁵, Fredrik Wiklund⁷⁰, Stephen Chanock⁶⁹, Susan Gapster⁷¹, Victoria L. Stevens⁷¹, Catherine M. Tangen⁷², Jyotsna Batra^{73,74}, Judith Clements^{73,74}, Australian Prostate Cancer Research Centre BioResource (APCB)⁷³, Henrik Gronberg⁷⁰, Johanna Schleutker^{75,76}, Demetrius Albanes⁶⁹, Alicja Wolk⁷⁷, Catharine West⁷⁸, Lorelei Mucci⁷⁹, Géraldine Cancel-Tassin^{80,81}, Stella Koutros⁶⁹, Karina Dalsgaard Sorensen^{82,83}, Lovise Maehle⁸⁴, David E. Neal^{85,86,87}, Freddie C. Hamdy⁸⁷, Jenny L. Donovan⁸⁸, Ruth C. Travis⁸⁹, Robert J. Hamilton⁹⁰, Sue Ann Ingles⁶⁵, Barry S. Rosenstein^{91,92}, Yong-Jie Lu⁹³, Graham G. Giles^{94,95}, Adam S. Kibel⁹⁶, Ana Vega⁹⁷, Manolis Kogevinas^{98,99,100,101}, Kathryn L. Penney¹⁰², Jong Y. Park¹⁰³, Janet L. Stanford^{104,105}, Cezary Cybulski¹⁰⁶, Børge G. Nordestgaard^{107,108}, Hermann Brenner^{109,110,111}, Christiane Maier¹¹², Jeri Kim¹¹³, Esther M.

John^{114,115}, Manuel R. Teixeira^{116,117}, Susan L. Neuhausen¹¹⁸, Kim De Ruyck¹¹⁹, Azad Razack¹²⁰, Lisa F. Newcomb^{104,121}, Davor Lessel¹²², Radka Kaneva¹²³, Nawaid Usmani^{124,125}, Frank Claessens¹²⁶, Paul A. Townsend¹²⁷, Manuela Gago Dominguez^{128,129}, Monique J. Roobol¹³⁰, Florence Menegaux¹³¹, Kay-Tee Khaw¹³², Lisa Cannon-Albrigh^{133,134}, Hardev Pandha¹³⁵, Stephen N. Thibodeau¹³⁶.

⁶⁵Department of Preventive Medicine, Keck School of Medicine, University of Southern California/Norris Comprehensive Cancer Center, Los Angeles, CA, USA.

⁶⁶Department of Epidemiology and Biostatistics, Case Western Reserve University, Cleveland, OH, USA.

⁶⁷Seidman Cancer Center, University Hospitals, Cleveland, OH, USA.

⁶⁸University of Cambridge, Department of Clinical Neurosciences, Cambridge, UK.

⁶⁹Division of Cancer Epidemiology and Genetics, National Cancer Institute, NIH, Bethesda, MD, USA.

⁷⁰Department of Medical Epidemiology and Biostatistics, Karolinska Institute, Stockholm, Sweden.

⁷¹Epidemiology Research Program, American Cancer Society, 250 Williams Street, Atlanta, GA, USA.

⁷²SWOG Statistical Center, Fred Hutchinson Cancer Research Center, Seattle, WA, USA.

⁷³Australian Prostate Cancer Research Centre-Qld, Institute of Health and Biomedical Innovation and School of Biomedical Science, Queensland University of Technology, Brisbane, Queensland, Australia.

⁷⁴Translational Research Institute, Brisbane, Queensland, Australia.

⁷⁵Department of Medical Biochemistry and Genetics, Institute of Biomedicine, University of Turku, Finland.

⁷⁶Tyks Microbiology and Genetics, Department of Medical Genetics, Turku University Hospital, Finland.

⁷⁷Division of Nutritional Epidemiology, Institute of Environmental Medicine, Karolinska Institutet, Sweden.

⁷⁸Division of Cancer Sciences, University of Manchester, Manchester Academic Health Science Centre, Radiotherapy Related Research, Manchester NIHR Biomedical Research Centre, The Christie Hospital NHS Foundation Trust, Manchester, UK.

⁷⁹Department of Epidemiology, Harvard T.H Chan School of Public Health, Boston, MA, USA.

⁸⁰CeRePP, Tenon Hospital, Paris, France.

⁸¹UPMC Sorbonne Universites, GRC N°5 ONCOTYPE-URO, Tenon Hospital, Paris, France.

⁸²Department of Molecular Medicine, Aarhus University Hospital, Denmark.

⁸³Department of Clinical Medicine, Aarhus University, Denmark.

⁸⁴Department of Medical Genetics, Oslo University Hospital, Norway.

⁸⁵University of Cambridge, Department of Oncology, Addenbrooke's Hospital, Cambridge, UK.

⁸⁶Cancer Research UK Cambridge Research Institute, Li Ka Shing Centre, Cambridge, UK.

⁸⁷Nuffield Department of Surgical Sciences, University of Oxford, Oxford, UK, Faculty of Medical Science, University of Oxford, John Radcliffe Hospital, Oxford, UK.

⁸⁸School of Social and Community Medicine, University of Bristol, Bristol, UK.

⁸⁹Cancer Epidemiology Unit, Nuffield Department of Population Health University of Oxford, Oxford, UK.

⁹⁰Dept. of Surgical Oncology, Princess Margaret Cancer Centre, Toronto, Canada.

⁹¹Department of Radiation Oncology, Icahn School of Medicine at Mount Sinai, New York, NY, USA.

⁹²Department of Genetics and Genomic Sciences, Icahn School of Medicine at Mount Sinai, New York, NY, USA.

⁹³Centre for Molecular Oncology, Barts Cancer Institute, Queen Mary University of London, John Vane Science Centre, London, UK.

⁹⁴Cancer Epidemiology & Intelligence Division, The Cancer Council Victoria, Melbourne, Victoria, Australia.

- ⁹⁵Centre for Epidemiology and Biostatistics, Melbourne School of Population and Global Health, The University of Melbourne, Melbourne, Australia.
- ⁹⁶Division of Urologic Surgery, Brigham and Womens Hospital, Boston, MA, USA.
- ⁹⁷Fundación Pública Galega de Medicina Xenómica-SERGAS, Grupo de Medicina Xenómica, CIBERER, IDIS, Santiago de Compostela, Spain.
- ⁹⁸Centre for Research in Environmental Epidemiology (CREAL), Barcelona Institute for Global Health (ISGlobal), Barcelona, Spain.
- ⁹⁹CIBER Epidemiología y Salud Pública (CIBERESP), Madrid, Spain.
- ¹⁰⁰IMIM (Hospital del Mar Research Institute), Barcelona, Spain.
- ¹⁰¹Universitat Pompeu Fabra (UPF), Barcelona, Spain.
- ¹⁰²Channing Division of Network Medicine, Department of Medicine, Brigham and Women's Hospital/Harvard Medical School, Boston, MA, USA.
- ¹⁰³Department of Cancer Epidemiology, Moffitt Cancer Center, Tampa, USA.
- ¹⁰⁴Division of Public Health Sciences, Fred Hutchinson Cancer Research Center, Seattle, Washington, USA.
- ¹⁰⁵Department of Epidemiology, School of Public Health, University of Washington, Seattle, Washington, USA.
- ¹⁰⁶International Hereditary Cancer Center, Department of Genetics and Pathology, Pomeranian Medical University, Szczecin, Poland.
- ¹⁰⁷Faculty of Health and Medical Sciences, University of Copenhagen, Denmark.
- ¹⁰⁸Department of Clinical Biochemistry, Herlev and Gentofte Hospital, Copenhagen University Hospital, Herlev, Denmark.
- ¹⁰⁹Division of Clinical Epidemiology and Aging Research, German Cancer Research Center (DKFZ), Heidelberg, Germany.
- ¹¹⁰German Cancer Consortium (DKTK), German Cancer Research Center (DKFZ), Heidelberg, Germany.
- ¹¹¹Division of Preventive Oncology, German Cancer Research Center (DKFZ) and National Center for Tumor Diseases (NCT), Heidelberg, Germany.
- ¹¹²Institute for Human Genetics, University Hospital Ulm, Ulm, Germany.

- ¹¹³The University of Texas MD Anderson Cancer Center, Department of Genitourinary Medical Oncology, Houston, TX, USA.
- ¹¹⁴Cancer Prevention Institute of California, Fremont, CA, USA.
- ¹¹⁵Department of Health Research & Policy (Epidemiology) and Stanford Cancer Institute, Stanford University School of Medicine, Stanford, CA , USA.
- ¹¹⁶Department of Genetics, Portuguese Oncology Institute of Porto, Porto, Portugal.
- ¹¹⁷Biomedical Sciences Institute (ICBAS), University of Porto, Porto, Portugal.
- ¹¹⁸Department of Population Sciences, Beckman Research Institute of the City of Hope, Duarte, CA, USA.
- ¹¹⁹Ghent University, Faculty of Medicine and Health Sciences, Basic Medical Sciences, Gent, Belgium.
- ¹²⁰Department of Surgery, Faculty of Medicine, University of Malaya, Kuala Lumpur, Malaysia.
- ¹²¹Department of Urology, University of Washington, Seattle, WA, USA.
- ¹²²Institute of Human Genetics, University Medical Center Hamburg-Eppendorf, Hamburg, Germany.
- ¹²³Molecular Medicine Center, Department of Medical Chemistry and Biochemistry, Medical University, Sofia, Bulgaria.
- ¹²⁴Department of Oncology, Cross Cancer Institute, University of Alberta, Edmonton, Alberta, Canada.
- ¹²⁵Division of Radiation Oncology, Cross Cancer Institute, Edmonton, Alberta, Canada.
- ¹²⁶Molecular Endocrinology Laboratory, Department of Cellular and Molecular Medicine, KU Leuven, Leuven, Belgium.
- ¹²⁷Institute of Cancer Sciences, Manchester Cancer Research Centre, University of Manchester, Manchester Academic Health Science Centre, St Mary's Hospital, Manchester, UK.
- ¹²⁸Genomic Medicine Group, Galician Foundation of Genomic Medicine, Instituto de Investigacion Sanitaria de Santiago de Compostela (IDIS), Complejo Hospitalario Universitario de Santiago, Servicio Galego de Saúde, SERGAS, Santiago De Compostela, Spain.
- ¹²⁹University of California San Diego, Moores Cancer Center, La Jolla, CA, USA.

¹³⁰Department of Urology, Erasmus University Medical Center, Rotterdam, the Netherlands.

¹³¹Cancer & Environment Group, Center for Research in Epidemiology and Population Health (CESP), INSERM, University Paris-Sud, University Paris-Saclay, Villejuif, France.

¹³²Clinical Gerontology Unit, University of Cambridge, Cambridge, UK.

¹³³Division of Genetic Epidemiology, Department of Medicine, University of Utah School of Medicine, Salt Lake City, Utah, USA.

¹³⁴George E. Wahlen Department of Veterans Affairs Medical Center, Salt Lake City, UT, USA.

¹³⁵The University of Surrey, Guildford, Surrey, UK.

¹³⁶Department of Laboratory Medicine and Pathology, Mayo Clinic, Rochester, MN, USA.