

## Mendelian randomization study of height and body mass index as modifiers of ovarian cancer risk in 22,588 *BRCA1* and *BRCA2* mutation carriers

Frank Qian<sup>1</sup>, Matti A. Rookus<sup>2</sup>, Goska Leslie<sup>3</sup>, Harvey A. Risch<sup>4</sup>, Mark H. Greene<sup>5</sup>, Cora M. Aalfs<sup>6</sup>, Muriel A. Adank<sup>7</sup>, Julian Adlard<sup>8</sup>, Bjarni A. Agnarsson<sup>9,10</sup>, Munaza Ahmed<sup>11</sup>, Kristiina Aittomäki<sup>12</sup>, Irene L. Andrulis<sup>13, 14</sup>, Norbert Arnold<sup>15</sup>, Banu K. Arun<sup>16</sup>, Margreet G.E.M. Ausems<sup>17</sup>, Jacopo Azzollini<sup>18</sup>, Daniel Barrowdale<sup>3</sup>, Julian Barwell<sup>19</sup>, Javier Benitez<sup>20, 21</sup>, Katarzyna Białkowska<sup>22</sup>, Valérie Bonadona<sup>23</sup>, Julika Borde<sup>24-26</sup>, Ake Borg<sup>27</sup>, Angela R. Bradbury<sup>28</sup>, Joan Brunet<sup>29</sup>, Saundra S. Buys<sup>30</sup>, Trinidad Caldés<sup>31</sup>, Maria A. Caligo<sup>32</sup>, Ian Campbell<sup>33, 34</sup>, Jonathan Carter<sup>35</sup>, Jocelyne Chiquette<sup>36</sup>, Wendy K. Chung<sup>37</sup>, Kathleen B.M. Claes<sup>38</sup>, J. Margriet Collée<sup>39</sup>, Marie-Agnès Collonge-Rame<sup>40</sup>, Fergus J. Couch<sup>41</sup>, Mary B. Daly<sup>42</sup>, Capucine Delnatte<sup>43</sup>, Orland Diez<sup>44</sup>, Susan M. Domchek<sup>28</sup>, Cecilia M. Dorfling<sup>45</sup>, Jacqueline Eason<sup>46</sup>, Douglas F. Easton<sup>3, 47</sup>, Ros Eeles<sup>48</sup>, Christoph Engel<sup>49, 50</sup>, D. Gareth Evans<sup>51, 52</sup>, Laurence Faivre<sup>53, 54</sup>, Lidia Feliubadaló<sup>55</sup>, Lenka Foretova<sup>56</sup>, Eitan Friedman<sup>57, 58</sup>, Debra Frost<sup>3</sup>, Patricia A. Ganz<sup>59</sup>, Judy Garber<sup>60</sup>, Vanesa Garcia-Barberan<sup>31</sup>, Andrea Gehrig<sup>61</sup>, Gord Glendon<sup>13</sup>, Andrew K. Godwin<sup>62</sup>, Encarna B. Gómez Garcia<sup>63</sup>, Ute Hamann<sup>64</sup>, Jan Hauke<sup>24-26</sup>, John L. Hopper<sup>65</sup>, Peter J. Hulick<sup>66, 67</sup>, Evgeny N. Imyanitov<sup>68</sup>, Claudine Isaacs<sup>69</sup>, Louise Izatt<sup>70</sup>, Anna Jakubowska<sup>22, 71</sup>, Ramunas Janavicius<sup>72</sup>, Esther M. John<sup>73</sup>, Beth Y. Karlan<sup>74</sup>, Carolien M. Kets<sup>75</sup>, Yael Laitman<sup>57</sup>, Conxi Lázaro<sup>55</sup>, Dominique Leroux<sup>76</sup>, Jenny Lester<sup>74</sup>, Fabienne Lesueur<sup>77-79</sup>, Jennifer T. Loud<sup>5</sup>, Jan Lubiński<sup>22</sup>, Alicja Łukomska<sup>22</sup>, Lesley McGuffog<sup>3</sup>, Noura Mebirouk<sup>77-79</sup>, Hanne E.J. Meijers-Heijboer<sup>80</sup>, Alfons Meindl<sup>81</sup>, Austin Miller<sup>82</sup>, Marco Montagna<sup>83</sup>, Thea M. Mooij<sup>2</sup>, Emmanuelle Mouret-Fourme<sup>84</sup>, Katherine L. Nathanson<sup>28</sup>, Bitu Nehoray<sup>85</sup>, Susan L. Neuhausen<sup>86</sup>, Heli Nevanlinna<sup>87</sup>, Finn C. Nielsen<sup>88</sup>, Kenneth Offit<sup>89, 90</sup>, Edith Olah<sup>91</sup>, Kai-ren Ong<sup>92</sup>, Jan C. Oosterwijk<sup>93</sup>, Laura Ottini<sup>94</sup>, Michael T. Parsons<sup>95</sup>, Paolo Peterlongo<sup>96</sup>, Georg Pfeiler<sup>97</sup>, Nisha Pradhan<sup>89</sup>, Paolo Radice<sup>98</sup>, Susan J. Ramus<sup>99, 100</sup>, Johanna Rantala<sup>101</sup>, Gad Rennert<sup>102</sup>, Mark Robson<sup>90</sup>, Gustavo C. Rodriguez<sup>103</sup>, Ritu Salani<sup>104</sup>, Maren T. Scheuner<sup>105</sup>, Rita K. Schmutzler<sup>25, 26</sup>, Payal D. Shah<sup>28</sup>, Lucy E. Side<sup>11</sup>, Jacques Simard<sup>106</sup>, Christian F. Singer<sup>97</sup>, Doris Steinemann<sup>107</sup>, Dominique Stoppa-Lyonnet<sup>84, 108, 109</sup>, Yen Yen Tan<sup>97</sup>, Manuel R. Teixeira<sup>110, 111</sup>, Mary Beth Terry<sup>112</sup>, Mads Thomassen<sup>113</sup>, Marc Tischkowitz<sup>114, 115</sup>, Silvia Tognazzo<sup>83</sup>, Amanda E. Toland<sup>116</sup>, Nadine Tung<sup>117</sup>, Christi J. van Asperen<sup>118</sup>, Klaartje van Engelen<sup>119</sup>, Elizabeth J. van Rensburg<sup>45</sup>, Laurence Venat-Bouvet<sup>120</sup>, Jeroen Vierstraete<sup>38</sup>, Gabriel Wagner<sup>97</sup>, Lisa Walker<sup>121</sup>, Jeffrey N. Weitzel<sup>85</sup>, Drakoulis Yannoukakos<sup>122</sup>, KConFab Investigators<sup>33, 34</sup>, HEBON Investigators<sup>123</sup>, GEMO Study Collaborators<sup>84</sup>, EMBRACE Collaborators<sup>3</sup>, Antonis C. Antoniou<sup>3</sup>, David E. Goldgar<sup>124</sup>, Olufunmilayo I. Olopade<sup>125</sup>, Georgia Chenevix-Trench<sup>95</sup>, Timothy R. Rebbeck<sup>126, 127</sup>, Dezheng Huo<sup>125, 128\*</sup>, on behalf of CIMBA

<sup>1</sup> Department of Medicine, University of Chicago, Chicago, IL, USA.

<sup>2</sup> Department of Epidemiology, The Netherlands Cancer Institute, Amsterdam, The Netherlands.

<sup>3</sup> Centre for Cancer Genetic Epidemiology, Department of Public Health and Primary Care, University of Cambridge, Cambridge, UK.

<sup>4</sup> Chronic Disease Epidemiology, Yale School of Public Health, New Haven, CT, USA.

- <sup>5</sup> Clinical Genetics Branch, Division of Cancer Epidemiology and Genetics, National Cancer Institute, Bethesda, MD, USA.
- <sup>6</sup> Department of Clinical Genetics, Amsterdam UMC, location AMC, Amsterdam, The Netherlands.
- <sup>7</sup> Family Cancer Clinic, The Netherlands Cancer Institute - Antoni van Leeuwenhoek hospital, Amsterdam, The Netherlands.
- <sup>8</sup> Yorkshire Regional Genetics Service, Chapel Allerton Hospital, Leeds, UK.
- <sup>9</sup> Department of Pathology, Landspítali University Hospital, Reykjavik, Iceland.
- <sup>10</sup> School of Medicine, University of Iceland, Reykjavik, Iceland.
- <sup>11</sup> Great Ormond Street Hospital for Children, London, UK.
- <sup>12</sup> Department of Clinical Genetics, Helsinki University Hospital, University of Helsinki, Helsinki, Finland.
- <sup>13</sup> Fred A. Litwin Center for Cancer Genetics, Lunenfeld-Tanenbaum Research Institute of Mount Sinai Hospital, Toronto, ON, Canada.
- <sup>14</sup> Department of Molecular Genetics, University of Toronto, Toronto, ON, Canada.
- <sup>15</sup> Department of Gynaecology and Obstetrics, University Hospital of Schleswig-Holstein, Campus Kiel, Christian-Albrechts University Kiel, Kiel, Germany.
- <sup>16</sup> Department of Breast Medical Oncology, University of Texas MD Anderson Cancer Center, Houston, TX, USA.
- <sup>17</sup> Division Laboratories, Pharmacy and Biomedical Genetics, Department of Medical Genetics, University Medical Center Utrecht, Utrecht, The Netherlands.
- <sup>18</sup> Unit of Medical Genetics, Department of Medical Oncology and Hematology, Fondazione IRCCS (Istituto di Ricovero e Cura a Carattere Scientifico) Istituto Nazionale dei Tumori di Milano, Milan, Italy.
- <sup>19</sup> Leicestershire Clinical Genetics Service, University Hospitals of Leicester NHS Trust, Leicester, UK.
- <sup>20</sup> Human Cancer Genetics Programme, Spanish National Cancer Research Centre (CNIO), Madrid, Spain.
- <sup>21</sup> Biomedical Network on Rare Diseases (CIBERER), Madrid, Spain.
- <sup>22</sup> Department of Genetics and Pathology, Pomeranian Medical University, Szczecin, Poland.
- <sup>23</sup> Unité de Prévention et d'Epidémiologie Génétique, Centre Léon Bérard, Lyon, France.
- <sup>24</sup> Center for Integrated Oncology (CIO), University Hospital of Cologne, Cologne, Germany.
- <sup>25</sup> Center for Molecular Medicine Cologne (CMMC), University of Cologne, Cologne, Germany.
- <sup>26</sup> Center for Hereditary Breast and Ovarian Cancer, University Hospital of Cologne, Cologne, Germany.
- <sup>27</sup> Department of Oncology, Lund University and Skåne University Hospital, Lund, Sweden.
- <sup>28</sup> Department of Medicine, Abramson Cancer Center, Perelman School of Medicine at the University of Pennsylvania, Philadelphia, PA, USA.
- <sup>29</sup> Genetic Counseling Unit, Hereditary Cancer Program, IDIBGI (Institut d'Investigació Biomèdica de Girona), Catalan Institute of Oncology, CIBERONC, Girona, Spain.
- <sup>30</sup> Department of Medicine, Huntsman Cancer Institute, Salt Lake City, UT, USA.

- <sup>31</sup> Medical Oncology Department, Hospital Clínico San Carlos, Instituto de Investigación Sanitaria San Carlos (IdISSC), Centro Investigación Biomédica en Red de Cáncer (CIBERONC), Madrid, Spain.
- <sup>32</sup> Section of Molecular Genetics, Dept. of Laboratory Medicine, University Hospital of Pisa, Pisa, Italy.
- <sup>33</sup> Peter MacCallum Cancer Center, Melbourne, Victoria, Australia.
- <sup>34</sup> Sir Peter MacCallum Department of Oncology, The University of Melbourne, Melbourne, Victoria, Australia.
- <sup>35</sup> Department of Gynaecological Oncology, Chris O'Brien Lifehouse and The University of Sydney, Camperdown, Australia.
- <sup>36</sup> CRCHU de Québec- oncologie, Centre des maladies du sein Deschênes-Fabia, Hôpital du Saint-Sacrement, Québec, QC, Canada.
- <sup>37</sup> Departments of Pediatrics and Medicine, Columbia University, New York, NY, USA.
- <sup>38</sup> Centre for Medical Genetics, Ghent University, Gent, Belgium.
- <sup>39</sup> Department of Clinical Genetics, Erasmus University Medical Center, Rotterdam, The Netherlands.
- <sup>40</sup> Service de Génétique, CHU de Besançon, Besançon, France.
- <sup>41</sup> Department of Laboratory Medicine and Pathology, Mayo Clinic, Rochester, MN, USA.
- <sup>42</sup> Department of Clinical Genetics, Fox Chase Cancer Center, Philadelphia, PA, USA.
- <sup>43</sup> Unité d'Oncogénétique, ICO-Centre René Gauducheau, Saint Herblain, France.
- <sup>44</sup> Oncogenetics Group, Clinical and Molecular Genetics Area, Vall d'Hebron Institute of Oncology (VHIO), University Hospital Vall d'Hebron, Barcelona, Spain.
- <sup>45</sup> Department of Genetics, University of Pretoria, Arcadia, South Africa.
- <sup>46</sup> Nottingham Clinical Genetics Service, Nottingham University Hospitals NHS Trust, Nottingham, UK.
- <sup>47</sup> Centre for Cancer Genetic Epidemiology, Department of Oncology, University of Cambridge, Cambridge, UK.
- <sup>48</sup> Oncogenetics Team, The Institute of Cancer Research and Royal Marsden NHS Foundation Trust, London, UK.
- <sup>49</sup> Institute for Medical Informatics, Statistics and Epidemiology, University of Leipzig, Leipzig, Germany.
- <sup>50</sup> LIFE - Leipzig Research Centre for Civilization Diseases, University of Leipzig, Leipzig, Germany.
- <sup>51</sup> Division of Evolution and Genomic Medicine, School of Biological Sciences, Faculty of Biology, Medicine and Health, University of Manchester, Manchester Academic Health Science Centre, Manchester, UK.
- <sup>52</sup> Manchester Centre for Genomic Medicine, St Mary's Hospital, Central Manchester University Hospitals NHS Foundation Trust, Manchester Academic Health Science Centre, Manchester, UK.
- <sup>53</sup> Unité d'oncogénétique, Centre de Lutte Contre le Cancer, Centre Georges-François Leclerc, Dijon, France.
- <sup>54</sup> Centre de Génétique, CHU Dijon, Dijon, France.
- <sup>55</sup> Molecular Diagnostic Unit, Hereditary Cancer Program, ICO-IDIBELL (Bellvitge Biomedical Research Institute, Catalan Institute of Oncology), CIBERONC, Barcelona, Spain.

<sup>56</sup> Department of Cancer Epidemiology and Genetics, Masaryk Memorial Cancer Institute, Brno, Czech Republic.

<sup>57</sup> The Susanne Levy Gertner Oncogenetics Unit, Chaim Sheba Medical Center, Ramat Gan, Israel.

<sup>58</sup> Sackler Faculty of Medicine, Tel Aviv University, Ramat Aviv, Israel.

<sup>59</sup> Schools of Medicine and Public Health, Division of Cancer Prevention & Control Research, Jonsson Comprehensive Cancer Centre, UCLA, Los Angeles, CA, USA.

<sup>60</sup> Cancer Risk and Prevention Clinic, Dana-Farber Cancer Institute, Boston, MA, USA.

<sup>61</sup> Centre of Familial Breast and Ovarian Cancer, Department of Medical Genetics, Institute of Human Genetics, University Würzburg, Würzburg, Germany.

<sup>62</sup> Department of Pathology and Laboratory Medicine, Kansas University Medical Center, Kansas City, KS, USA.

<sup>63</sup> Department of Clinical Genetics and GROW, School for Oncology and Developmental Biology, Maastricht University Medical Center, Maastricht, The Netherlands.

<sup>64</sup> Molecular Genetics of Breast Cancer, German Cancer Research Center (DKFZ), Heidelberg, Germany.

<sup>65</sup> Centre for Epidemiology and Biostatistics, Melbourne School of Population and Global Health, The University of Melbourne, Melbourne, Victoria, Australia.

<sup>66</sup> Center for Medical Genetics, NorthShore University HealthSystem, Evanston, IL, USA.

<sup>67</sup> The University of Chicago Pritzker School of Medicine, Chicago, IL, USA.

<sup>68</sup> N.N. Petrov Institute of Oncology, St. Petersburg, Russia.

<sup>69</sup> Lombardi Comprehensive Cancer Center, Georgetown University, Washington, DC, USA.

<sup>70</sup> Clinical Genetics, Guy's and St Thomas' NHS Foundation Trust, London, UK.

<sup>71</sup> Independent Laboratory of Molecular Biology and Genetic Diagnostics, Pomeranian Medical University, Szczecin, Poland.

<sup>72</sup> Hematology, oncology and transfusion medicine center, Dept. of Molecular and Regenerative Medicine, Vilnius University Hospital Santariskiu Clinics, Vilnius, Lithuania.

<sup>73</sup> Department of Medicine, Division of Oncology, Stanford Cancer Institute, Stanford University School of Medicine, Stanford, CA, USA.

<sup>74</sup> Women's Cancer Program at the Samuel Oschin Comprehensive Cancer Institute, Cedars-Sinai Medical Center, Los Angeles, CA, USA.

<sup>75</sup> Department of Human Genetics, Radboud University Medical Center, Nijmegen, The Netherlands.

<sup>76</sup> Département de Génétique, CHU de Grenoble, Grenoble, France.

<sup>77</sup> Genetic Epidemiology of Cancer team, Inserm U900, Paris, France.

<sup>78</sup> Institut Curie, Paris, France.

<sup>79</sup> Mines ParisTech, Fontainebleau, France.

<sup>80</sup> Department of Clinical Genetics, Amsterdam UMC, location VU University Medical Center, Amsterdam, The Netherlands.

<sup>81</sup> Department of Gynecology and Obstetrics, Ludwig Maximilian University of Munich, Munich, Germany.

<sup>82</sup> NRG Oncology, Statistics and Data Management Center, Roswell Park Cancer Institute, Buffalo, NY, USA.

- <sup>83</sup> Immunology and Molecular Oncology Unit, Veneto Institute of Oncology IOV - IRCCS, Padua, Italy.
- <sup>84</sup> Service de Génétique, Institut Curie, Paris, France.
- <sup>85</sup> Clinical Cancer Genomics, City of Hope, Duarte, CA, USA.
- <sup>86</sup> Department of Population Sciences, Beckman Research Institute of City of Hope, Duarte, CA, USA.
- <sup>87</sup> Department of Obstetrics and Gynecology, Helsinki University Hospital, University of Helsinki, Helsinki, Finland.
- <sup>88</sup> Center for Genomic Medicine, Rigshospitalet, Copenhagen University Hospital, Copenhagen, Denmark.
- <sup>89</sup> Clinical Genetics Research Lab, Department of Cancer Biology and Genetics, Memorial Sloan-Kettering Cancer Center, New York, NY, USA.
- <sup>90</sup> Clinical Genetics Service, Department of Medicine, Memorial Sloan-Kettering Cancer Center, New York, NY, USA.
- <sup>91</sup> Department of Molecular Genetics, National Institute of Oncology, Budapest, Hungary.
- <sup>92</sup> West Midlands Regional Genetics Service, Birmingham Women's Hospital Healthcare NHS Trust, Birmingham, UK.
- <sup>93</sup> Department of Genetics, University Medical Center Groningen, University Groningen, Groningen, The Netherlands.
- <sup>94</sup> Department of Molecular Medicine, University La Sapienza, Rome, Italy.
- <sup>95</sup> Department of Genetics and Computational Biology, QIMR Berghofer Medical Research Institute, Brisbane, Queensland, Australia.
- <sup>96</sup> Genome Diagnostics Program, IFOM - the FIRC (Italian Foundation for Cancer Research) Institute of Molecular Oncology, Milan, Italy.
- <sup>97</sup> Dept of OB/GYN and Comprehensive Cancer Center, Medical University of Vienna, Vienna, Austria.
- <sup>98</sup> Unit of Molecular Bases of Genetic Risk and Genetic Testing, Department of Research, Fondazione IRCCS (Istituto Di Ricovero e Cura a Carattere Scientifico) Istituto Nazionale dei Tumori (INT), Milan, Italy.
- <sup>99</sup> School of Women's and Children's Health, Faculty of Medicine, University of NSW Sydney, Sydney, New South Wales, Australia.
- <sup>100</sup> The Kinghorn Cancer Centre, Garvan Institute of Medical Research, Sydney, New South Wales, Australia.
- <sup>101</sup> Clinical Genetics, Karolinska Institutet, Stockholm, Sweden.
- <sup>102</sup> Clalit National Cancer Control Center, Carmel Medical Center and Technion Faculty of Medicine, Haifa, Israel.
- <sup>103</sup> Division of Gynecologic Oncology, NorthShore University HealthSystem, University of Chicago, Evanston, IL, USA.
- <sup>104</sup> Wexner Medical Center, The Ohio State University, Columbus, OH, USA.
- <sup>105</sup> Cancer Genetics and Prevention Program, University of California San Francisco, San Francisco, CA, USA.
- <sup>106</sup> Genomics Center, Centre Hospitalier Universitaire de Québec – Université Laval, Research Center, Québec City, QC, Canada.

- <sup>107</sup> Institute of Cell and Molecular Pathology, Hannover Medical School, Hannover, Germany.
- <sup>108</sup> Department of Tumour Biology, INSERM U830, Paris, France.
- <sup>109</sup> Université Paris Descartes, Paris, France.
- <sup>110</sup> Department of Genetics, Portuguese Oncology Institute, Porto, Portugal.
- <sup>111</sup> Biomedical Sciences Institute (ICBAS), University of Porto, Porto, Portugal.
- <sup>112</sup> Department of Epidemiology, Mailman School of Public Health, Columbia University, New York, NY, USA.
- <sup>113</sup> Department of Clinical Genetics, Odense University Hospital, Odense C, Denmark.
- <sup>114</sup> Program in Cancer Genetics, Departments of Human Genetics and Oncology, McGill University, Montréal, QC, Canada.
- <sup>115</sup> Department of Medical Genetics, University of Cambridge, Cambridge, UK.
- <sup>116</sup> Department of Cancer Biology and Genetics, The Ohio State University, Columbus, OH, USA.
- <sup>117</sup> Department of Medical Oncology, Beth Israel Deaconess Medical Center, Boston, MA, USA.
- <sup>118</sup> Department of Clinical Genetics, Leiden University Medical Center, Leiden, The Netherlands.
- <sup>119</sup> Department of Clinical Genetics, Amsterdam UMC, Vrije Universiteit Amsterdam, Amsterdam, The Netherlands.
- <sup>120</sup> Department of Medical Oncology, CHU Dupuytren, Limoges, France.
- <sup>121</sup> Oxford Regional Genetics Service, Churchill Hospital, Oxford, UK.
- <sup>122</sup> Molecular Diagnostics Laboratory, INRASTES, National Centre for Scientific Research 'Demokritos', Athens, Greece.
- <sup>123</sup> The Hereditary Breast and Ovarian Cancer Research Group Netherlands (HEBON), Coordinating center: The Netherlands Cancer Institute, Amsterdam, The Netherlands.
- <sup>124</sup> Department of Dermatology, Huntsman Cancer Institute, University of Utah School of Medicine, Salt Lake City, UT, USA.
- <sup>125</sup> Center for Clinical Cancer Genetics, The University of Chicago, Chicago, IL, USA.
- <sup>126</sup> Harvard T.H. Chan School of Public Health, Boston, MA, USA.
- <sup>127</sup> Dana-Farber Cancer Institute, Boston, MA, USA.
- <sup>128</sup> Department of Public Health Sciences, University of Chicago, Chicago, IL, USA.

**Corresponding author:** Dezheng Huo, 5841 S. Maryland Ave., MC 2000, Chicago, Illinois 60637. Phone: (773) 834-0843; E-mail: [dhuo@health.bsd.uchicago.edu](mailto:dhuo@health.bsd.uchicago.edu)

**Running title:** Height, body mass index, and ovarian cancer risk

**Abstract:** 200 words (limits 200)

**Text:** 3322 words (limits 5000)

## Abstract

**Background** Height and body mass index (BMI) are associated with higher ovarian cancer risk in the general population, but whether such associations exist among *BRCA1/2* mutation carriers is unknown.

**Methods** We applied a Mendelian randomization approach to examine height/BMI with ovarian cancer risk using the Consortium of Investigators for the Modifiers of *BRCA1/2* (CIMBA) dataset, comprising 14,676 *BRCA1* and 7,912 *BRCA2* mutation carriers, with 2,923 ovarian cancer cases. We created a height genetic-score (height-GS) using 586 height-associated variants and a BMI genetic-score (BMI-GS) using 93 BMI-associated variants. Associations were assessed using weighted Cox models.

**Results** Observed height had no significant association with ovarian cancer risk (hazard ratio [HR]: 1.07 per 10-cm increase in height, 95% confidence interval [CI]: 0.94–1.23). Height-GS showed similar results (HR=1.02, 95% CI: 0.85–1.23). Higher BMI appeared to increase risk in premenopausal women significantly with HR=1.25 (95% CI: 1.06–1.48) and HR=1.59 (95% CI: 1.08–2.33) per 5-kg/m<sup>2</sup> increase in observed and genetically determined BMI, respectively. No association was found for postmenopausal women. Interaction between menopausal status and BMI was significant ( $P_{interaction} < 0.05$ ).

**Conclusion** Our observation of a positive association between BMI and ovarian cancer risk in premenopausal *BRCA1/2* mutation carriers was consistent with findings in the general population.

**Keywords:** ovarian neoplasms, Mendelian randomization, *BRCA1*, *BRCA2*, body height, body mass index

## Introduction

Ovarian cancer is the fifth leading cause of cancer deaths in US women, due to its typically advanced stage at presentation.<sup>1,2</sup> Furthermore, unlike breast or colorectal cancer, there is no proven screening method for ovarian cancer to identify early disease and initiate treatment to improve survival.<sup>3,4</sup> Family history, oral contraceptive use, parity, body mass index (BMI), and genetic variants are potentially useful in estimating lifetime risk.<sup>1</sup> In particular, inherited *BRCA1* and *BRCA2* mutations are associated with increased lifetime risk of ovarian cancer and account for ~10-15% of overall disease incidence.<sup>5-7</sup> However, among mutation carriers, age at diagnosis is variable. Penetrance of *BRCA1/2* mutations is likely modified by other genetic variants and lifestyle/reproductive factors.<sup>8,9</sup> Investigation of these factors could aid in implementation of strategies to reduce ovarian cancer risk among mutation carriers.

Both height and BMI are quantitative traits with substantial genetic bases. In recent genome-wide association studies (GWAS), numerous genetic variants were found to be associated with these traits.<sup>10,11</sup> In the general population, both height and BMI appear to be positively but inconsistently associated with risk of ovarian cancer.<sup>12-14</sup> Previous studies also showed that the association between BMI and ovarian cancer was stronger in premenopausal women.<sup>12,15,16</sup> Because of differences in age at onset and tumor histology/grade, risk factors for ovarian cancer might be different for *BRCA1/2* mutation carriers and general populations.<sup>17</sup> Only one case-control study, with 469 ovarian cancer cases, has examined anthropometric measurements in *BRCA1/2* mutation carriers and found that neither height nor BMI were related to ovarian cancer risk.<sup>18</sup> Larger, adequately-powered studies are needed to assess whether a relationship between either height or BMI and ovarian cancer risk exists for *BRCA1/2* mutation carriers, and if the direction of association is concordant with that in the general population.



Mendelian randomization (MR) methods use genetic markers associated with a trait as an instrumental variable (IV) to assess their potential relationship with a disease outcome.<sup>19-21</sup> Compared to traditional epidemiologic approaches, MR can reduce biases such as reverse causation and residual confounding, that can interfere with causal interpretations. However, the MR approach requires that the genetic variants are associated with the exposure, the variants are not or are only weakly associated with confounding factors in the causal pathway, and the variants only affect disease risk through the exposure (i.e. absence of pleiotropic effects).<sup>20,21</sup> To the degree that these assumptions are met, the MR approach can strengthen the evidence for a causal relationship between exposure and disease.

Herein, using traditional epidemiologic and MR methods, we conducted analyses of height and BMI on ovarian cancer risk in the Consortium of Investigators for the Modifiers of *BRCA1/2* (CIMBA), with 22,588 participants. We examined heterogeneity of these associations with respect to the mutation carried (*BRCA1* vs *BRCA2*), menopausal status, tumor histology, and tumor grade.

## **Methods**

Characteristics of the CIMBA consortium and information on specific genotyping protocols are provided in the **Supplementary Methods** and were described previously.<sup>22-24</sup>

### *Selection of Genetic Variants*

From the latest publications of the Genetic Investigation of Anthropometric Traits, we identified SNPs associated with height or BMI at genome-wide significance level ( $P < 5 \times 10^{-7}$ )

<sup>8</sup>).<sup>11,25</sup> SNPs with low imputation quality (<0.5) were excluded, leaving 586 SNPs for height and 93 for BMI. **Supplementary Tables 1 and 2** provide additional details on these SNPs.

### *Statistical Analysis*

Calculation of the height- and BMI-genetic scores (GS) was described in detail previously.<sup>24</sup> Briefly, we calculated the weighted sums of all of the height- and BMI-associated variants under additive models, which do not include interactions between variants. Namely, we used the formulas:  $Height - GS = \sum_{i=1}^{586} \beta_{XGi} SNP_i$  and  $BMI - GS = \sum_{i=1}^{93} \beta_{XGi} SNP_i$ , where  $\beta_{XGi}$  is the literature-reported per-allele magnitude of association of the *i*th-SNP for height and BMI, respectively. A scaling factor was calculated by regressing each GS against its respective trait among non-case carriers. The corresponding regression coefficients were  $\beta_0$  (intercept=165.455) and  $\beta_1$  (slope=5.217) for height and  $\beta_0$  (22.607) and  $\beta_1$  (5.523) for BMI. In the present study, BMI-GS was scaled to BMI at date of questionnaire, rather than BMI at age 18, as previous GWAS have been based on BMI measurements in middle-aged adults.

We subsequently modeled each scaled-GS against ovarian cancer risk using weighted Cox models. Our primary outcome of interest was ovarian cancer diagnosis, with individuals censored for breast cancer diagnosis, risk-reducing bilateral salpingo-oophorectomy, death, or end of follow-up, whichever occurred first. Due to the study design of CIMBA, weights in the model were applied for cases and non-cases based on previously observed incidence of ovarian cancer in *BRCA1/2* carriers.<sup>26,27</sup> We applied a robust sandwich variance-estimation approach to the risk estimates to account for non-independence among multiple carriers per family. Additionally, we performed subgroup analyses by *BRCA1/2* mutations and menopausal status. Menopausal status was defined as a time-varying covariate, coded as premenopausal from birth

until age at natural menopause or bilateral salpingo-oophorectomy. For individuals with missing age at menopause, we imputed the age as 50. Imputing missing age at menopause as 46 did not materially change the results. The mean and median ages at natural menopause in this population were 46 and 48, respectively. All analyses were adjusted for the first eight principal components (to account for ethnicity and population stratification), birth cohort, and country of enrollment. Additional analyses assessed the associations of height and BMI with ovarian cancer subgroups, by histological type (serous vs. non-serous) and by tumor grade (well- or moderately-differentiated tumors vs. poorly- or un-differentiated).

Additionally, phenotype associations with each individual height- and BMI-variant were assessed and pooled using inverse variance-weighted meta-analysis. The individual associations were obtained by first extracting  $\beta_{XGi}$  for each SNP  $i$ , which represents the per-allele magnitude of association with height or BMI from previous GWAS. Next, we calculated  $\beta_{YGi}$  and  $SE(\beta_{YGi})$  using multivariate-adjusted weighted Cox models for each SNP using the CIMBA data, where ovarian cancer risk is predicted by genotype  $G$  (with  $G=0,1,2$  for the allele corresponding to greater height or BMI), principal components, birth cohort, *BRCA* mutation, and country of enrollment. The overall causal association ( $\beta_{YX}$ ) is calculated using inverse-variance weighted

estimate of each variant's effect:  $\beta_{YX} = \frac{\sum_i \beta_{XGi} \beta_{YGi} SE(\beta_{YGi})^{-2}}{\sum_i \beta_{XGi}^2 SE(\beta_{YGi})^{-2}}$ . Standard error was estimated as

$SE_{YX} = \sqrt{\frac{1}{\sum_i \beta_{XGi}^2 SE(\beta_{YGi})^{-2}}}$  using the Burgess's method.<sup>19,28</sup> Egger's test was used to assess for

possible pleiotropic effects of the variants (i.e. whether variants influence the outcome through other pathways), to ensure that this assumption held.<sup>29</sup>

Finally, in participants with available data on height and BMI, we conducted a formal IV analysis using the method of two-stage residual inclusion regression.<sup>30</sup> In stage one, observed height or BMI was regressed against the corresponding GS, principal components, birth cohort,

country, and mutation status. In the second stage, we used a Cox model to fit ovarian cancer risk against height or BMI, birth cohort, country, mutation status, and residuals from stage one. Variance estimates were obtained through 10,000 boot-straps (see details in **Supplementary Methods**). In these individuals, we also analyzed the association between observed measurements of height or BMI and ovarian cancer risk using weighted Cox models, adjusted for established ovarian cancer risk factors including birth cohort, menopausal status, age at menarche (years) and parity (continuous). The BMI values used were obtained at the date of questionnaire, usually close to the date of genetic testing and recalled for BMI at age 18.

In models with menopausal status as time-varying variable, the test for heterogeneity by menopausal status was essentially a test of the proportional hazards assumption. All analyses were performed using SAS 9.4 (SAS Institute, Cary, NC) and Stata 14.0 (StataCorp, College Station, TX). A two-sided  $P$ -value $<0.05$  was considered statistically significant unless stated otherwise.

## Results

### *Demographic and Clinical Characteristics*

Characteristics for the 22,588 individuals in the CIMBA consortium, comprising 14,676 *BRCA1* and 7,912 *BRCA2* mutation carriers, are shown in **Table 1**. We documented 2,923 women with ovarian cancer (*BRCA1*: 2,319; *BRCA2*: 604). Compared with non-cases, participants who developed ovarian cancer were more often parous women, were younger at first live birth, and were from earlier birth cohorts. At the date of questionnaire/interview, height measurement was available for 7,657 participants and BMI measurement for 7,516 participants.

Most tumors for *BRCA1/2* mutation carriers were invasive, of serous, poorly/undifferentiated grade, and stages 3 or 4 at diagnosis, characteristics which are consistent with prior reports.<sup>31</sup>

### *Observed and Predicted Height on Risk of Ovarian Cancer*

In the survival modelling of ovarian cancer risk, age was used as the underlying time scale and the numbers of individuals retained in the analysis were 20535, 14647, 7375, and 2832 at ages 30, 40, 50, and 60 years, respectively, suggesting that statistical power for the late age is limited. After adjustment for birth cohort, country of enrollment, mutation, menopausal status, and principal components, a nonsignificant association was found for observed height and ovarian cancer risk (HR=1.07 per 10-cm increase, 95%CI: 0.94–1.23,  $P=0.31$ ) (**Table 2**). We found broadly consistent associations of height in both *BRCA1* and *BRCA2* mutation carriers, by menopausal status, and by tumor histological type and grade.

The height-GS was significantly associated with height in all participants, in ovarian cancer cases, and in non-case participants (all  $P<10^{-24}$ ) (**Supplementary Table 3**). Overall, approximately 13.4% of the variation in height was explained by the height-GS. Besides height, we found weaker associations between the height-GS and body weight, and age-at-menarche.

In MR analysis, height-GS had a nonsignificant positive association with ovarian cancer risk, HR=1.02 per 10-cm increase in genetically-predicted height, 95%CI: 0.85–1.23,  $P=0.82$  (**Table 3**). We found similar associations by subgroups of mutation, menopausal status, and tumor grade.

Combining the effects of all 586 height-associated variants using inverse-variance weighted meta-analysis, we obtained similar findings (HR=1.02, 95%CI: 0.83-1.26,  $P=0.83$ ) (**Table 3**). Among the SNPs that were combined, there was a low degree of heterogeneity

( $I^2=0\%$ ). Examining small-study effects using Egger's test did not suggest likely pleiotropic effects. In the two-stage residual inclusion analysis, the estimated relative risk was larger though with wide CIs, which overlapped with those derived using other methods (HR=1.20, 95%CI: 0.86–1.69,  $P=0.29$ ).

#### *Observed and Predicted BMI on Risk of Ovarian Cancer*

After multivariate adjustment, we found a nonsignificant positive association between BMI at date of questionnaire and ovarian cancer risk, HR=1.04 per 5-kg/m<sup>2</sup> increase in BMI, 95%CI: 0.95–1.14,  $P=0.42$  (**Table 4**). In a pre-specified analysis, the association between BMI and ovarian cancer risk was stronger in premenopausal women (HR=1.25, 95%CI: 1.06–1.48;  $P=0.009$ ) but not in postmenopausal women (HR=0.98, 95%CI: 0.88–1.10), with significant interaction ( $P=0.02$ ). We found that BMI was a significant predictor of non-serous ovarian cancer risk (HR=1.25, 95%CI: 1.06–1.49), but not for serous ovarian cancer (HR=0.98, 95%CI: 0.84–1.15).

Similar to BMI at date of questionnaire, we detected a significant interaction of BMI in young adulthood and menopausal status ( $P=0.01$ ), with stronger magnitude in premenopausal (HR=1.34, 95%CI: 0.97–1.84) compared with postmenopausal women (HR=0.82, 95%CI: 0.65–1.04).

BMI-GS was strongly associated with BMI at both date of questionnaire and young adulthood (**Supplementary Table 4**). Overall, the BMI-GS explained 2.6% of the variation in BMI at date of questionnaire and 1.7% of the variation in young adulthood BMI. We found associations between the BMI-GS and height and age-at-menarche, though the strength of the association was weaker than with BMI.

In the entire consortium, the BMI-GS had a nonsignificant positive association with ovarian cancer risk with a HR=1.10 per 5-kg/m<sup>2</sup> of genetically predicted BMI, 95%CI: 0.86–1.42,  $P=0.44$  (**Table 5**). We found heterogeneity by menopausal status ( $P=0.006$ ). BMI-GS was positively associated with ovarian cancer risk in premenopausal women (HR=1.59, 95%CI: 1.08–2.33) but not in postmenopausal women (HR=0.80, 95%CI: 0.58–1.11). BMI-GS also tended to be more associated with non-serous (HR=1.60, 95%CI: 0.83–3.08) than serous tumors (HR=0.92, 95%CI: 0.59–1.43).

We found similar results when we statistically combined the associations of the 93 BMI-associated variants, with an overall HR=1.12, 95%CI: 0.86–1.46. Heterogeneity was low ( $I^2=15.9\%$ ), indicating a low likelihood of pleiotropic associations. Using the two-stage residual inclusion approach, we found a generally similar association (HR=1.37, 95%CI: 0.84–2.24,  $P=0.21$ ).

#### *Individual SNPs and Ovarian Cancer Risk*

We found 22 height-associated and 4 BMI-associated SNPs that were nominally associated with ovarian cancer risk ( $P<0.05$ ) (**Table 6**). None of these SNPs were significantly associated with ovarian cancer risk after correcting for multiple testing. We cross-checked these identified SNPs with the most up-to-date list of ovarian cancer susceptibility SNPs and did not find any overlaps.<sup>32</sup>

## **Discussion**

Using data from a large international consortium of *BRCA1/2* mutation carriers, we found no statistically significant association between height and ovarian cancer risk. Interestingly, we

observed interactions between BMI (both observed and genetically predicted) and menopausal status on ovarian cancer risk, with increasing BMI associated with increased risk in premenopausal but not in postmenopausal women.

Our finding of a positive association between BMI and overall ovarian cancer risk in *BRCA* mutation carriers is corroborated by several prior studies in the general populations.<sup>12,14,15,33</sup> One MR analysis using 77 BMI-associated SNPs, conducted in the general population, found that each 1-standard deviation (SD) increment in genetically-predicted adult BMI corresponded to an odds ratio (OR) of 1.35 (95%CI 1.05-1.72).<sup>34</sup> We found that 5-kg/m<sup>2</sup> (about 1 SD) increment in genetically-predicted BMI was associated with a HR=1.10 (95%CI: 0.86–1.42) in mutation carriers. However, the association of BMI with ovarian cancer risk is likely to vary by menopausal status. In the general populations, significant differential association of BMI with ovarian cancer risk by menopausal status has been found in some studies<sup>15,16,35,36</sup> but not others.<sup>12,37</sup> A pooled analysis of 47 epidemiologic studies with 25,157 ovarian cancer cases showed that the relative risk per 5-kg/m<sup>2</sup> increase in BMI was 1.12 (95%CI: 1.07–1.17) in premenopausal women and 1.08 (95%CI: 1.04–1.12) in postmenopausal women.<sup>12</sup> The largest single cohort study, with 3,686 ovarian cancer cases, found the HR per 5-kg/m<sup>2</sup> increase in BMI was 1.21 (99%CI: 1.09–1.33) in premenopausal and 1.07 (99%CI: 1.02–1.12) in postmenopausal women.<sup>15</sup> An MR analysis conducted in the general population also observed stronger associations for non-high grade serous carcinomas in premenopausal women (OR=1.62, 95%CI: 0.88–3.01) compared with postmenopausal hormone replacement therapy (HRT) users (OR=1.26, 95%CI: 0.57–2.82) and postmenopausal HRT non-users (OR=1.17, 95%CI: 0.61–2.24), though no formal statistical tests examining heterogeneity were performed.<sup>14</sup> Similarly, we found in *BRCA1/2* mutation carriers that 5-kg/m<sup>2</sup> increment in genetically-predicted BMI was



associated with a HR=1.59 (95%CI: 1.08-2.33) for premenopausal ovarian cancer and a HR=0.80 (95%CI: 0.58-1.11) for postmenopausal ovarian cancer. Studies which have not demonstrated significant variation by menopausal status tended to show that the positive association between BMI and ovarian cancer risk was primarily among those who had never used HRT.<sup>12</sup> Taken together, our results and previous literature are suggestive that higher BMI may increase ovarian cancer risk in premenopausal women, but not in postmenopausal women.

Additionally, several studies that had sufficient numbers of patients to evaluate the relationship between BMI and ovarian cancer risk by histologic subtype have shown significant heterogeneity. Observational studies in the Ovarian Cancer Cohort Consortium found stronger associations between BMI and endometrioid (OR=1.17 per 5-kg/m<sup>2</sup>, 95%CI: 1.11–1.23) or mucinous ovarian cancer (OR=1.19, 95%CI: 1.06–1.32), but no association with serous ovarian cancer (OR=0.98, 95%CI: 0.94–1.02).<sup>16</sup> A more recent MR analysis in the same consortium using a genetic score comprised of 87 SNPs showed that a genetically predicted BMI had stronger association with endometrioid (OR=1.17, 95%CI: 0.87-1.59) or mucinous ovarian cancer (OR=1.18, 95%CI: 0.84-1.67) than high grade serous cancer (OR=1.06, 95%CI: 0.89-1.27), though the 95%CIs for these estimates were largely overlapping.<sup>14</sup> Consistent with findings in the general population, our study in *BRCA1/2* mutation carriers showed that BMI was positively associated with non-serous ovarian cancer (HR=1.25 per 5-kg/m<sup>2</sup>, 95%CI: 1.06-1.49 in observed BMI and HR=1.60, 95%CI: 0.83-3.08, per 5-kg/m<sup>2</sup> in genetically predicted BMI), of which endometrioid is a major subtype. Of note, obesity is an established risk factor for endometrial cancer.<sup>38</sup> However, subsequent studies with greater number of cases of different ovarian cancer subtypes are needed to assess whether the effect of obesity truly differs by tumor subtype.

Our finding of a nonsignificant positive association between height and ovarian cancer risk is also consistent with prior epidemiological studies in general populations.<sup>12,37,39</sup> In the general population, 5-cm increment in height was associated with a 7% increase (95%CI: 5-9%) in ovarian cancer risk,<sup>12</sup> and 5-cm increment in genetically predicted height was associated with a 6% (95%CI: 1–11%) increase in ovarian cancer risk.<sup>39</sup> The associations for observed height did not differ significantly between ovarian histological types,<sup>2,12</sup> while genetically predicted height has stronger association with clear cell (OR=1.20, 95%CI: 1.04–1.38) or low grade/borderline serous ovarian cancers (OR=1.15, 95%CI: 1.01–1.30), compared to high-grade serous (OR=1.05, 95%CI: 0.99–1.11).<sup>39</sup> We did not find statistically significant heterogeneity by histology in our study of mutation carriers, though point estimates varied across histology.

Several biological mechanisms potentially explain the associations observed in our study. Overweight/obese women are more likely to have anovulatory cycles and fertility issues, particularly when caused by polycystic ovarian syndrome (PCOS), and thus have an increased risk of ovarian cancer.<sup>40,41</sup> The association of PCOS and ovarian cancer was mainly confined to premenopausal women.<sup>42</sup> Some studies have suggested that *BRCA1/2* mutation carriers may have subclinical ovarian insufficiency, which could mediate the relationship between obesity-related infertility and increased ovarian cancer risk.<sup>43</sup> Obesity itself also creates a proinflammatory state and adipocyte secreted inflammatory markers have been implicated in ovarian cancer development.<sup>44</sup> Circulating levels of estradiol, androgen, and progesterone have also been implicated in the risk of ovarian cancer.<sup>45,46</sup> One study in *BRCA1/2* mutation carriers showed higher estradiol levels during each menstrual cycle compared with non-carriers, supporting the potential role of sex hormones in ovarian tumorigenesis in this population.<sup>47</sup> Obese premenopausal women tend to have lower circulating levels of progesterone compared with

normal weight women.<sup>48</sup> Higher progesterone levels may reduce ovarian cancer risk, through upregulation of p53, leading to tumor cell apoptosis.<sup>46,49-51</sup> Taken together, these pathways may explain the association of higher BMI with premenopausal ovarian cancer risk. Additionally, height has been associated with higher levels of circulating insulin-like growth factor-1 (IGF-1),<sup>52,53</sup> a pathway that has been implicated in tumor transformation and may exert anti-apoptotic and mitogenic effects.<sup>54,55</sup> Moreover, *BRCA1* may directly interact with the IGF-1 pathway to mediate cancer risk.<sup>56</sup>

Our study has several strengths, including large sample size, genetic scores utilizing most identified height- and BMI-variants, several MR methods, and consistent findings between observed and genetically-predicted phenotypes. Several limitations of our study should be considered. First, even with large sample size, the CIs for most risk estimates were wide, which limits inferences about causation. While both the height- and BMI-GS were clearly associated with their respective traits, they were only able to explain 13.4% and 2.6% of the variation, respectively. This reduced the statistical precision of our risk estimates. During the preparation of our manuscript, a new genome-wide meta-analysis<sup>57</sup> found a substantial number of new genetic loci related to height and BMI, increasing the amount of variation that could be explained for these two traits to 24.6% and 6.0%, though the variation that could be explained when examining these SNPs in a validation cohort was 14.0% and 2.3%. This is comparable to the amount of variation that could be explained using the set of genetic variants in our study. Including these additional SNPs may be able to improve the precision of our estimates for both height and BMI. Moreover, the inclusion of rare variants to strengthen the height and BMI genetic instruments should also be considered in future studies.<sup>58</sup> Our study did not explicitly examine whether adding height or BMI (either observed or genetically predicted) to existing

polygenic risk scores for ovarian cancer could further refine risk prediction. Histology was only available in a subset of ovarian cancer patients, which limits our capacity to understand subtype-specific effects of BMI and height. Our study only included women of European ancestry, which may preclude generalization to women of other racial/ethnic groups.

In summary, our study suggests that higher BMI may be causally associated with ovarian cancer risk in *BRCA1/2* carriers, possibly more so for premenopausal women. BMI could be used to identify premenopausal women at elevated risk of ovarian cancer. Our finding of stronger association between BMI and non-serous ovarian cancer warrants confirmation in future studies.

**Tables**

Variable	Ovarian cancer cases, <i>N</i> = 2923	Non-cases, <i>N</i> = 19,665	<i>P</i> value <sup>b</sup>
Mutation carrier status			<0.0001
<i>BRCA1</i>	2319 (79.3)	12,357 (62.8)	
<i>BRCA2</i>	604 (20.7)	7308 (37.2)	
Year of birth, median (IQR)	1948 (1940, 1955)	1960 (1951, 1969)	<0.0001
Age at diagnosis or censoring, years (mean ± SD)	52.5 ± 9.8	44.7 ± 12.4	<0.0001
Ethnicity, <i>n</i> (%)			0.07
Caucasian, not otherwise specified	2060 (89.7)	13,613 (88.4)	
Ashkenazi Jewish	237 (10.3)	1780 (11.6)	
Height in cm, <i>n</i>	784	6873	
Mean ± SD	163.2 ± 6.5	164.8 ± 6.9	<0.0001
Weight at baseline <sup>a</sup> in kg, <i>n</i>	780	6789	
Mean ± SD	69.0 ± 14.6	68.5 ± 14.1	0.32
Body mass index at baseline <sup>a</sup> in kg/m <sup>2</sup> , <i>n</i>	772	6744	
Mean ± SD	25.9 ± 5.3	25.2 ± 5.1	0.0002
Weight in early adulthood in kg, <i>n</i>	536	4,912	
Mean ± SD	56.5 ± 8.3	57.9 ± 9.5	0.0007
Body mass index in early adulthood in kg/m <sup>2</sup> , <i>n</i>	536	4881	
Mean ± SD	21.2 ± 3.0	21.3 ± 3.3	0.43
Age at menarche in years, <i>n</i>	771	6688	
Mean ± SD	13.0 ± 1.5	13.0 ± 1.5	0.90
Parous, <i>n</i> (%)			<0.0001
Yes	805 (88.3)	5790 (77.4)	
No	107 (11.7)	1692 (22.6)	
Age at first live birth in years, <i>n</i>	735	5555	
Mean ± SD	24.4 ± 4.5	25.4 ± 4.9	<0.0001
Menopausal status, <i>n</i> (%)			<0.0001
Premenopausal	112 (11.5)	3816 (51.1)	
Postmenopausal	863 (88.5)	3654 (48.9)	
Age at menopause, years (mean ± SD)	46.8 ± 5.7	44.7 ± 6.1	<0.0001
Tumour behaviour, <i>n</i> (%)			
Invasive	1228 (99.2)		
Borderline	10 (0.8)		
Tumour histotype, <i>n</i> (%)			
Serous	892 (67.9)		
Mucinous	20 (1.5)		
Endometrioid	141 (10.7)		
Clear cell	17 (1.3)		
Other	243 (18.5)		
Tumour grade, <i>n</i> (%)			
Well differentiated	43 (4.6)		
Moderately differentiated	196 (21.0)		

Poorly/ undifferentiated	696 (74.4)
Tumour stage, <i>n</i> (%)	
Borderline	2 (0.3)
Stage 1	121 (16.4)
Stage 2	93 (12.6)
Stage 3	412 (55.7)
Stage 4	112 (15.1)
<p><i>CIMBA</i> Consortium of Investigators for the Modifiers of <i>BRCA1/2</i>, <i>IQR</i> interquartile range, <i>SD</i> standard deviation  <sup>a</sup>Reported at the date of questionnaire  <sup>b</sup><i>P</i> values for comparing cases and non-cases were calculated from logistic regression models with robust sandwich variance estimator</p>	

Table 1. Baseline characteristics of participants in the CIMBA consortium with genotype information

	N/events	HR (95% CI)	P value
Per 10 cm increase in observed height			
All participants (confounding adjustment sequentially)			
Adjusted for principal components	7657/784	1.12 (0.97–1.29)	0.12
Additionally adjusted for country	7657/784	1.15 (1.00–1.32)	0.06
Additionally adjusted for birth cohort	7657/784	1.05 (0.91–1.21)	0.53
Additionally adjusted for mutation status	7657/784	1.06 (0.92–1.22)	0.42
<b>Additionally adjusted for menopausal status</b>	<b>7657/784</b>	<b>1.07 (0.94–1.23)</b>	<b>0.31</b>
Additionally adjusted for parity and age at menarche	7090/724	1.09 (0.94–1.26)	0.24
By mutation status <sup>a</sup>			
<i>BRCA1</i> carrier	4502/552	1.07 (0.91–1.24)	0.42
<i>BRCA2</i> carrier	3155/232	1.11 (0.85–1.45)	0.44
<i>P</i> <sub>interaction</sub>			0.64
By menopausal status <sup>b</sup>			
Premenopausal	7657/105	1.02 (0.72–1.42)	0.93
Postmenopausal	4328/679	1.09 (0.94–1.26)	0.27
<i>P</i> <sub>interaction</sub>			0.71
By tumour subtype <sup>c</sup>			
Serous	7360/319	1.07 (0.87–1.31)	0.52
Non-serous <sup>d</sup>	7360/168	1.30 (1.01–1.68)	0.045
<i>P</i> <sub>het</sub>			0.24
By tumour grade <sup>e</sup>			
Well or moderately differentiated	7252/111	1.12 (0.83–1.52)	0.46
Poorly/undifferentiated	7252/268	1.15 (0.93–1.43)	0.19
<i>P</i> <sub>het</sub>			0.89
<i>HR</i> hazard ratio, <i>CI</i> confidence interval			
<sup>a</sup> Adjusted for principal components, birth cohort, country of enrolment, and menopausal status in weighted Cox model			
<sup>b</sup> Adjusted for principal components, mutation status, birth cohort, and country of enrolment			
<sup>c</sup> Adjusted for principal components, birth cohort, country of enrolment, mutation status, and menopausal status			
<sup>d</sup> Includes endometrioid, mucinous, clear cell, and other histologic types			
Bolded line refers to the model corresponding to our main results			

Table 2. Association of height and ovarian cancer risk using observed height among 7657 participants



	N/events	HR (95% CI)	P value	Heterogeneity ( $I^2$ )
<b>Height GS<sup>a</sup></b>				
All participants (confounding adjustment sequentially)				
Adjusted for principal components	22,588/2923	0.99 (0.82–1.19)	0.89	
Additionally adjusted for country	22,588/2923	0.97 (0.81–1.17)	0.77	
Additionally adjusted for birth cohort	22,588/2923	0.98 (0.82–1.18)	0.83	
Additionally adjusted for mutation status	22,588/2923	1.02 (0.85–1.22)	0.13	
<b>Additionally adjusted for menopausal status</b>	22,588/2923	<b>1.02 (0.85–1.23)</b>	<b>0.82</b>	
By mutation status <sup>b</sup>				
BRCA1 carrier	14,676/2319	1.02 (0.83–1.25)	0.87	
BRCA2 carrier	7912/604	1.04 (0.68–1.57)	0.87	
$P_{\text{interaction}}$			0.99	
By menopausal status <sup>c</sup>				
Premenopausal	22,588/967	0.96 (0.73–1.26)	0.77	
Postmenopausal	9219/1955	1.08 (0.85–1.38)	0.52	
$P_{\text{interaction}}$			0.50	
By tumour subtype <sup>d</sup>				
Serous	20,978/892	1.36 (0.97–1.90)	0.08	
Non-serous	20,978/421	0.95 (0.58–1.56)	0.84	
$P_{\text{het}}$			0.25	
By tumour grade <sup>d</sup>				
Well or moderately differentiated	20,600/239	1.63 (0.86–3.09)	0.14	
Poorly/undifferentiated	20,600/696	1.20 (0.82–1.74)	0.35	
$P_{\text{het}}$			0.42	
Meta-analysis method <sup>e</sup>				
All participants	22,588/2923	1.02 (0.83–1.26)	0.83	0.0%
BRCA1 carrier	14,676/2319	1.02 (0.81–1.28)	0.89	0.0%
BRCA2 carrier	7912/604	1.05 (0.67–1.66)	0.82	7.0%
$P_{\text{interaction}}$			0.89	
Two-stage residual inclusion method <sup>f</sup>				
All participants	7657/784	1.20 (0.86–1.69)	0.29	
BRCA1 carrier	4502/552	1.40 (0.94–2.10)	0.10	
BRCA2 carrier	3155/232	0.93 (0.49–1.74)	0.81	

HR hazard ratio, CI confidence interval, CIMBA Consortium of Investigators for the Modifiers of BRCA1/2, GS genetic score  
<sup>a</sup>Height genetic score combining 586 height-associated single-nucleotide polymorphisms (SNPs)  
<sup>b</sup>Adjusted for principal components, birth cohort, country of enrolment, and menopausal status in weighted Cox model  
<sup>c</sup>Adjusted for principal components, mutation status, birth cohort, and country of enrolment  
<sup>d</sup>Adjusted for principal components, mutation status, menopausal status, birth cohort, and country of enrolment  
<sup>e</sup>HRs were calculated using inverse-variance meta-analysis and re-scaled to the corresponding units by calculating the height measurements per z-score among controls. Effect estimates for ovarian cancer for each SNP were calculated from weighted Cox model adjusting for principal components, birth cohort, country of enrolment, menopausal status, and mutation status  
<sup>f</sup>Analysis was performed among 7657 participants with measured height  
**Bolded line refers to the model corresponding to our main results**

Table 3. Association of height and ovarian cancer risk among 22,588 participants in CIMBA per 10-cm increase in genetically predicted height

	N/events	HR (95% CI)	P value
Per 5 kg/m <sup>2</sup> increase in BMI at date of questionnaire			
All participants (confounding adjustment sequentially)			
Adjusted for principal components	7516/772	1.00 (0.90–1.10)	0.96
Additionally adjusted for country	7516/772	0.99 (0.90–1.09)	0.84
Additionally adjusted for birth cohort	7516/772	1.02 (0.93–1.12)	0.72
Additionally adjusted for mutation status	7516/772	1.06 (0.96–1.16)	0.26
<b>Additionally adjusted for menopausal status</b>	<b>7516/772</b>	<b>1.04 (0.95–1.14)</b>	<b>0.42</b>
Additionally adjusted for parity and age at menarche	6964/715	1.04 (0.94–1.14)	0.48
By mutation status <sup>a</sup>			
BRCA1 carrier	4401/543	1.06 (0.95–1.17)	0.31
BRCA2 carrier	3115/229	0.96 (0.81–1.15)	0.67
<i>P</i> <sub>interaction</sub>			0.35
By menopausal status <sup>b</sup>			
<b>Premenopausal</b>	<b>7516/102</b>	<b>1.25 (1.06–1.48)</b>	<b>0.009</b>
Postmenopausal	4257/670	0.98 (0.88–1.10)	0.78
<i>P</i> <sub>interaction</sub>			<b>0.02</b>
By tumour subtype <sup>c</sup>			
Serous	7223/312	0.98 (0.84–1.15)	0.83
<b>Non-serous<sup>d</sup></b>	<b>7223/167</b>	<b>1.25 (1.06–1.49)</b>	<b>0.01</b>
<i>P</i> <sub>het</sub>			<b>0.04</b>
By tumour grade <sup>c</sup>			
Well or moderately differentiated	7252/109	1.05 (0.84–1.32)	0.65
Poorly/undifferentiated	7252/268	0.95 (0.82–1.11)	0.54
<i>P</i> <sub>het</sub>			0.47
Per 5 kg/m <sup>2</sup> increase in BMI in young adulthood			
All participants (confounding adjustment sequentially)			
Unadjusted	5417/536	0.86 (0.69–1.07)	0.17
Adjusted for country	5417/536	0.86 (0.69–1.08)	0.19
Additionally adjusted for birth cohort	5417/536	0.87 (0.70–1.08)	0.21
Additionally adjusted for mutation status	5417/536	0.91 (0.73–1.13)	0.39
Additionally adjusted for menopausal status	5417/536	0.93 (0.76–1.16)	0.53
Additionally adjusted for parity and age at menarche	5210/516	0.92 (0.74–1.14)	0.42
By mutation status <sup>a</sup>			
BRCA1 carrier	3134/380	0.92 (0.71–1.18)	0.50
BRCA2 carrier	2283/156	1.00 (0.74–1.36)	0.99
<i>P</i> <sub>interaction</sub>			0.73
By menopausal status <sup>b</sup>			
<b>Premenopausal</b>	<b>5417/67</b>	<b>1.34 (0.97–1.84)</b>	<b>0.07</b>
Postmenopausal	3094/469	0.82 (0.65–1.04)	0.11
<i>P</i> <sub>interaction</sub>			<b>0.01</b>
<i>HR</i> hazard ratio, <i>CI</i> confidence interval			
<sup>a</sup> Adjusted for principal components, birth cohort, country of enrolment, and menopausal status in weighted Cox model			
<sup>b</sup> Adjusted for principal components, mutation status, birth cohort, and country of enrolment			
<sup>c</sup> Adjusted for principal components, birth cohort, country of enrolment, mutation status, and menopausal status			
<sup>d</sup> Includes endometrioid, mucinous, clear cell, and other histological types			
Bolded lines refer to the model corresponding to our main results			

Table 4. Association of body mass index (BMI) and ovarian cancer risk using observed BMI

Breast cancer group	N/events	HR (95% CI)	P value	Heterogeneity ( $I^2$ )
<b>BMI-GS<sup>a</sup></b>				
All participants (confounding adjustment sequentially)				
Adjusted for principal components	22,588/2923	1.12 (0.87–1.45)	0.37	
Additionally adjusted for country	22,588/2923	1.11 (0.86–1.44)	0.41	
Additionally adjusted for birth cohort	22,588/2923	1.12 (0.87–1.45)	0.36	
Additionally adjusted for mutation status	22,588/2923	1.11 (0.86–1.42)	0.43	
<b>Additionally adjusted for menopausal status</b>	<b>22,588/2923</b>	<b>1.10 (0.86–1.42)</b>	<b>0.44</b>	
By mutation status <sup>b</sup>				
<i>BRCA1</i> carrier	14,676/2319	1.16 (0.88–1.53)	0.31	
<i>BRCA2</i> carrier	7912/604	0.81 (0.46–1.43)	0.46	
$P_{\text{interaction}}$			0.27	
By menopausal status <sup>c</sup>				
<b>Premenopausal</b>	<b>22,588/967</b>	<b>1.59 (1.08–2.33)</b>	<b>0.02</b>	
Postmenopausal	9219/1955	0.80 (0.58–1.11)	0.18	
$P_{\text{interaction}}$			<b>0.006</b>	
By tumour subtype <sup>d</sup>				
Serous	20,978/892	0.92 (0.59–1.43)	0.71	
<b>Non-serous</b>	<b>20,978/421</b>	<b>1.60 (0.83–3.08)</b>	<b>0.16</b>	
$P_{\text{het}}$			<b>0.17</b>	
By tumour grade <sup>d</sup>				
Well or moderately differentiated	20,600/239	1.20 (0.52–2.75)	0.67	
Poorly/undifferentiated	20,600/696	0.74 (0.45–1.21)	0.23	
$P_{\text{het}}$			0.33	
Meta-analysis method <sup>e</sup>				
All participants	22,588/2923	1.12 (0.86–1.46)	0.39	15.9%
<i>BRCA1</i> carrier	14,676/2319	1.18 (0.88–1.57)	0.26	17.2%
<i>BRCA2</i> carrier	7912/604	0.80 (0.45–1.43)	0.45	0.0%
$P_{\text{interaction}}$			0.24	
Two-stage residual inclusion method <sup>f</sup>				
All participants	7516/772	1.37 (0.84–2.24)	0.21	
<i>BRCA1</i> carrier	4401/543	1.24 (0.67–2.27)	0.49	
<i>BRCA2</i> carrier	3115/229	1.57 (0.67–3.66)	0.30	
<i>HR</i> hazard ratio, <i>CI</i> confidence interval, <i>CIMBA</i> Consortium of Investigators for the Modifiers of <i>BRCA1/2</i> <sup>a</sup> BMI-GS was constructed by combining 93 BMI-associated single-nucleotide polymorphisms (SNPs) <sup>b</sup> Adjusted for principal components, birth cohort, country of enrolment, and menopausal status in weighted Cox model <sup>c</sup> Adjusted for principal components, mutation status, birth cohort, and country of enrolment <sup>d</sup> Adjusted for principal components, mutation status, menopausal status, birth cohort, and country of enrolment <sup>e</sup> Hazard ratios were calculated using inverse-variance meta-analysis and re-scaled to the corresponding units by calculating the height measurements per z-score among controls. Effect estimates for ovarian cancer for each SNP were calculated from weighted Cox model adjusting for principal components, birth cohort, country of enrolment, menopausal status, and mutation status <sup>f</sup> Analysis was performed among 7516 participants with measured BMI Bolded lines refer to the model corresponding to our main results				

Table 5. Association of body mass index genetic score (BMI-GS) and ovarian cancer risk among 22,588 participants in CIMBA, per 5 kg/m<sup>2</sup> increase in genetically predicted BMI

rs ID	Chromosome	Position	Nearest gene	Reference allele in CIMBA	Effect allele in CIMBA	Effect allele frequency in CIMBA	Imputation quality <sup>a</sup>	Association with ovarian cancer in CIMBA		
								Log hazard ratio <sup>b</sup>	Standard error	P value
<b>Height</b>										
rs11049611	12	28600244	CCDC91	C	T	0.28	1	0.127	0.036	0.0004
rs6902771	6	152157881	ESR1	C	T	0.46	0.98	0.091	0.032	0.005
rs584828	17	38599230	IGFBP4	C	T	0.39	0.68	0.109	0.040	0.006
rs3817428	15	89415247	ACAN	C	G	0.22	0.51	0.144	0.053	0.006
rs7517682	1	103519589	COL11A1	G	A	0.56	0.98	0.085	0.033	0.009
rs12470505	2	219908369	CCDC108	T	G	0.10	0.97	-0.143	0.055	0.009
rs26024	5	127696022	FBN2	A	C	0.34	1	-0.087	0.034	0.011
rs13113518	4	56399648	CLOCK	T	C	0.37	0.99	0.081	0.033	0.014
rs7319045	13	92024574	GPC5	A	G	0.61	0.92	0.084	0.035	0.017
rs2044124	17	61845425	CCDC47	T	C	0.95	0.91	0.187	0.079	0.018
rs9309101	2	43629612	THADA	A	G	0.35	1	0.076	0.033	0.021
rs11867943	17	54229842	ANKFN1	A	T	0.11	0.96	0.118	0.051	0.022
rs12779328	10	12943973	CCDC3	C	T	0.30	0.94	-0.080	0.036	0.026
rs8073371	17	46096276	COP22	C	T	0.20	1.00	-0.095	0.043	0.029
rs2013265	8	24092500	ADAM28	C	T	0.22	0.62	0.104	0.047	0.029
rs11687941	2	242191410	HDLBP	C	G	0.26	0.96	-0.079	0.037	0.031
rs6838153	4	122720999	EXOSC9	A	G	0.33	0.99	-0.072	0.034	0.033
rs7112925	11	66826160	RHOD	C	T	0.36	0.95	-0.071	0.034	0.037
rs16942341	15	89388905	ACAN	C	T	0.03	0.60	0.255	0.123	0.039
rs6080830	20	17771113	BANF2	A	G	0.43	0.68	-0.080	0.039	0.041
rs867245	4	2218888	POLN	C	G	0.07	1.00	0.122	0.060	0.043
rs1155939	6	126866133	C6orf173	C	A	0.51	0.99	0.064	0.033	0.049
<b>BMI</b>										
rs16851483	3	141275436	RASA2	G	T	0.07	1	-0.203	0.068	0.003
rs2207139	6	50845490	TFAP2B	A	G	0.16	0.99	0.120	0.043	0.005
rs2033732	8	85079709	RALYL	T	C	0.75	0.72	-0.088	0.042	0.037
rs6804842	3	25106437	RARB	A	G	0.58	0.58	0.087	0.044	0.046
CIMBA Consortium of Investigators for the Modifiers of BRCA1/2										
<sup>a</sup> Imputation quality of 1 indicates genotyped SNPs										
<sup>b</sup> Per-allele association with ovarian cancer was adjusted for principal components, birth cohort, menopausal status, age at menopause, country of enrolment, and mutation status in weighted Cox models										

Table 6. Height or body mass index (BMI) single-nucleotide polymorphisms (SNPs) statistically significantly associated ( $P < 0.05$ ) with ovarian cancer risk in CIMBA

## **Additional Information**

### **Ethics approval and consent to participate**

The current work and all contributing studies in CIMBA received approval from the local institutional review board or ethics committee. Written informed consent was provided by all of the participants participating in each individual CIMBA study.

### **Conflict of interest**

Dr. Georg Pfeiler received honoraria and grant from Pfizer, Roche, Novartis, Accord, AstraZeneca, Amgen, Accord, and Lilly. Dr. Ritu Salani served on advisory board for Tesaro, Clovis, Astra Zeneca, Ethicon, and Genmab, and speaker's bureau for Tesaro, Genentech. Other authors declare no conflicts of interest.

### **Availability of data and materials**

Due to the sensitive nature of the data used in this study, data requests by researchers trained in maintaining human subject confidentiality may be directed to the corresponding author of this study.

### **Authors' contributions**

F.Q., M.A.R., O.I.O, G.C., T.R.R., and D.H. were responsible for study design. F.Q. and D.H. performed the data analysis and wrote the first draft of the manuscript. All authors provided critical feedback on the contents of the manuscript and approved the final version for submission.

### **Funding**

CIMBA: The CIMBA data management and data analysis were supported by Cancer Research – UK grants C12292/A20861, C12292/A11174. ACA is a Cancer Research -UK Senior Cancer Research Fellow. GCT and ABS are NHMRC Research Fellows. iCOGS: the European Community's Seventh Framework Programme under grant agreement No. 223175 (HEALTH-F2-2009-223175) (COGS), Cancer Research UK (C1287/A10118, C1287/A 10710, C12292/A11174, C1281/A12014, C5047/A8384, C5047/A15007, C5047/A10692, C8197/A16565), the National Institutes of Health (CA128978) and Post-Cancer GWAS initiative (1U19 CA148537, 1U19 CA148065 and 1U19 CA148112 - the GAME-ON initiative), the Department of Defence (W81XWH-10-1-0341), the Canadian Institutes of Health Research (CIHR) for the CIHR Team in Familial Risks of Breast Cancer (CRN-87521), and the Ministry of Economic Development, Innovation and Export Trade (PSR-SIIRI-701), Komen Foundation for the Cure, the Breast Cancer Research Foundation, and the Ovarian Cancer Research Fund. The PERSPECTIVE project was supported by the Government of Canada through Genome Canada and the Canadian Institutes of Health Research, the Ministry of Economy, Science and Innovation through Genome Québec, and The Quebec Breast Cancer Foundation. BCFR: UM1 CA164920 from the National Cancer Institute. The content of this manuscript does not necessarily reflect the views or policies of the National Cancer Institute or any of the collaborating centers in the Breast Cancer Family Registry (BCFR), nor does mention of trade

names, commercial products, or organizations imply endorsement by the US Government or the BCFR. BFBOCC: Lithuania (BFBOCC-LT): Research Council of Lithuania grant SEN-18/2015. BIDMC: Breast Cancer Research Foundation. BMBSA: Cancer Association of South Africa (PI Elizabeth J. van Rensburg). CNIO: Spanish Ministry of Health PI16/00440 supported by FEDER funds, the Spanish Ministry of Economy and Competitiveness (MINECO) SAF2014-57680-R and the Spanish Research Network on Rare diseases (CIBERER). COH-CCGCRN: Research reported in this publication was supported by the National Cancer Institute of the National Institutes of Health under grant number R25CA112486, and RC4CA153828 (PI: J. Weitzel) from the National Cancer Institute and the Office of the Director, National Institutes of Health. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. CONSTIT: Associazione Italiana Ricerca sul Cancro (AIRC; IG2014 no.15547) to P. Radice. Italian Association for Cancer Research (AIRC; grant no.16933) to L. Ottini. Associazione Italiana Ricerca sul Cancro (AIRC; IG2015 no.16732) to P. Peterlongo. Jacopo Azzollini is supported by funds from Italian citizens who allocated the 5x1000 share of their tax payment in support of the Fondazione IRCCS Istituto Nazionale Tumori, according to Italian laws (INT-Institutional strategic projects '5x1000'). DEMOKRITOS: European Union (European Social Fund – ESF) and Greek national funds through the Operational Program "Education and Lifelong Learning" of the National Strategic Reference Framework (NSRF) - Research Funding Program of the General Secretariat for Research & Technology: SYN11\_10\_19 NBCA. Investing in knowledge society through the European Social Fund. DFKZ: German Cancer Research Center. EMBRACE: Cancer Research UK Grants C1287/A10118 and C1287/A11990. D. Gareth Evans and Fiona Laloo are supported by an NIHR grant to the Biomedical Research Centre, Manchester. The Investigators at The Institute of Cancer Research and The Royal Marsden NHS Foundation Trust are supported by an NIHR grant to the Biomedical Research Centre at The Institute of Cancer Research and The Royal Marsden NHS Foundation Trust. Ros Eeles and Elizabeth Bancroft are supported by Cancer Research UK Grant C5047/A8385. Ros Eeles is also supported by NIHR support to the Biomedical Research Centre at The Institute of Cancer Research and The Royal Marsden NHS Foundation Trust. FCCC: The University of Kansas Cancer Center (P30 CA168524) and the Kansas Bioscience Authority Eminent Scholar Program. A.K.G. was funded by R01 CA140323, R01 CA214545, and by the Chancellors Distinguished Chair in Biomedical Sciences Professorship. FPGMX: FISPI05/2275 and Mutua Madrileña Foundation (FMMA). GC-HBOC: German Cancer Aid (grant no 110837, Rita K. Schmutzler) and the European Regional Development Fund and Free State of Saxony, Germany (LIFE - Leipzig Research Centre for Civilization Diseases, project numbers 713-241202, 713-241202, 14505/2470, 14575/2470). GEMO: Ligue Nationale Contre le Cancer; the Association "Le cancer du sein, parlons-en!" Award, the Canadian Institutes of Health Research for the "CIHR Team in Familial Risks of Breast Cancer" program and the French National Institute of Cancer (INCa grants 2013-1-BCB-01-ICH-1 and SHS-E-SP 18-015). GEORGETOWN: the Non-Therapeutic Subject Registry Shared Resource at Georgetown University (NIH/NCI grant P30-CA051008), the Fisher Center for Hereditary Cancer and Clinical Genomics Research, and Swing Fore the Cure. G-FAST: Bruce Poppe is a senior clinical investigator of FWO. Mattias Van Heetvelde obtained funding from IWT. HCSC: Spanish Ministry of Health PI15/00059, PI16/01292, and CB-161200301

CIBERONC from ISCIII (Spain), partially supported by European Regional Development FEDER funds. HEBCS: Helsinki University Hospital Research Fund, the Finnish Cancer Society and the Sigrid Juselius Foundation. HEBON: the Dutch Cancer Society grants NKI1998-1854, NKI2004-3088, NKI2007-3756, the Netherlands Organisation of Scientific Research grant NWO 91109024, the Pink Ribbon grants 110005 and 2014-187.WO76, the BBMRI grant NWO 184.021.007/CP46 and the Transcan grant JTC 2012 Cancer 12-054. HRBCP: Hong Kong Sanatorium and Hospital, Dr Ellen Li Charitable Foundation, The Kerry Group Kuok Foundation, National Institute of Health IR 03CA130065, and North California Cancer Center. HUNBOCS: Hungarian Research Grants KTIA-OTKA CK-80745 and OTKA K-112228. ICO: The authors would like to particularly acknowledge the support of the Asociación Española Contra el Cáncer (AECC), the Instituto de Salud Carlos III (organismo adscrito al Ministerio de Economía y Competitividad) and “Fondo Europeo de Desarrollo Regional (FEDER), una manera de hacer Europa” (PI10/01422, PI13/00285, PIE13/00022, PI15/00854, PI16/00563 and CIBERONC) and the Institut Català de la Salut and Autonomous Government of Catalonia (2009SGR290, 2014SGR338 and PERIS Project MedPerCan). IHCC: PBZ\_KBN\_122/P05/2004. ILUH: Icelandic Association “Walking for Breast Cancer Research” and by the Landspítali University Hospital Research Fund. INHERIT: Canadian Institutes of Health Research for the “CIHR Team in Familial Risks of Breast Cancer” program – grant # CRN-87521 and the Ministry of Economic Development, Innovation and Export Trade – grant # PSR-SIIRI-701. IOVHBOCS: Ministero della Salute and “5x1000” Istituto Oncologico Veneto grant. IPOBCS: Liga Portuguesa Contra o Cancro. kConFab: The National Breast Cancer Foundation, and previously by the National Health and Medical Research Council (NHMRC), the Queensland Cancer Fund, the Cancer Councils of New South Wales, Victoria, Tasmania and South Australia, and the Cancer Foundation of Western Australia. MAYO: NIH grants CA116167, CA192393 and CA176785, an NCI Specialized Program of Research Excellence (SPORE) in Breast Cancer (CA116201), and a grant from the Breast Cancer Research Foundation. MCGILL: Jewish General Hospital Weekend to End Breast Cancer, Quebec Ministry of Economic Development, Innovation and Export Trade. Marc Tischkowitz is supported by the funded by the European Union Seventh Framework Program (2007Y2013)/European Research Council (Grant No. 310018). MODSQUAD: MH CZ - DRO (MMCI, 00209805), MEYS - NPS I - LO1413 to LF and by the European Regional Development Fund and the State Budget of the Czech Republic (RECAMO, CZ.1.05/2.1.00/03.0101) to LF, and by Charles University in Prague project UNCE204024 (MZ). MSKCC: the Breast Cancer Research Foundation, the Robert and Kate Niehaus Clinical Cancer Genetics Initiative, the Andrew Sabin Research Fund and a Cancer Center Support Grant/Core Grant (P30 CA008748). NAROD: 1R01 CA149429-01. NCI: the Intramural Research Program of the US National Cancer Institute, NIH, and by support services contracts NO2-CP-11019-50, N02-CP-21013-63 and N02-CP-65504 with Westat, Inc, Rockville, MD. NICCC: Clalit Health Services in Israel, the Israel Cancer Association and the Breast Cancer Research Foundation (BCRF), NY. NNPIO: the Russian Foundation for Basic Research (grants 17-54-12007, 17-00-00171 and 18-515-12007). NRG Oncology: U10 CA180868, NRG SDMC grant U10 CA180822, NRG Administrative Office and the NRG Tissue Bank (CA 27469), the NRG Statistical and Data Center (CA 37517) and the Intramural Research Program, NCI. OSUCCG: Ohio State University Comprehensive Cancer Center. PBCS: Italian



Association of Cancer Research (AIRC) [IG 2013 N.14477] and Tuscany Institute for Tumors (ITT) grant 2014-2015-2016. SEABASS: Ministry of Science, Technology and Innovation, Ministry of Higher Education (UM.C/HIR/MOHE/06) and Cancer Research Initiatives Foundation. SMC: the Israeli Cancer Association. SWE-BRCA: the Swedish Cancer Society. UCHICAGO: NCI Specialized Program of Research Excellence (SPORE) in Breast Cancer (CA125183), R01 CA142996, 1U01CA161032, P20CA233307, American Cancer Society (MRSF-13-063-01-TBG, CRP-10-119-01-CCE), Breast Cancer Research Foundation, Susan G. Komen Foundation (SAC110026), and Ralph and Marion Falk Medical Research Trust, the Entertainment Industry Fund National Women's Cancer Research Alliance. Mr. Qian was supported by the Alpha Omega Alpha Carolyn L. Cuckein Student Research Fellowship. UCLA: Jonsson Comprehensive Cancer Center Foundation; Breast Cancer Research Foundation. UCSF: UCSF Cancer Risk Program and Helen Diller Family Comprehensive Cancer Center. UKFOCR: Cancer Research UK. UPENN: Breast Cancer Research Foundation; Susan G. Komen Foundation for the cure, Basser Center for BRCA. UPITT/MWH: Hackers for Hope Pittsburgh. VFCTG: Victorian Cancer Agency, Cancer Australia, National Breast Cancer Foundation. WCP: Dr Karlan is funded by the American Cancer Society Early Detection Professorship (SIOP-06-258-01-COUN) and the National Center for Advancing Translational Sciences (NCATS), Grant UL1TR000124.

## **Acknowledgements**

All the families and clinicians who contribute to the studies; Sue Healey, in particular taking on the task of mutation classification with the late Olga Sinilnikova; Maggie Angelakos, Judi Maskiell, Gillian Dite, Helen Tsimiklis; members and participants in the New York site of the Breast Cancer Family Registry; members and participants in the Ontario Familial Breast Cancer Registry; Vilius Rudaitis and Laimonas Griškevičius; Drs Janis Eglitis, Anna Krilova and Aivars Stengrevics; Yuan Chun Ding and Linda Steele for their work in participant enrollment and biospecimen and data management; Bent Ejlerlsen and Anne-Marie Gerdes for the recruitment and genetic counseling of participants; Alicia Barroso, Rosario Alonso and Guillermo Pita; Manoukian Siranoush, Bernard Peissel, Cristina Zanzottera, Milena Mariani, Daniela Zaffaroni, Bernardo Bonanni, Monica Barile, Irene Feroce, Mariarosaria Calvello, Alessandra Viel, Riccardo Dolcetti, Laura Ottini, Giuseppe Giannini, Laura Papi, Gabriele Lorenzo Capone, Liliana Varesco, Viviana Gismondi, Maria Grazia Tibiletti, Daniela Furlan, Antonella Savarese, Aline Martayan, Stefania Tommasi, Brunella Pilato; the personnel of the Cogentech Cancer Genetic Test Laboratory, Milan, Italy. Ms. JoEllen Weaver and Dr. Betsy Bove; Marta Santamariña, Ana Blanco, Miguel Aguado, Uxía Esperón and Belinda Rodríguez; IFE - Leipzig Research Centre for Civilization Diseases (Markus Loeffler, Joachim Thiery, Matthias Nüchter, Ronny Baber); We thank all participants, clinicians, family doctors, researchers, and technicians for their contributions and commitment to the DKFZ study. Genetic Modifiers of Cancer Risk in BRCA1/2 Mutation Carriers (GEMO) study is a study from the National Cancer Genetics Network UNICANCER Genetic Group, France. We wish to pay a tribute to Olga M. Sinilnikova, who with Dominique Stoppa-Lyonnet initiated and coordinated



GEMO until she sadly passed away on the 30th June 2014. The team in Lyon (Olga Sinilnikova, Mélanie Léoné, Laure Barjhoux, Carole Verny-Pierre, Sylvie Mazoyer, Francesca Damiola, Valérie Sornin) managed the GEMO samples until the biological resource centre was transferred to Paris in December 2015 (Noura Mebirouk, Fabienne Lesueur, Dominique Stoppa-Lyonnet). We want to thank all the GEMO collaborating groups for their contribution to this study: Coordinating Centre, Service de Génétique, Institut Curie, Paris, France: Muriel Belotti, Ophélie Bertrand, Anne-Marie Birot, Bruno Buecher, Sandrine Caputo, Anaïs Dupré, Emmanuelle Fourme, Marion Gauthier-Villars, Lisa Golmard, Claude Houdayer, Marine Le Mentec, Virginie Moncoutier, Antoine de Pauw, Claire Saule, Dominique Stoppa-Lyonnet, and Inserm U900, Institut Curie, Paris, France: Fabienne Lesueur, Noura Mebirouk. Contributing Centres : Unité Mixte de Génétique Constitutionnelle des Cancers Fréquents, Hospices Civils de Lyon - Centre Léon Bérard, Lyon, France: Nadia Boutry-Kryza, Alain Calender, Sophie Giraud, Mélanie Léoné. Institut Gustave Roussy, Villejuif, France: Brigitte Bressac-de-Paillerets, Olivier Caron, Marine Guillaud-Bataille. Centre Jean Perrin, Clermont-Ferrand, France: Yves-Jean Bignon, Nancy Uhrhammer. Centre Léon Bérard, Lyon, France: Valérie Bonadona, Christine Lasset. Centre François Baclesse, Caen, France: Pascaline Berthet, Laurent Castera, Dominique Vaur. Institut Paoli Calmettes, Marseille, France: Violaine Bourdon, Catherine Noguès, Tetsuro Noguchi, Cornel Popovici, Audrey Remenieras, Hagay Sobol. CHU Arnaud-de-Villeneuve, Montpellier, France: Isabelle Coupier, Pascal Pujol. Centre Oscar Lambret, Lille, France: Claude Adenis, Aurélie Dumont, Françoise Révillion. Centre Paul Strauss, Strasbourg, France: Danièle Muller. Institut Bergonié, Bordeaux, France: Emmanuelle Barouk-Simonet, Françoise Bonnet, Virginie Bubien, Michel Longy, Nicolas Sevenet, Institut Claudius Regaud, Toulouse, France: Laurence Gladieff, Rosine Guimbaud, Viviane Feillel, Christine Toulas. CHU Grenoble, France: Héléne Dreyfus, Christine Dominique Leroux, Magalie Peysselon, Rebischung. CHU Dijon, France: Amandine Baurand, Geoffrey Bertolone, Fanny Coron, Laurence Faivre, Caroline Jacquot, Sarab Lizard. CHU St-Etienne, France: Caroline Kientz, Marine Lebrun, Fabienne Prieur. Hôtel Dieu Centre Hospitalier, Chambéry, France: Sandra Fert Ferrer. Centre Antoine Lacassagne, Nice, France: Véronique Mari. CHU Limoges, France: Laurence Vénat-Bouvet. CHU Nantes, France: Stéphane Bézieau, Capucine Delnatte. CHU Bretonneau, Tours and Centre Hospitalier de Bourges France: Isabelle Mortemousque. Groupe Hospitalier Pitié-Salpêtrière, Paris, France: Chrystelle Colas, Florence Coulet, Florent Soubrier, Mathilde Warcoin. CHU Vandoeuvre-les-Nancy, France: Myriam Bronner, Johanna Sokolowska. CHU Besançon, France: Marie-Agnès Collonge-Rame, Alexandre Damette. CHU Poitiers, Centre Hospitalier d'Angoulême and Centre Hospitalier de Niort, France: Paul Gesta. Centre Hospitalier de La Rochelle : Hakima Lallaoui. CHU Nîmes Carémeau, France : Jean Chiesa. CHI Poissy, France: Denise Molina-Gomes. CHU Angers, France : Olivier Ingster; Ilse Coene en Brecht Crombez; Ilse Coene and Brecht Crombez; Alicia Tosar and Paula Diaque; Drs .Sofia Khan, Taru A. Muranen, Carl Blomqvist, Irja Erkkilä and Virpi Palola; The Hereditary Breast and Ovarian Cancer Research Group Netherlands (HEBON) consists of the following Collaborating Centers: Netherlands Cancer Institute (coordinating center), Amsterdam, NL: M.A. Rookus, F.B.L. Hogervorst, F.E. van Leeuwen, M.A. Adank, M.K. Schmidt, N.S. Russell, D.J. Jenner; Erasmus Medical Center, Rotterdam, NL: J.M. Collée, A.M.W. van den Ouweland, M.J. Hooning, C.M. Seynaeve, C.H.M. van Deurzen, I.M. Obdeijn, I.A. Boere; Leiden University Medical Center, NL: C.J. van Asperen, P. Devilee,

T.C.T.E.F. van Cronenburg; Radboud University Nijmegen Medical Center, NL: C.M. Kets, A.R. Mensenkamp; University Medical Center Utrecht, NL: M.G.E.M. Ausems, M.J. Koudijs; Amsterdam Medical Center, NL: C.M. Aalfs, H.E.J. Meijers-Heijboer; VU University Medical Center, Amsterdam, NL: K. van Engelen, J.J.P. Gille; Maastricht University Medical Center, NL: E.B. Gómez-García, M.J. Blok, M. de Boer; University of Groningen, NL: J.C. Oosterwijk, A.H. van der Hout, M.J. Mourits, G.H. de Bock; The Netherlands Comprehensive Cancer Organisation (IKNL): S. Siesling, J. Verloop; The nationwide network and registry of histo- and cytopathology in The Netherlands (PALGA): A.W. van den Belt-Dusebout. HEBON thanks the study participants and the registration teams of IKNL and PALGA for part of the data collection; Hong Kong Sanatorium and Hospital; the Hungarian Breast and Ovarian Cancer Study Group members (Janos Papp, Aniko Bozsik, Timea Pocza, Zoltan Matrai, Miklos Kasler, Judit Franko, Maria Balogh, Gabriella Domokos, Judit Ferenczi, Department of Molecular Genetics, National Institute of Oncology, Budapest, Hungary) and the clinicians and patients for their contributions to this study; the Oncogenetics Group (VHIO) and the High Risk and Cancer Prevention Unit of the University Hospital Vall d'Hebron, and the Cellex Foundation for providing research facilities and equipment; the ICO Hereditary Cancer Program team led by Dr. Gabriel Capella; the ICO Hereditary Cancer Program team led by Dr. Gabriel Capella; Dr Martine Dumont for sample management and skillful assistance; Ana Peixoto, Catarina Santos and Pedro Pinto; members of the Center of Molecular Diagnosis, Oncogenetics Department and Molecular Oncology Research Center of Barretos Cancer Hospital; Heather Thorne, Eveline Niedermayr, all the kConFab research nurses and staff, the heads and staff of the Family Cancer Clinics, and the Clinical Follow Up Study (which has received funding from the NHMRC, the National Breast Cancer Foundation, Cancer Australia, and the National Institute of Health (USA)) for their contributions to this resource, and the many families who contribute to kConFab; the KOBRA Study Group; Csilla Szabo (National Human Genome Research Institute, National Institutes of Health, Bethesda, MD, USA); Lenka Foretova and Eva Machackova (Department of Cancer Epidemiology and Genetics, Masaryk Memorial Cancer Institute and MF MU, Brno, Czech Republic); and Michal Zikan, Petr Pohlreich and Zdenek Kleibl (Oncogynecologic Center and Department of Biochemistry and Experimental Oncology, First Faculty of Medicine, Charles University, Prague, Czech Republic); Anne Lincoln, Lauren Jacobs; the NICCC National Familial Cancer Consultation Service team led by Sara Dishon, the lab team led by Dr. Flavio Lejbkowitz, and the research field operations team led by Dr. Mila Pinchev; the investigators of the Australia New Zealand NRG Oncology group; members and participants in the Ontario Cancer Genetics Network; Leigha Senter, Kevin Sweet, Caroline Craven, Julia Cooper, and Michelle O'Connor; Yip Cheng Har, Nur Aishah Mohd Taib, Phuah Sze Yee, Norhashimah Hassan and all the research nurses, research assistants and doctors involved in the MyBrCa Study for assistance in patient recruitment, data collection and sample preparation, Philip Iau, Sng Jen-Hwei and Sharifah Nor Akmal for contributing samples from the Singapore Breast Cancer Study and the HUKM-HKL Study respectively; the Meirav Comprehensive breast cancer center team at the Sheba Medical Center; Christina Selkirk; Åke Borg, Håkan Olsson, Helena Jernström, Karin Henriksson, Katja Harbst, Maria Soller, Ulf Kristoffersson; from Gothenburg Sahlgrenska University Hospital: Anna Öfverholm, Margareta Nordling, Per Karlsson, Zakaria Einbeigi; from Stockholm and Karolinska University Hospital: Anna von Wachenfeldt, Annelie

Liljegren, Annika Lindblom, Brita Arver, Gisela Barbany Bustinza, Johanna Rantala; from Umeå University Hospital: Beatrice Melin, Christina Edwinsdotter Ardnor, Monica Emanuelsson; from Uppsala University: Hans Ehrencrona, Maritta Hellström Pigg, Richard Rosenquist; from Linköping University Hospital: Marie Stenmark-Askmal, Sigrun Liedgren; Cecilia Zvocec, Qun Niu; Joyce Seldon and Lorna Kwan; Dr. Robert Nussbaum, Beth Crawford, Kate Loranger, Julie Mak, Nicola Stewart, Robin Lee, Amie Blanco and Peggy Conrad and Salina Chan; Simon Gayther, Susan Ramus, Paul Pharoah, Carole Pye, Patricia Harrington and Eva Wozniak; Geoffrey Lindeman, Marion Harris, Martin Delatycki, Sarah Sawyer, Rebecca Driessen, and Ella Thompson for performing all DNA amplification.

## References

1. Salehi F, Dunfield L, Phillips KP, Krewski D, Vanderhyden BC. Risk factors for ovarian cancer: an overview with emphasis on hormonal factors. *J Toxicol Environ Health B Crit Rev* 2008; **11**(3-4): 301-321; doi 10.1080/10937400701876095.
2. Wentzensen N, Poole EM, Trabert B, White E, Arslan AA, Patel AV *et al.* Ovarian Cancer Risk Factors by Histologic Subtype: An Analysis From the Ovarian Cancer Cohort Consortium. *J Clin Oncol* 2016; **34**(24): 2888-2898; doi 10.1200/JCO.2016.66.8178.
3. Rauh-Hain JA, Krivak TC, Del Carmen MG, Olawaiye AB. Ovarian cancer screening and early detection in the general population. *Rev Obstet Gynecol* 2011; **4**(1): 15-21.
4. Jacobs IJ, Menon U, Ryan A, Gentry-Maharaj A, Burnell M, Kalsi JK *et al.* Ovarian cancer screening and mortality in the UK Collaborative Trial of Ovarian Cancer Screening (UKCTOCS): a randomised controlled trial. *Lancet* 2016; **387**(10022): 945-956; doi 10.1016/S0140-6736(15)01224-6.
5. Neff RT, Senter L, Salani R. BRCA mutation in ovarian cancer: testing, implications and treatment considerations. *Ther Adv Med Oncol* 2017; **9**(8): 519-531; doi 10.1177/1758834017714993.
6. King MC, Marks JH, Mandell JB, New York Breast Cancer Study G. Breast and ovarian cancer risks due to inherited mutations in BRCA1 and BRCA2. *Science* (Research Support, U.S. Gov't, Non-P.H.S.) 2003; **302**(5645): 643-646; e-pub ahead of print 2003/10/25; doi 10.1126/science.1088759.
7. Kuchenbaecker KB, Hopper JL, Barnes DR, Phillips KA, Mooij TM, Roos-Blom MJ *et al.* Risks of Breast, Ovarian, and Contralateral Breast Cancer for BRCA1 and BRCA2 Mutation Carriers. *JAMA* 2017; **317**(23): 2402-2416; doi 10.1001/jama.2017.7112.
8. Friebel TM, Domchek SM, Rebbeck TR. Modifiers of cancer risk in BRCA1 and BRCA2 mutation carriers: systematic review and meta-analysis. *J Natl Cancer Inst* (Meta-Analysis Research Support, N.I.H., Extramural Research Support, Non-U.S. Gov't Review) 2014; **106**(6): dju091; e-pub ahead of print 2014/05/16; doi 10.1093/jnci/dju091.
9. Gayther SA, Russell P, Harrington P, Antoniou AC, Easton DF, Ponder BA. The contribution of germline BRCA1 and BRCA2 mutations to familial ovarian cancer: no evidence for other ovarian cancer-susceptibility genes. *Am J Hum Genet* 1999; **65**(4): 1021-1029; doi 10.1086/302583.
10. Lango Allen H, Estrada K, Lettre G, Berndt SI, Weedon MN, Rivadeneira F *et al.* Hundreds of variants clustered in genomic loci and biological pathways affect human height. *Nature* 2010; **467**(7317): 832-838; doi 10.1038/nature09410.
11. Locke AE, Kahali B, Berndt SI, Justice AE, Pers TH, Day FR *et al.* Genetic studies of body mass index yield new insights for obesity biology. *Nature* (Meta-Analysis) 2015; **518**(7538): 197-206; e-pub ahead of print 2015/02/13; doi 10.1038/nature14177.
12. Collaborative Group on Epidemiological Studies of Ovarian C. Ovarian cancer and body size: individual participant meta-analysis including 25,157 women with ovarian cancer from 47 epidemiological studies. *PLoS Med* 2012; **9**(4): e1001200; doi 10.1371/journal.pmed.1001200.

13. Beehler GP, Sekhon M, Baker JA, Teter BE, McCann SE, Rodabaugh KJ *et al.* Risk of ovarian cancer associated with BMI varies by menopausal status. *J Nutr* 2006; **136**(11): 2881-2886.
14. Dixon SC, Nagle CM, Thrift AP, Pharoah PD, Pearce CL, Zheng W *et al.* Adult body mass index and risk of ovarian cancer by subtype: a Mendelian randomization study. *Int J Epidemiol* 2016; **45**(3): 884-895; doi 10.1093/ije/dyw158.
15. Bhaskaran K, Douglas I, Forbes H, dos-Santos-Silva I, Leon DA, Smeeth L. Body-mass index and risk of 22 specific cancers: a population-based cohort study of 5.24 million UK adults. *Lancet* 2014; **384**(9945): 755-765; doi 10.1016/S0140-6736(14)60892-8.
16. Olsen CM, Nagle CM, Whiteman DC, Ness R, Pearce CL, Pike MC *et al.* Obesity and risk of ovarian cancer subtypes: evidence from the Ovarian Cancer Association Consortium. *Endocr Relat Cancer* 2013; **20**(2): 251-262; doi 10.1530/ERC-12-0395.
17. Cunningham JM, Cicek MS, Larson NB, Davila J, Wang C, Larson MC *et al.* Clinical characteristics of ovarian cancer classified by BRCA1, BRCA2, and RAD51C status. *Sci Rep* 2014; **4**: 4026; doi 10.1038/srep04026.
18. McGee J, Kotsopoulos J, Lubinski J, Lynch HT, Rosen B, Tung N *et al.* Anthropometric measures and risk of ovarian cancer among BRCA1 and BRCA2 mutation carriers. *Obesity (Silver Spring)* 2012; **20**(6): 1288-1292; doi 10.1038/oby.2011.394.
19. Burgess S, Butterworth A, Thompson SG. Mendelian randomization analysis with multiple genetic variants using summarized data. *Genet Epidemiol* 2013; **37**(7): 658-665; doi 10.1002/gepi.21758.
20. Paternoster L, Tilling K, Davey Smith G. Genetic epidemiology and Mendelian randomization for informing disease therapeutics: Conceptual and methodological challenges. *PLoS Genet* 2017; **13**(10): e1006944; doi 10.1371/journal.pgen.1006944.
21. VanderWeele TJ, Tchetgen Tchetgen EJ, Cornelis M, Kraft P. Methodological challenges in mendelian randomization. *Epidemiology* 2014; **25**(3): 427-435; doi 10.1097/EDE.0000000000000081.
22. Couch FJ, Wang X, McGuffog L, Lee A, Olswold C, Kuchenbaecker KB *et al.* Genome-wide association study in BRCA1 mutation carriers identifies novel loci associated with breast and ovarian cancer risk. *PLoS Genet* 2013; **9**(3): e1003212; doi 10.1371/journal.pgen.1003212.
23. Rebbeck TR, Mitra N, Wan F, Sinilnikova OM, Healey S, McGuffog L *et al.* Association of type and location of BRCA1 and BRCA2 mutations with risk of breast and ovarian cancer. *JAMA* 2015; **313**(13): 1347-1361; doi 10.1001/jama.2014.5985.
24. Qian F, Wang S, Mitchell J, McGuffog L, Barrowdale D, Leslie G *et al.* Height and Body Mass Index as Modifiers of Breast Cancer Risk in BRCA1/2 Mutation Carriers: A Mendelian Randomization Study. *J Natl Cancer Inst* 2018; doi 10.1093/jnci/djy132.
25. Wood AR, Esko T, Yang J, Vedantam S, Pers TH, Gustafsson S *et al.* Defining the role of common variation in the genomic and biological architecture of adult human height. *Nat Genet* 2014; **46**(11): 1173-1186; doi 10.1038/ng.3097.
26. Antoniou A, Pharoah PD, Narod S, Risch HA, Eyfjord JE, Hopper JL *et al.* Average risks of breast and ovarian cancer associated with BRCA1 or BRCA2 mutations detected in case Series unselected for family history: a combined analysis of 22 studies. *Am J Hum Genet* 2003; **72**(5): 1117-1130; doi 10.1086/375033.
27. Antoniou AC, Goldgar DE, Andrieu N, Chang-Claude J, Brohet R, Rookus MA *et al.* A weighted cohort approach for analysing factors modifying disease risks in carriers of

- high-risk susceptibility genes. *Genet Epidemiol* 2005; **29**(1): 1-11; doi 10.1002/gepi.20074.
28. Bautista LE, Smeeth L, Hingorani AD, Casas JP. Estimation of bias in nongenetic observational studies using "mendelian triangulation". *Ann Epidemiol* 2006; **16**(9): 675-680; doi 10.1016/j.annepidem.2006.02.001.
  29. Bowden J, Davey Smith G, Burgess S. Mendelian randomization with invalid instruments: effect estimation and bias detection through Egger regression. *Int J Epidemiol* 2015; **44**(2): 512-525; doi 10.1093/ije/dyv080.
  30. Terza JV, Basu A, Rathouz PJ. Two-stage residual inclusion estimation: addressing endogeneity in health econometric modeling. *J Health Econ* 2008; **27**(3): 531-543; doi 10.1016/j.jhealeco.2007.09.009.
  31. Ramus SJ, Gayther SA. The contribution of BRCA1 and BRCA2 to ovarian cancer. *Mol Oncol* 2009; **3**(2): 138-150; doi 10.1016/j.molonc.2009.02.001.
  32. Jones MR, Kamara D, Karlan BY, Pharoah PDP, Gayther SA. Genetic epidemiology of ovarian cancer and prospects for polygenic risk prediction. *Gynecol Oncol* 2017; **147**(3): 705-713; doi 10.1016/j.ygyno.2017.10.001.
  33. Reeves GK, Pirie K, Beral V, Green J, Spencer E, Bull D *et al.* Cancer incidence and mortality in relation to body mass index in the Million Women Study: cohort study. *BMJ* 2007; **335**(7630): 1134; doi 10.1136/bmj.39367.495995.AE.
  34. Gao C, Patel CJ, Michailidou K, Peters U, Gong J, Schildkraut J *et al.* Mendelian randomization study of adiposity-related traits and risk of breast, ovarian, prostate, lung and colorectal cancer. *Int J Epidemiol* 2016; **45**(3): 896-908; doi 10.1093/ije/dyw129.
  35. Liu Z, Zhang TT, Zhao JJ, Qi SF, Du P, Liu DW *et al.* The association between overweight, obesity and ovarian cancer: a meta-analysis. *Jpn J Clin Oncol* 2015; **45**(12): 1107-1115; doi 10.1093/jjco/hyv150.
  36. Schouten LJ, Rivera C, Hunter DJ, Spiegelman D, Adami HO, Arslan A *et al.* Height, body mass index, and ovarian cancer: a pooled analysis of 12 cohort studies. *Cancer Epidemiol Biomarkers Prev* 2008; **17**(4): 902-912; doi 10.1158/1055-9965.EPI-07-2524.
  37. Lahmann PH, Cust AE, Friedenreich CM, Schulz M, Lukanova A, Kaaks R *et al.* Anthropometric measures and epithelial ovarian cancer risk in the European Prospective Investigation into Cancer and Nutrition. *Int J Cancer* 2010; **126**(10): 2404-2415; doi 10.1002/ijc.24952.
  38. Cook LS, Meisner ALW, Weiss NS. Endometrial Cancer. In: Michael J. Thun MSL, James R. Cerhan, Christopher A. Haiman, David Schottenfeld (ed.) *Cancer Epidemiology and Prevention*, 4th edn. Oxford University Press 2018.
  39. Dixon-Suen SC, Nagle CM, Thrift AP, Pharoah PDP, Ewing A, Pearce CL *et al.* Adult height is associated with increased risk of ovarian cancer: a Mendelian randomisation study. *Br J Cancer* 2018; **118**(8): 1123-1129; doi 10.1038/s41416-018-0011-3.
  40. Schildkraut JM, Schwingl PJ, Bastos E, Evanoff A, Hughes C. Epithelial ovarian cancer risk among women with polycystic ovary syndrome. *Obstet Gynecol* 1996; **88**(4 Pt 1): 554-559.
  41. Tworoger SS, Fairfield KM, Colditz GA, Rosner BA, Hankinson SE. Association of oral contraceptive use, other contraceptive methods, and infertility with ovarian cancer risk. *Am J Epidemiol* 2007; **166**(8): 894-901; doi 10.1093/aje/kwm157.

42. Yin W, Falconer H, Yin L, Xu L, Ye W. Association Between Polycystic Ovary Syndrome and Cancer Risk. *JAMA oncology* 2018; e-pub ahead of print 2018/11/30; doi 10.1001/jamaoncol.2018.5188.
43. Oktay K, Kim JY, Barad D, Babayev SN. Association of BRCA1 mutations with occult primary ovarian insufficiency: a possible explanation for the link between infertility and breast/ovarian cancer risks. *J Clin Oncol* 2010; **28**(2): 240-244; doi 10.1200/JCO.2009.24.2057.
44. Iyengar NM, Gucaip A, Dannenberg AJ, Hudis CA. Obesity and Cancer Mechanisms: Tumor Microenvironment and Inflammation. *J Clin Oncol* 2016; **34**(35): 4270-4276; doi 10.1200/JCO.2016.67.4283.
45. Eliassen AH, Hankinson SE. Endogenous hormone levels and risk of breast, endometrial and ovarian cancers: prospective studies. *Adv Exp Med Biol* 2008; **630**: 148-165.
46. Risch HA. Hormonal etiology of epithelial ovarian cancer, with a hypothesis concerning the role of androgens and progesterone. *J Natl Cancer Inst* 1998; **90**(23): 1774-1786.
47. Widschwendter M, Rosenthal AN, Philpott S, Rizzuto I, Fraser L, Hayward J *et al.* The sex hormone system in carriers of BRCA1/2 mutations: a case-control study. *Lancet Oncol* 2013; **14**(12): 1226-1232; doi 10.1016/S1470-2045(13)70448-0.
48. Jain A, Polotsky AJ, Rochester D, Berga SL, Loucks T, Zeitlian G *et al.* Pulsatile luteinizing hormone amplitude and progesterone metabolite excretion are reduced in obese women. *J Clin Endocrinol Metab* 2007; **92**(7): 2468-2473; doi 10.1210/jc.2006-2274.
49. Bu SZ, Yin DL, Ren XH, Jiang LZ, Wu ZJ, Gao QR *et al.* Progesterone induces apoptosis and up-regulation of p53 expression in human ovarian carcinoma cell lines. *Cancer* 1997; **79**(10): 1944-1950.
50. Schildkraut JM, Calingaert B, Marchbanks PA, Moorman PG, Rodriguez GC. Impact of progestin and estrogen potency in oral contraceptives on ovarian cancer risk. *J Natl Cancer Inst* 2002; **94**(1): 32-38.
51. Rodriguez GC, Nagarsheth NP, Lee KL, Bentley RC, Walmer DK, Cline M *et al.* Progestin-induced apoptosis in the Macaque ovarian epithelium: differential regulation of transforming growth factor-beta. *J Natl Cancer Inst* 2002; **94**(1): 50-60.
52. Crowe FL, Key TJ, Allen NE, Appleby PN, Overvad K, Gronbaek H *et al.* A cross-sectional analysis of the associations between adult height, BMI and serum concentrations of IGF-I and IGFBP-1 -2 and -3 in the European Prospective Investigation into Cancer and Nutrition (EPIC). *Ann Hum Biol* 2011; **38**(2): 194-202; doi 10.3109/03014460.2010.507221.
53. Beauchamp MC, Yasmeen A, Knafo A, Gotlieb WH. Targeting insulin and insulin-like growth factor pathways in epithelial ovarian cancer. *J Oncol* 2010; **2010**: 257058; doi 10.1155/2010/257058.
54. Pollak M. The insulin and insulin-like growth factor receptor family in neoplasia: an update. *Nat Rev Cancer* 2012; **12**(3): 159-169; doi 10.1038/nrc3215.
55. Lukanova A, Lundin E, Toniolo P, Micheli A, Akhmedkhanov A, Rinaldi S *et al.* Circulating levels of insulin-like growth factor-I and risk of ovarian cancer. *Int J Cancer* 2002; **101**(6): 549-554; doi 10.1002/ijc.10613.
56. Werner H, Bruchim I. IGF-1 and BRCA1 signalling pathways in familial cancer. *Lancet Oncol* 2012; **13**(12): e537-544; doi 10.1016/S1470-2045(12)70362-5.

57. Yengo L, Sidorenko J, Kemper KE, Zheng Z, Wood AR, Weedon MN *et al.* Meta-analysis of genome-wide association studies for height and body mass index in approximately 700000 individuals of European ancestry. *Hum Mol Genet* 2018; **27**(20): 3641-3649; doi 10.1093/hmg/ddy271.
58. Visscher PM, Wray NR, Zhang Q, Sklar P, McCarthy MI, Brown MA *et al.* 10 Years of GWAS Discovery: Biology, Function, and Translation. *Am J Hum Genet* 2017; **101**(1): 5-22; doi 10.1016/j.ajhg.2017.06.005.