


Genetic variation in *ST6GAL1* is a determinant of capecitabine and oxaliplatin induced hand-foot syndrome

Katie Watts¹ | Christopher Wills¹ | Ayman Madi² | Claire Palles³ | Timothy S. Maughan⁴ | Richard Kaplan⁵ | Nada A. Al-Tassan⁶ | Rachel Kerr⁷ | David J. Kerr⁸ | Richard S. Houlston⁹ | Valentina Escott-Price¹⁰ | Jeremy P. Cheadle¹ 

¹Division of Cancer and Genetics, School of Medicine, Cardiff University, Cardiff, UK

²The Clatterbridge Cancer Centre NHS Foundation Trust, Wirral, UK

³Institute of Cancer and Genomic Sciences, Institute of Biomedical Research, University of Birmingham, Birmingham, UK

⁴CRUK/MRC Oxford Institute for Radiation Oncology, University of Oxford, Oxford, UK

⁵MRC Clinical Trials Unit, University College of London, London, UK

⁶Department of Genetics, King Faisal Specialist Hospital and Research Center, Riyadh, Saudi Arabia

⁷Department of Oncology, Old Road Campus Research Building, University of Oxford, Oxford, UK

⁸Nuffield Department of Clinical Laboratory Sciences, University of Oxford, John Radcliffe Hospital, Oxford, UK

⁹Division of Genetics and Epidemiology, The Institute of Cancer Research, London, UK

¹⁰Institute of Psychological Medicine and Clinical Neurosciences, School of Medicine, Cardiff University, Cardiff, UK

Correspondence

Jeremy P. Cheadle, Division of Cancer and Genetics, School of Medicine, Cardiff University, Heath Park, Cardiff, CF14 4XN, UK.

Email: cheadlejp@cardiff.ac.uk

Funding information

This work was supported by Cardiff University School of Medicine (to KW), Tenovus Cancer Care (to JPC and CW), KFSHRC (to NAA) and

Abstract

Cancer patients treated with capecitabine and oxaliplatin (XELOX) often develop hand-foot syndrome (HFS) or palmar-plantar erythrodysesthesia. Genetic variation in *ST6GAL1* is a risk factor for type-2 diabetes (T2D), a disease also associated with HFS. We analysed genome-wide association data for 10 toxicities in advanced colorectal cancer (CRC) patients from the COIN and COIN-B trials. One thousand and fifty-five patients were treated with XELOX ± cetuximab and 745 with folinic acid, fluorouracil and oxaliplatin ± cetuximab. We also analysed rs6783836 in *ST6GAL1* with HFS in CRC patients from QUASAR2. Using UK Biobank data, we sought to confirm an association between *ST6GAL1* and T2D (17 384 cases, 317 887 controls) and analysed rs6783836 against markers of diabetes, inflammation and psoriasis. We found that 68% of patients from COIN and COIN-B with grade 2-3 HFS responded to treatment as compared to 58% with grade 0-1 HFS (odds ratio [OR] = 1.1, 95% confidence interval [CI] = 1.02-1.2, $P = 2.0 \times 10^{-4}$). HFS was also associated with improved overall survival (hazard ratio = 0.92, 95% CI = 0.84-0.99, $P = 4.6 \times 10^{-2}$). rs6783836 at *ST6GAL1* was associated with HFS in patients treated with XELOX (OR = 3.1, 95% CI = 2.1-4.6, $P = 4.3 \times 10^{-8}$) and was borderline significant in patients receiving capecitabine from QUASAR2, but with an opposite allele effect (OR = 0.66, 95% CI = 0.42-1.03, $P = .05$). *ST6GAL1* was associated with T2D (lead SNP rs3887925, OR = 0.94, 95% CI = 0.92-0.96, $P = 1.2 \times 10^{-8}$) and the rs6783836-T allele was associated with lowered HbA1c levels ($P = 5.9 \times 10^{-3}$) and lymphocyte count ($P = 2.7 \times 10^{-3}$), and psoriasis ($P = 7.5 \times 10^{-3}$) beyond thresholds for multiple testing. In conclusion, HFS is a biomarker of treatment outcome and rs6783836 in *ST6GAL1* is a potential biomarker for HFS with links to T2D and inflammation.

KEYWORDS

chemotherapy, colorectal cancer, genetics, *ST6GAL1*, toxicity, XELOX

Abbreviations: CRC, colorectal cancer; CTCAE, Common Terminology Criteria for Adverse Events; eQTL, expression quantitative trait loci; GTEx, The Genotype-Tissue Expression project database; GWAS, Genome-Wide Association Study; QC, quality control; QUASAR2, QUick And Simple And Reliable trial; SNP, single nucleotide polymorphism.

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2022 The Authors. *International Journal of Cancer* published by John Wiley & Sons Ltd on behalf of UICC.

Cancer Research UK (grant number C1298/A8362 to RSH). The COIN and COIN-B trials were funded by Cancer Research UK, the Medical Research Council and an unrestricted educational grant from Merck-Serono and were conducted with the support of the National Institute of Health Research Cancer Research Network. None of the sponsors played a role in the study design; the collection, analysis, and interpretation of data; the writing of the report; and the decision to submit the article for publication.

What's new?

Combination therapy with capecitabine and oxaliplatin (XELOX) in cancer patients can lead to hand-foot syndrome (HFS), risk factors for which include older age, being female, and having pre-existing conditions, such as diabetes and neuropathy. Here, investigation of HFS in XELOX-treated patients with advanced colorectal cancer (CRC) shows that HFS toxicity is associated with improvements in treatment response and overall survival. In particular, the variant rs6783836 in *ST6GAL1*, a gene with roles in inflammation and type-2 diabetes, was linked to HFS and glycated haemoglobin levels. The findings identify rs6783836 as a promising HFS biomarker in XELOX-treated patients with advanced CRC.

1 | INTRODUCTION

Toxicity from chemotherapy may result in treatment discontinuation or dose reduction affecting the prospect of a cure in patients with cancer. Patients treated with capecitabine and oxaliplatin (XELOX) often develop hand-foot syndrome (HFS), in which small amounts of the chemotherapeutic agent leaks out of capillaries into the hands and feet and damages the surrounding tissues.¹ HFS has been suggested to be a biomarker of treatment efficacy with post hoc analyses from clinical trials of colorectal and breast cancer patients finding that grade 1+ HFS was associated with improved overall and progression-free survival.^{2,3} Established risk factors for HFS include being older, female, having pre-existing peripheral neuropathy, circulation problems and diabetes.^{4,5}

Genetic variation in *ST6* β -galactoside α -2,6-sialyltransferase 1 (*ST6GAL1*) is associated with risk of developing type-2 diabetes (T2D).^{6,7} *ST6GAL1* catalyses the addition of α 2,6-linked sialic acids onto key surface glycoproteins. Increases in α 2,6-linked sialic acids have been linked to inflammatory conditions⁸ and *ST6GAL1* deficiency leads to increased inflammatory cell production,⁹ granulocyte recruitment¹⁰ and cytokine release.¹¹ There is also substantial evidence that *ST6GAL1* plays an important role in cancer progression, and is overexpressed in numerous cancers including colorectal.¹² High *ST6GAL1* expression has been associated with radioresistance and chemoresistance to several anticancer treatments, which ultimately leads to poorer patient outcomes.¹³⁻¹⁶

Inherited genetic factors are being recognised to affect toxicity from chemotherapeutic agents; notably, rare variants in the dihydropyrimidine dehydrogenase gene are associated with 5-fluorouracil toxicity. We have previously studied the relationship between common single nucleotide polymorphisms (SNPs) and 10 of the major toxicities, in patients with advanced colorectal cancer (CRC) from the COIN and COIN-B¹⁷⁻¹⁹ clinical trials.²⁰ Here, we extended our analysis and meta-analysed those patients receiving XELOX \pm cetuximab and, separately, folinic acid, fluorouracil and oxaliplatin (FOLFOX) \pm cetuximab. We also sought to confirm an association between *ST6GAL1* and T2D, and understand their interrelationship with HFS by studying biomarkers of inflammation and psoriasis using data from the UK Biobank.

2 | MATERIALS AND METHODS

2.1 | Patients and samples

Two thousand six hundred and seventy-one patients (mean age at randomisation of 62 years, range 18-87, 36% female) with metastatic or locally advanced CRC recruited into the MRC clinical trials COIN (ISRCTN27286448)^{17,18} and COIN-B (ISRCTN3837568)¹⁹ were studied. None of the patients had previously received chemotherapy for advanced disease. COIN patients were randomised 1:1:1 to receive continuous oxaliplatin and fluoropyrimidine chemotherapy (n = 815), continuous chemotherapy with cetuximab (n = 815), or intermittent chemotherapy (n = 815). COIN-B patients were randomised 1:1 to receive intermittent chemotherapy and cetuximab (n = 112) or intermittent chemotherapy and continuous cetuximab (n = 114). For the first 12-weeks, treatments were identical in all patients apart from the choice of fluoropyrimidine (n = 1603, 60% received XELOX and n = 1068, 40% received FOLFOX) together with the randomisation of \pm cetuximab (n = 1041, 39% received cetuximab). Blood DNA samples were prepared from 2244 of the 2671 patients.

2.2 | Toxicities assessed

Assessment of toxicities was performed at 12 weeks, since at this point patients from all trial arms received identical levels of chemotherapy with or without cetuximab. This time point was also prior to any interruption to treatment for the intermittent therapy arms. Toxicities assessed were diarrhoea, neutropenic sepsis, peripheral neuropathy, HFS, neutropenia, lethargy, stomatitis, nausea, vomiting and rash graded by critical adverse events as per the Common Terminology Criteria for Adverse Events (CTCAE version 4.0) with the highest grade noted within the first 12 weeks of treatment. Patients with toxicities graded 2-5 (G2-5) were grouped and compared against those graded 0-1 (G0-1). For HFS, we compared patients with G2-3 (G3 is the maximum) against those with G0-1; we also considered a linear model of toxicity.

2.3 | Patient outcome

Assessment of response was also performed at 12 weeks. Response was defined as complete or partial response using RECIST 1.0 guidelines and no response was defined as stable or progressive disease. Overall survival (OS) was defined as time from randomisation to death or date of last assessment.

2.4 | Genotyping

Two thousand two hundred and forty-four patient DNA samples were genotyped using Affymetrix Axiom Arrays according to the manufacturer's recommendations (Affymetrix, Santa Clara, California).²¹ After quality control (QC), SNP genotypes were available for 1950 patients.²¹ For 150 patients, no data on toxicity had been collected and these were excluded leaving 1800 for analysis. Prediction of untyped SNPs was carried out using IMPUTEv2 (v2.3.0) based on data from the 1000 Genomes Project as reference. We restricted our analysis to directly typed SNPs and imputed SNPs with INFO scores ≥ 0.8 , a Hardy-Weinberg equilibrium $\geq 1.0 \times 10^{-6}$ and a minor allele frequency (MAF) ≥ 0.05 .

2.5 | Statistical analyses

We previously analysed 4 million SNPs for a relationship with each toxicity under univariate models in patients that received XELOX (n = 707), XELOX + cetuximab (n = 348), FOLFOX (n = 385) and FOLFOX + cetuximab (n = 360).²⁰ Here, we incorporated covariates associated at $P < .05$ (Table S1) into the additive logistic models in Plink v1.9²² and meta-analysed those patients receiving XELOX \pm cetuximab (n = 1055) and, separately, FOLFOX \pm cetuximab (n = 745). Meta-analyses were run under a random effects model to account for the effect of cetuximab on toxicity, and results plotted in R studio using qqman.²³ SNPs associated at genome-wide significance ($P < 5.0 \times 10^{-8}$) were selected for further analyses. For survival analyses, cox proportional hazard regression models were used for both univariate and multivariate analyses. Results are reported in accordance with STREGA guidelines.²⁴

MAGMA²⁵ was used for gene and gene set analyses using data files from the NCBI 37.3 gene definitions and ~8500 predefined gene sets. Gene analyses were run under a SNP-wise univariate model imposing a Bonferroni corrected significance threshold of $P = 2.5 \times 10^{-6}$. Gene set analyses were run under competitive models with a corrected significance threshold of $P = 5.8 \times 10^{-6}$.

Power to detect toxicity effect sizes was calculated using the genpwr package in R,²⁶ based upon 70% power, $P = 5.0 \times 10^{-8}$ and SNPs with MAFs = 0.20; under these conditions we could identify SNPs with a mean OR of 2.8 (range 2-4 dependent upon toxicity, Table 1).

TABLE 1 Patients with grade 2-5 CTCAE toxicities at 12 weeks and detectable odds ratios at 70% power

Toxicity	Frequency		Detectable odds ratio	
	XELOX \pm cetuximab n (%)	FOLFOX \pm cetuximab n (%)	XELOX \pm cetuximab	FOLFOX \pm cetuximab
Diarrhoea	288 (27)	187 (25)	2	3
Neutropenic sepsis	6 (1)	63 (8)	NA	4
Peripheral neuropathy	154 (15)	73 (10)	2	3
Hand-foot syndrome	109 (10)	65 (9)	3	4
Neutropenia	42 (4)	209 (28)	4	2
Lethargy	361 (34)	256 (34)	2	2
Stomatitis	61 (6)	150 (20)	4	3
Nausea	210 (20)	88 (12)	2	3
Vomiting	122 (12)	59 (8)	3	4
Rash	177 (17)	201 (27)	2	2

Note: Percentage of patients in parentheses. NA—for neutropenic sepsis in patients treated with XELOX \pm cetuximab we had insufficient power to perform the genome-wide association study. Patients with hand-foot syndrome were graded 2-3.

2.6 | Independent replication

We attempted to replicate the association of rs6783836 with HFS using data from 927 patients with stage II or III CRCs enrolled in the Quick and Simple and Reliable trial (QUASAR2) comparing capecitabine or capecitabine plus bevacizumab.²⁷ Patients were genotyped using the Illumina genome-wide SNP panels (Human Hap 370, Human Hap 610 or Human Omni 2.5). Imputation was performed using IMPUTEv2 with 1000 genomes as reference. The INFO score for rs6783836 was 0.89. HFS was graded using the CTCAE scale and patients with G2-3 (46%) were compared to those with G0-1. Age was used as a covariate.

2.7 | ST6GAL1 variants and T2D

Six hundred and fourteen SNPs spanning ST6GAL1 were tested for an association with T2D in UK Biobank participants under project application number 65833 (17 384 cases and 317 887 controls as of 1 January 2021). We restricted our analysis to directly typed SNPs and imputed SNPs with INFO scores ≥ 0.8 , a Hardy-Weinberg equilibrium $\geq 1.0 \times 10^{-6}$ and a MAF ≥ 0.01 . We also analysed the relationship between rs6783836 and diabetic skin lesions by logistic regression on 617 diabetic individuals with self-reported open sores and 6605 diabetic controls (as of 1 July 2021).

2.8 | Potential biomarkers of HFS

We analysed rs6783836 and potential biomarkers of HFS using participant data from the UK Biobank. We assessed seven markers of

wound healing and/or inflammation: lymphocyte, neutrophil, monocyte, eosinophil, platelet and basophil counts (10^9 cells/L) and C-reactive protein levels (mg/L), and one marker for diabetes: glycated haemoglobin (HbA1c) levels (mmol/mol). Analyses were run using PHESANT.²⁸ Lymphocyte count, HbA1c levels, platelet count, neutrophil count, c-reactive protein levels and monocyte count were analysed under a linear regression and, basophil count and eosinophil count were analysed under an ordered logistic model as software default due to limited variation in the data. Results were held to a significance threshold of $P = 6.3 \times 10^{-3}$ (Bonferroni correction for eight

tests, $P = .05/8$). We analysed rs6783836 as a potential regulator of inflammation by performing a univariate logistic regression on 4228 individuals from the UK Biobank with self-reported psoriasis and 331 043 controls (as of 1 January 2021).

2.9 | Additional bioinformatic analyses

The Genotype-Tissue Expression (GTEx) project database was used to identify expression quantitative trait loci (eQTLs) for relevant SNPs

TABLE 2 Relationship between hand-foot syndrome (HFS) and patient outcome in COIN and COIN-B

Model	Grade of HFS (n)	Response at 12 weeks				Overall survival			
		% Responders	OR	95% CI	P (multivariate)	Median survival (days)	HR	95% CI	P (multivariate)
Grouped	0-1 (1626)	58	1.6	1.1-2.2	1.4×10^{-2} (2.0×10^{-2})	503	0.81	0.67-0.97	2.4×10^{-2} (0.15)
	2-3 (174)	68				596			
Linear	0 (1264)	56	1.3	1.2-1.6	1.4×10^{-4} (2.0×10^{-4})	499	0.90	0.83-0.97	5.8×10^{-3} (4.6×10^{-2})
	1 (362)	66				514			
	2 (144)	68				596			
	3 (30)	67				687			

Note: Response was defined as complete or partial response using RECIST 1.0 guidelines and no response was defined as stable or progressive disease. One thousand and eight hundred patients had data on overall survival and 1590 had data on response at 12 weeks. Covariates included in the multivariate analysis were age, sex, disease site, World Health Organisation performance status, primary tumour resection status, white blood cell count, chemotherapy regimen and cetuximab status.




FIGURE 1 Kaplan-Meier plot showing the relationship between hand-foot syndrome (HFS) and overall survival (OS). The y-axis represents survival probability and the x-axis represents time (days). The blue line represents patients with G0-1 HFS and the red line patients with G2-3 HFS. Dotted lines show the median OS (596 days in those with G2-3 HFS and 503 days in those with G0-1 HFS). The P-value was calculated using a cox proportional hazard regression [Color figure can be viewed at wileyonlinelibrary.com]




FIGURE 2 Regional plots for the association of rs6783836 with hand-foot syndrome (HFS). (A) Manhattan plot of the association between single-nucleotide polymorphism (SNP) genotype and HFS in patients treated with capecitabine and oxaliplatin. The red line corresponds to a $P = 5.0 \times 10^{-8}$ and the blue line $P = 1.0 \times 10^{-5}$. (B) Locus zoom plot shows results of the analysis for SNPs and recombination rates. $-\log_{10}(P)$ (y axis) of the SNPs are shown according to their chromosomal positions (x axis). The sentinel SNP (purple) is labelled by its rsID. The colour intensity of each symbol reflects the extent of linkage disequilibrium with the sentinel SNP, deep blue ($r^2 = 0$) through to dark red ($r^2 = 1.0$). Genetic recombination rates, estimated using 1000 Genomes Project samples, are shown with a blue line. Physical positions are based on NCBI build 37 of the human genome. Also shown are the relative positions of genes and transcripts mapping to the region of association. Genes have been redrawn to show their relative positions; therefore, maps are not to physical scale. Fine-mapping identified a credible set of 3 SNPs with rs6783836 having the highest posterior probability of 0.53 [Color figure can be viewed at wileyonlinelibrary.com]

(<https://gtexportal.org/home>). Fine-mapping was used for SNPs at significant loci; conditional regression was first used to identify the number of causal variants and fine-mapping was then run using PAINTOR.²⁹ Credible sets of causal SNPs were assembled for 95% coverage.

3 | RESULTS

3.1 | Relationship between HFS and patient outcome

Overall, 174/1800 (10%) patients from COIN and COIN-B developed G2-3 HFS at 12 weeks (109/1055, 10% in the XELOX group and

65/745, 9% in the FOLFOX group, Table 1). HFS was predictive of treatment outcome (Table 2). 105/154 (68%) patients with G2-3 HFS responded (had complete or partial response) to chemotherapy ± cetuximab at 12 weeks as compared to 831/1436 (58%) with G0-1 HFS (odds ratio [OR] = 1.6, 95% confidence intervals [CI] = 1.1-2.2, $P = 1.4 \times 10^{-2}$, univariate model). Under a multivariate model accounting for age, sex, disease site, World Health Organisation performance status, primary tumour resection status, white blood cell count, chemotherapy regimen and cetuximab status, this remained significant (OR = 1.1, 95% CI = 1.02-1.2, $P = 2.0 \times 10^{-2}$). Median OS was 596 days in those with G2-3 HFS and 503 days in those with G0-1 HFS (hazard ratio [HR] = 0.81, 95% CI = 0.67-0.97, $P = 2.4 \times 10^{-2}$, Figure 1); although, this did not remain significant under multivariate analysis ($P = .15$, Table 2). However, when HFS

TABLE 3 Relationship between rs6783836 and hand-foot syndrome (HFS) in patients from COIN and COIN-B treated with XELOX ± cetuximab

Treatment groups analysed	Total patients	Patients G0-1 HFS			Patients G2-3 HFS			OR	95% CI	P-value
		wild type	heterozygous	homozygous	wild type	heterozygous	homozygous			
Meta-analysis	1042	734	190	10	58	48	2	3.1	2.1-4.6	4.3×10^{-8}
<i>Subgroups:</i>										
XELOX	699	520	121	5	30	21	2	3.3	1.9-5.7	2.7×10^{-5}
XELOX + cetuximab	343	214	69	5	28	27	0	2.9	1.6-5.1	3.0×10^{-4}

Abbreviations: CI, confidence intervals; OR, odds ratio; T, reference allele.




FIGURE 3 Layered locus zoom plot showing single-nucleotide polymorphisms (SNPs) in *ST6GAL1* associated with hand-foot syndrome (HFS) and type-2 diabetes (T2D). Plot shows results of the analysis for SNPs and recombination rates. $-\log_{10}(P)$ (y axis) of the SNPs are shown according to their chromosomal positions (x axis). The dashed line corresponds to a $P = 5.0 \times 10^{-8}$. Genetic recombination rates, estimated using 1000 Genomes Project samples, are shown with a blue line. Physical positions are based on NCBI build 37 of the human genome. Lead SNPs for HFS and T2D are indicated by their rsIDs. Also shown is the relative coding region of *ST6GAL1* and chromatin state annotations from ENCODE [Color figure can be viewed at wileyonlinelibrary.com]




FIGURE 4 Relationship between rs6783836 and (A) continuous and (B) ordinal phenotypes. The x axis shows phenotype and respective UK Biobank ID, and the y axis shows SD change or odds ratio. Only lymphocyte count and glycated haemoglobin (HbA1c) were significantly associated with rs6783836 after Bonferroni correction for eight tests ($P < 6.3 \times 10^{-3}$) [Color figure can be viewed at wileyonlinelibrary.com]

was assessed as a linear trait, the relationship with OS was significant under such analyses (G0 median survival = 499 days, G1 = 514 days, G2 = 596 days, G3 = 687 days, HR = 0.92, 95% CI = 0.84-0.99, $P = 4.6 \times 10^{-2}$, Table 2). Cetuximab increased the frequency of HFS in patients treated with XELOX (56/348, 16% with and 53/707, 8% without cetuximab, $P = 1.5 \times 10^{-5}$) and FOLFOX (56/360, 16% with and 9/385, 2% without cetuximab, $P = 2.7 \times 10^{-8}$).

3.2 | Relationship between genetic variation at *ST6GAL1* and HFS

rs6783836 at 3q27.3 was associated with HFS at genome-wide significant levels in patients treated with XELOX (OR = 3.1, 95% CI = 2.1-4.6, $P = 4.3 \times 10^{-8}$, Figure 2). Forty-six percent (50/108) of patients with G2-3 HFS carried rs6783836 in a heterozygous or homozygotes state for the minor allele as compared to 21% (200/934) of patients with G0-1 HFS (Table 3). The association between rs6783836 and HFS was seen in patients treated with XELOX alone (OR = 3.3, 95% CI = 1.9-5.7, $P = 2.7 \times 10^{-5}$) and in those treated with XELOX + cetuximab (OR = 2.9, 95% CI = 1.6-5.1, $P = 3.0 \times 10^{-4}$, Table 3); cetuximab did not affect this relationship ($P_{\text{interaction}} = 0.98$). rs6783836 was not associated with HFS in patients treated with FOLFOX (OR = 0.86, 95% CI = 0.44-1.7, $P = .65$) and the difference between regimens was significant ($P_{\text{interaction}} = 1.0 \times 10^{-3}$). rs6783836 was not associated with patient outcome regardless of chemotherapy regime (XELOX ± cetuximab, response OR = 1.0, 95% CI = 0.78-1.4, $P = .82$ and OS HR = 0.95, 95% CI = 0.82-1.1, $P = .46$; FOLFOX ± cetuximab, response OR = 0.77, 95% CI = 0.54-1.1, $P = .15$ and OS HR = 1.0, 95% CI = 0.86-1.2, $P = .78$). rs6783836 maps to intron 4 of *ST6GAL1* in a region involved in transcriptional elongation (Figure 3) and was not an eQTL.

3.3 | Investigating the relationship between rs6783836 and HFS in an independent cohort

rs6783836 was borderline significant for HFS in patients treated with capecitabine from QUASAR2 (OR = 0.66, 95% CI = 0.42-1.03, $P = .05$) but with an opposite direction of effect to that found in COIN and COIN-B (Table S2).

3.4 | Investigating other variants, genes and pathways associated with toxicities

No other SNPs were associated with toxicities in COIN and COIN-B at genome-wide significant levels, but eight SNPs were suggestive of association ($P < 1.0 \times 10^{-6}$, Table S3). No genes were associated with toxicities after correction for multiple testing (data not shown). Four gene sets—peripheral neuropathy with response to food, neutropenia with dendritic spine development, diarrhoea with co-receptor activity and skin rash with blood vessel endothelial cell migration, were associated after correction for multiple testing (Table S4).

3.5 | Understanding the interrelationship between genetic variation in *ST6GAL1*, T2D and HFS

rs3887925 in intron 1 of *ST6GAL1* was the lead SNP associated with T2D (OR = 0.94, 95% CI = 0.92-0.96, $P = 1.2 \times 10^{-8}$, Figure 3), although rs6783836 was not associated with T2D (OR = 0.93, 95% CI = 0.85-1.0, $P = .07$) nor diabetic skin lesions (OR = 1.1, 95% CI = 0.89-1.3, $P = .44$). rs3887925 and rs6783836 were not in linkage disequilibrium (LD) ($D' = 0.26$, $R^2 = .01$). The rs6783836-T allele

was associated with lowered lymphocyte count (beta = -0.0052 , 95% CI = -0.0087 to -0.0018 , $P = 2.7 \times 10^{-3}$) and lowered HbA1c levels (beta = -0.0047 , 95% CI = -0.0080 to -0.0013 , $P = 5.9 \times 10^{-3}$, Figure 4) that withstood correction for multiple testing. rs6783836 was also associated with psoriasis (OR = 0.91, 95% CI = 0.85-0.98, $P = 7.5 \times 10^{-3}$).

4 | DISCUSSION

It has previously been suggested that HFS may be a biomarker of efficacy to chemotherapy.^{2,3} However, others have suggested that since HFS is a cumulative toxicity, there may be a bias for those living longer simply having more HFS due to having more treatment. Our data only considered HFS after 12 weeks of treatment and we found that patients with HFS had better response to chemotherapy at 12 weeks. We also observed an improvement in OS when analysed under a linear model of toxicity. Similarly, an exploratory analysis of two German trials noted an association between HFS and OS, and found no difference between patients with early and late HFS.³⁰ Together, these data suggest that HFS should be tolerated where possible and that an understanding of the underlying mechanism may help improve treatment efficacy.

With regards to the underlying mechanism, we identified rs6783836 in *ST6GAL1* as a genome-wide significant biomarker for HFS in patients treated with XELOX, with or without, cetuximab. Diabetics are at an increased risk of developing HFS and we confirmed an association for *ST6GAL1* with T2D, and also found that rs6783836 was associated with glycated haemoglobin levels, a marker routinely used in the diagnosis and monitoring of diabetes, supporting an inter-relationship. *ST6GAL1* has a known role in inflammation³¹ and we found an association between rs6783836 and lymphocyte count and psoriasis. Given that others have associated *ST6GAL1* with psoriasis in an Asian population,³² these data support a link between these syndromes and an underlying defect in the inflammatory pathway. Further studies are necessary to understand whether rs6783836 activates or inactivates *ST6GAL1*; however, it is interesting to note that *ST6GAL1* knockout mice are susceptible to ionising radiation and exhibit weight loss, gastrointestinal permeability and diarrhoea.³³

The odds ratios and betas for rs6783836 / *ST6GAL1* with T2D, lymphocyte count and psoriasis were in the opposite direction to HFS. Lin et al³⁴ proposed a flip-flop mechanism for allelic heterogeneity caused by interacting loci in weak LD and this has gained support from recent studies³⁵⁻³⁹ and may help explain our observations. Interestingly, the association with HFS was not found in patients treated with FOLFOX and was borderline significant, but with allele flipping, in patients from QUASAR2 treated with capecitabine alone. Further studies are therefore warranted to understand the underlying process and its specificity to particular therapeutic combinations.

We noted an association between peripheral neuropathy and genes involved in response to food which is supported by a previous observation linking diet to chemotherapy-induced peripheral neuropathy (CIMP).⁴⁰ Other forms of peripheral neuropathy have also been

linked with diet.^{41,42} Adopting vegetarianism has been shown to relieve symptoms in patients with diabetic neuropathy,^{43,44} and there is evidence that taking multivitamins reduces the likelihood of a patient experiencing CIMP.⁴⁵ Our data adds weight to this promising avenue for treatment of this toxicity.

AUTHOR CONTRIBUTIONS

The work reported in the article has been performed by the authors, unless specified in the text. Jeremy P. Cheadle obtained funding for our study. The study was designed and directed by Jeremy P. Cheadle. Timothy S. Maughan was CI of COIN and Ayman Madi was a COIN trial fellow—both provided clinical advice and supported the translational research. Richard Kaplan managed the COIN and COIN-B trials and facilitated access to the clinical data. Nada A. Al-Tassan oversaw the genotyping and Richard S. Houlston oversaw the imputation and quality control. Claire Palles, Rachel Kerr and David J. Kerr provided data from QUASAR2. Christopher Wills oversaw quality control of the UK Biobank data. Katie Watts undertook all of the statistical analyses with supervision from Valentina Escott-Price and Jeremy P. Cheadle. Katie Watts and Jeremy P. Cheadle interpreted the data with input from Christopher Wills and Valentina Escott-Price. Katie Watts wrote the first draft of the article with subsequent input from Jeremy P. Cheadle and all authors provided comments.

ACKNOWLEDGEMENTS

We thank Hywel Williams for helpful advice and the patients and their families who participated and gave their consent for this research, and the investigators throughout the UK who submitted samples for assessment. The Wales Gene Park provided compute storage.

CONFLICT OF INTEREST

Timothy S. Maughan consults for AstraZeneca and receives personal fees from Pierre Fabre (IDMC services). Timothy S. Maughan received research funding from Merck KgAa and AstraZeneca. The institution where Timothy S. Maughan works receives funding from Bayer. David J. Kerr is a director of Oxford Cancer Biomarkers. All other authors have declared no conflicts of interest.

DATA AVAILABILITY STATEMENT

The GWAS summary statistics are available through the NHGRI-EBI GWAS Catalogue under study accession numbers GCST90095054-GCST9009572. Further details and other data that supports the findings of our study are available from the corresponding author upon request. This research was also conducted using the UK Biobank Resource under application number 65833.

ETHICS STATEMENT

All patients gave fully informed consent for bowel cancer research (approved by REC [04/MRE06/60]).

ORCID

Jeremy P. Cheadle  <https://orcid.org/0000-0001-9453-8458>

REFERENCES

1. Milano G, Etienne-Grimaldi MC, Mari M, et al. Candidate mechanisms for capecitabine-related hand-foot syndrome. *Br J Clin Pharmacol*. 2008;66(1):88-95. doi:10.1111/J.1365-2125.2008.03159.X
2. Stintzing S, Fischer von Weikersthal L, Vehling-Kaiser U, et al. Correlation of capecitabine-induced skin toxicity with treatment efficacy in patients with metastatic colorectal cancer: results from the German AIO KRK-0104 trial. *Br J Cancer*. 2011;105(2):206-211. doi:10.1038/bjc.2011.227
3. Zielinski C, Lang I, Beslija S, et al. Predictive role of hand-foot syndrome in patients receiving first-line capecitabine plus bevacizumab for HER2-negative metastatic breast cancer. *Br J Cancer*. 2016;114(2):163-170. doi:10.1038/bjc.2015.419
4. Diasio R. Oral DPD-inhibitory fluoropyrimidine drugs - PubMed. *Oncology (Williston Park)*. 2000;14:19-23.
5. Kooner JS, Saleheen D, Sim X, et al. Genome-wide association study in individuals of south Asian ancestry identifies six new type 2 diabetes susceptibility loci. *Nat Genet*. 2011;43(10):984-989. doi:10.1038/ng.921
6. Mahajan A, Taliun D, Thurner M, et al. Fine-mapping type 2 diabetes loci to single-variant resolution using high-density imputation and islet-specific epigenome maps. *Nat Genet*. 2018;50(11):1505-1513. doi:10.1038/s41588-018-0241-6
7. Kaburagi T, Kizuka Y, Kitazume S, Taniguchi N. The inhibitory role of α 2,6-sialylation in adipogenesis. *J Biol Chem*. 2017;292(6):2278-2286. doi:10.1074/jbc.M116.747667
8. Yasukawa Z, Sato C, Kitajima K. Inflammation-dependent changes in α 2,3-, α 2,6-, and α 2,8-sialic acid glycotopes on serum glycoproteins in mice. *Glycobiology*. 2005;15(9):827-837. doi:10.1093/glycob/cwi068
9. Nasirikenari M, Segal BH, Ostberg JR, Urbasic A, Lau JTY. Altered granulopoietic profile and exaggerated acute neutrophilic inflammation in mice with targeted deficiency in the sialyltransferase ST6Gal I. *Blood*. 2006;108(10):3397-3405. doi:10.1182/blood-2006-04-014779
10. Nasirikenari M, Chandrasekaran EV, Matta KL, et al. Altered eosinophil profile in mice with ST6Gal-1 deficiency: an additional role for ST6Gal-1 generated by the P1 promoter in regulating allergic inflammation. *J Leukoc Biol*. 2010;87(3):457-466. doi:10.1189/jlb.1108704
11. Nasirikenari M, Lugade AA, Neelamegham S, et al. Recombinant sialyltransferase infusion mitigates infection-driven acute lung inflammation. *Front Immunol*. 2019;10:48. doi:10.3389/fimmu.2019.00048
12. Dorsett KA, Marciel MP, Hwang J, Ankenbauer KE, Bhalerao N, Bellis SL. Regulation of ST6GAL1 sialyltransferase expression in cancer cells. *Glycobiology*. 2021;31(5):530-539. doi:10.1093/GLYCOB/CWAA110
13. Lee M, Lee HJ, Bae S, Lee YS. Protein sialylation by sialyltransferase involves radiation resistance. *Mol Cancer Res*. 2008;6(8):1316-1325. doi:10.1158/1541-7786.MCR-07-2209
14. Schultz MJ, Swindall AF, Wright JW, Sztul ES, Landen CN, Bellis SL. ST6Gal-I sialyltransferase confers cisplatin resistance in ovarian tumor cells. *J Ovarian Res*. 2013;6(1):25. doi:10.1186/1757-2215-6-25/
15. Britain CM, Holdbrooks AT, Anderson JC, Willey CD, Bellis SL. Sialylation of EGFR by the ST6Gal-I sialyltransferase promotes EGFR activation and resistance to gefitinib-mediated cell death. *J Ovarian Res*. 2018;11(12):12. doi:10.1186/S13048-018-0385-0
16. Duarte HO, Rodrigues JG, Gomes C, et al. ST6Gal1 targets the ectodomain of ErbB2 in a site-specific manner and regulates gastric cancer cell sensitivity to trastuzumab. *Oncogene*. 2021;40(21):3719-3733. doi:10.1038/s41388-021-01801-w
17. Maughan TS, Adams RA, Smith CG, et al. Addition of cetuximab to oxaliplatin-based first-line combination chemotherapy for treatment of advanced colorectal cancer: results of the randomised phase 3 MRC COIN trial. *Lancet*. 2011;377(9783):2103-2114. doi:10.1016/S0140-6736(11)60613-2
18. Adams RA, Meade AM, Seymour MT, et al. Intermittent versus continuous oxaliplatin and fluoropyrimidine combination chemotherapy for first-line treatment of advanced colorectal cancer: results of the randomised phase 3 MRC COIN trial. *Lancet Oncol*. 2011;12(7):642-653. doi:10.1016/S1470-2045(11)70102-4
19. Wasan H, Meade AM, Adams R, et al. Intermittent chemotherapy plus either intermittent or continuous cetuximab for first-line treatment of patients with KRAS wild-type advanced colorectal cancer (COIN-B): a randomised phase 2 trial. *Lancet Oncol*. 2014;15(6):631-639. doi:10.1016/S1470-2045(14)70106-8
20. Watts K, Wills C, Madi A, et al. Genome-wide association studies of toxicity to oxaliplatin and fluoropyrimidine chemotherapy with or without cetuximab in 1800 patients with advanced colorectal cancer. *Int J Cancer*. 2021;149(9):1713-1722. doi:10.1002/ijc.33739
21. Al-Tassan NA, Whiffin N, Hosking FJ, et al. A new GWAS and meta-analysis with 1000Genomes imputation identifies novel risk variants for colorectal cancer. *Sci Rep*. 2015;5:10442. doi:10.1038/srep10442
22. Purcell S, Neale B, Todd-Brown K, et al. PLINK: a toolset for whole-genome association and population-based linkage analysis. *Am J Hum Genet*. 2007;81:559-575.
23. Turner SD. qqman: an R package for visualizing GWAS results using Q-Q and Manhattan plots. *J Open Source Softw*. 2018;3(25):731. doi:10.1101/005165
24. Little J, Higgins JPT, Ioannidis JPA, et al. Strengthening the Reporting of Genetic Association Studies (STREGA): an extension of the STROBE statement. *PLoS Med*. 2009;6(2):e22. doi:10.1371/journal.pmed.1000022
25. de Leeuw CA, Mooij JM, Heskes T, Posthuma D. MAGMA: generalized gene-set analysis of GWAS data. *PLOS Comput Biol*. 2015;11(4):e1004219. doi:10.1371/journal.pcbi.1004219
26. Moore CM, Jacobson SA, Fingerlin TE. Power and sample size calculations for genetic association studies in the presence of genetic model misspecification. *Hum Hered*. 2019;84(6):256-271. doi:10.1159/000508558
27. Kerr RS, Love S, Segelov E, et al. Adjuvant capecitabine plus bevacizumab versus capecitabine alone in patients with colorectal cancer (QUASAR 2): an open-label, randomised phase 3 trial. *Lancet Oncol*. 2016;17(11):1543-1557. doi:10.1016/S1470-2045(16)30172-3
28. Millard LAC, Davies NM, Gaunt TR, Smith GD, Tilling K. Software application profile: PHEASANT: a tool for performing automated phenotype scans in UK biobank. *Int J Epidemiol*. 2018;47(1):29-35. doi:10.1093/ije/dyx204
29. Kichaev G, Yang W-Y, Lindstrom S, et al. Integrating functional data to prioritize causal variants in statistical fine-mapping studies. *PLoS Genet*. 2014;10(10):e1004722. doi:10.1371/journal.pgen.1004722
30. Hofheinz RD, Heinemann V, Von Weikersthal LF, et al. Capecitabine-associated hand-foot-skin reaction is an independent clinical predictor of improved survival in patients with colorectal cancer. *Br J Cancer*. 2012;107(10):1678-1683. doi:10.1038/BJC.2012.434
31. Holdbrooks AT, Ankenbauer KE, Hwang J, Bellis SL. Regulation of inflammatory signaling by the ST6Gal-I sialyltransferase. *PLoS One*. 2020;15:e0241850. doi:10.1371/journal.pone.0241850
32. Wang H, Wang Z, Rani PL, et al. Identification of PTPN22, ST6GAL1 and JAZF1 as psoriasis risk genes demonstrates shared pathogenesis between psoriasis and diabetes. *Exp Dermatol*. 2017;26(11):1112-1117. doi:10.1111/exd.13393
33. Punch PR, Irons EE, Manhardt CT, Marathe H, Lau JTY. The sialyltransferase ST6GAL1 protects against radiation-induced gastrointestinal damage. *Glycobiology*. 2020;30(7):446-453. doi:10.1093/GLYCOB/CWZ108
34. Lin PI, Vance JM, Pericak-Vance MA, Martin ER. No gene is an Island: the flip-flop phenomenon. *Am J Hum Genet*. 2007;80(3):531-538. doi:10.1086/512133

35. Wang S, Qian F, Zheng Y, et al. Genetic variants demonstrating flip-flop phenomenon and breast cancer risk prediction among women of African ancestry. *Breast Cancer Res Treat*. 2018;168(3):703-712. doi:[10.1007/s10549-017-4638-1](https://doi.org/10.1007/s10549-017-4638-1)
36. Maher BS, Reimers MA, Riley BP, Kendler KS. Allelic heterogeneity in genetic association meta-analysis: an application to DTNBP1 and schizophrenia. *Hum Hered*. 2010;69(2):71-79. doi:[10.1159/000264445](https://doi.org/10.1159/000264445)
37. Zaykin DV, Shibata K. Genetic flip-flop without an accompanying change in linkage disequilibrium. *Am J Hum Genet*. 2008;82(3):794-796. doi:[10.1016/j.ajhg.2008.02.001](https://doi.org/10.1016/j.ajhg.2008.02.001)
38. Shao S, Niu Y, Zhang X, et al. Opposite associations between individual KIAA0319 polymorphisms and developmental dyslexia risk across populations: a stratified meta-analysis by the study population. *Sci Rep*. 2016;6(1):1-9. doi:[10.1038/srep30454](https://doi.org/10.1038/srep30454)
39. Mersha TB, Martin LJ, Biagini Myers JM, et al. Genomic architecture of asthma differs by sex. *Genomics*. 2015;106(1):15-22. doi:[10.1016/j.ygeno.2015.03.003](https://doi.org/10.1016/j.ygeno.2015.03.003)
40. Mongiovi JM, Zirpoli GR, Cannioto R, et al. Associations between self-reported diet during treatment and chemotherapy-induced peripheral neuropathy in a cooperative group trial (S0221). *Breast Cancer Res*. 2018;20(1):146. doi:[10.1186/s13058-018-1077-9](https://doi.org/10.1186/s13058-018-1077-9)
41. Spagnoli C, Pisani F, Di Mario F, et al. Peripheral neuropathy and gastroenterologic disorders: an overview on an underrecognized association. *Acta Biomed*. 2018;89(9-S):22-32. doi:[10.23750/abm.v89i9-S.7956](https://doi.org/10.23750/abm.v89i9-S.7956)
42. Chopra K, Tiwari V. Alcoholic neuropathy: possible mechanisms and future treatment possibilities. *Br J Clin Pharmacol*. 2012;73(3):348-362. doi:[10.1111/j.1365-2125.2011.04111.x](https://doi.org/10.1111/j.1365-2125.2011.04111.x)
43. Crane MG, Sample C. Regression of diabetic neuropathy with total vegetarian (vegan) diet. *J Nutr Med*. 1994;4(4):431-439. doi:[10.3109/13590849409003592](https://doi.org/10.3109/13590849409003592)
44. Bunner AE, Wells CL, Gonzales J, Agarwal U, Bayat E, Barnard ND. A dietary intervention for chronic diabetic neuropathy pain: a randomized controlled pilot study. *Nutr Diabetes*. 2015;5(5):e158. doi:[10.1038/nutd.2015.8](https://doi.org/10.1038/nutd.2015.8)
45. Zirpoli GR, McCann SE, Sucheston-Campbell LE, et al. Supplement use and chemotherapy-induced peripheral neuropathy in a cooperative group trial (S0221): the DELCaP study. *J Natl Cancer Inst*. 2017;109(12):dix098. doi:[10.1093/jnci/djx098](https://doi.org/10.1093/jnci/djx098)

SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.

How to cite this article: Watts K, Wills C, Madi A, et al. Genetic variation in *ST6GAL1* is a determinant of capecitabine and oxaliplatin induced hand-foot syndrome. *Int J Cancer*. 2022;151(6):957-966. doi:[10.1002/ijc.34046](https://doi.org/10.1002/ijc.34046)