

1 **Capture Hi-C identifies putative target genes at 33 breast cancer risk loci.**

2 Joseph S Baxter^{1*}, Olivia C Leavy^{2,3*}, Nicola H Dryden^{1*}, Sarah Maguire¹, Nichola Johnson¹, Vita
3 Fedele¹, Nikiana Simigdala¹, Lesley-Ann Martin¹, Simon Andrews⁴, Steven W Wingett⁴, Ioannis
4 Assiotis⁵, Kerry Fenwick⁵, Ritika Chauhan¹, Alistair G Rust¹, Nick Orr¹, Frank Dudbridge^{2,3}, Syed
5 Haider^{1#}, Olivia Fletcher^{1#}.

6 ¹Breast Cancer Now Toby Robins Research Centre, The Institute of Cancer Research, London, SW3
7 6JB, UK

8 ²Department of Non-communicable Disease Epidemiology, The London School of Hygiene and
9 Tropical Medicine, London, WC1E 7HT, UK

10 ³Department of Health Sciences, University of Leicester, Leicester, LE1 7RH, UK

11 ⁴Bioinformatics Group, The Babraham Institute, Cambridge CB22 3AT, UK

12 ⁵Tumour Profiling Unit, The Institute of Cancer Research, London, SW3 6JB, UK

13 * Contributed equally #Joint corresponding authors

14

15 Olivia Fletcher	Syed Haider
16 The Institute of Cancer Research,	The Institute of Cancer Research,
17 London, SW3 6JB, UK	London, SW3 6JB, UK
18 Tel: +44 (0) 207 153 5177	Tel: +44 (0) 207 153 5338
19 Fax: +44 (0) 207 153 5340	Fax: +44 (0) 207 153 5340
20 E-mail: Olivia.Fletcher@icr.ac.uk	E-mail: Syed.Haider@icr.ac.uk

21

22 **Key words:** Breast cancer; risk locus; target gene.

23

24

25 **ABSTRACT**

26 Genome-wide association studies (GWAS) have identified approximately 100 breast cancer risk loci.
27 Translating these findings into a greater understanding of the mechanisms that influence disease risk
28 requires identification of the genes or non-coding RNAs that mediate these associations. We used
29 Capture Hi-C (CHi-C) to annotate 63 loci; we identified 110 putative target genes at 33 loci. All CHi-C
30 interaction peaks can be viewed at bit.ly/CHiC-BC. To assess the support for these target genes in
31 other data sources we tested for associations between levels of expression and SNP genotype
32 (eQTLs), disease-specific survival (DSS), and compared them with somatically mutated cancer genes.
33 22 putative target genes were eQTLs, 32 were associated with DSS and 14 have been shown to be
34 somatically mutated in breast, or other, cancers. Identifying the target genes at GWAS risk loci will
35 lead to a greater understanding of the mechanisms that influence breast cancer risk and prognosis.

36

37

38 Genome-wide association studies (GWAS) coupled with large-scale replication and fine mapping
39 studies have led to the identification of approximately 100 breast cancer risk loci. Breast cancer is a
40 heterogeneous disease with two main subtypes defined by the presence (ER+) or absence (ER-) of
41 the estrogen receptor. Approximately 80% of newly diagnosed breast cancers are ER+ although this
42 proportion varies with age at diagnosis and ethnicity¹. The majority of breast cancer GWAS risk loci
43 have been identified on the basis of their association with overall breast cancer risk, or risk of ER+
44 disease². Most of the risk single nucleotide polymorphisms (SNPs) map to non-protein-coding
45 regions and are thought to influence transcriptional regulation^{3,4}; many map to gene deserts with
46 the nearest known protein-coding genes mapping several hundred kilobases (kb) away. Translating
47 these findings into a greater understanding of the mechanisms that influence an individual woman's
48 risk requires the identification of causal variants and the targets of these causal variants (i.e. genes
49 or non-coding RNAs that mediate the associations observed in GWAS). Systematic approaches to the
50 functional characterisation of cancer risk loci have been proposed^{4,5}. These include fine mapping of
51 potentially large genomic regions (defined as regions that include all SNPs correlated with the
52 published SNP with an $r^2 \leq 0.2$), the analysis of SNP genotypes in relation to expression of nearby
53 genes (eQTL) and the use of chromatin association methods (chromosome conformation capture
54 (3C) and Chromatin Interaction Analysis by Paired-End Tag Sequencing (ChIA-PET)) of regulatory
55 regions to determine the identities of target genes. To facilitate a high-throughput approach to the
56 identification of target genes at GWAS risk loci we developed Capture Hi-C (ChI-C)⁶. This novel Hi-C
57 protocol⁷ allows high-throughput, high resolution analysis of physical interactions between
58 regulatory elements and their target genes. We have used ChI-C previously, to characterise three
59 breast cancer risk loci mapping to gene deserts at 2q35, 8q24.21 and 9q31.2⁶. Here we selected 63
60 established breast cancer risk loci (Supplementary Table 1); we identify ChI-C interaction peaks
61 involving 110 putative target genes mapping to 33 loci and demonstrate long-range interaction
62 peaks some of which span megabase (Mb) distances and involve adjacent risk loci. We carry out

63 eQTL analyses, analyses of disease specific survival (DSS) and compare our putative target genes
64 with somatically mutated cancer genes to assess the orthogonal support for these putative target
65 genes.

66 **RESULTS**

67 Generating CHI-C libraries representing 63 breast cancer GWAS risk loci

68 We generated CHI-C libraries in two ER+ breast cancer cell lines (T-47D, ZR-75-1), two ER- breast
69 cancer cell lines (BT-20, MDA-MB-231), one “normal” breast epithelial cell line (Bre80-Q-TERT
70 (Bre80)) and a control, non-breast lymphoblastoid cell line (GM06990) (Supplementary Figure 1). We
71 defined an interaction peak as any pair of HindIII fragments for which the number of di-tags was
72 significantly (FDR adjusted $P < 0.01$) greater than expected under a negative binomial model, taking
73 into account both the distance between the HindIII fragments and the propensity of the bait
74 fragment to form interactions (“interactability”; Methods). The number of di-tags that constituted
75 an interaction peak depended on the distance between the interacting fragments and ranged from 5
76 to 14,151. We defined a locus as a single continuous capture region, annotated by at least one risk
77 SNP (Methods).

78 Distribution of CHI-C interaction peaks across the 63 risk loci

79 The number of interaction peaks at each locus, in each cell line ranged from zero to 1,744
80 (Supplementary Table 2, Supplementary Figure 2), with two outliers (1,744 at 8q21.11-rs2943559 in
81 ZR-75-1 and 1,007 at 8q24.21-rs13281615 in T-47D). There were 12 loci (19.0%) at which there were
82 no interaction peaks in any of the cell lines we examined (Supplementary Table 2); these loci were
83 excluded from further analyses. 46 (90.2%) of the 51 loci that we were able to analyse were
84 identified on the basis of their association with overall breast cancer risk or risk of ER+ disease; the
85 exceptions were 2p24.1-rs12710696, 5p15.33-rs10069690, 6q25.1-rs12662670, rs2046210, 16q12.2-

86 rs11075995 and 19p13.11-rs8170 which were associated with ER- and/or triple negative breast
87 cancer (TNBC)⁸⁻²⁰.

88 We first tested for differences in the median number of interaction peaks across the six different cell
89 lines according to ER status and cell type (breast/non-breast). The median number of interaction
90 peaks per locus varied significantly between cell lines (Kruskal-Wallis test $P = 0.0006$; Table 1, Figure
91 1a). There were, on average, more statistically significant interaction peaks per locus in the ER+
92 breast cancer cell lines (T-47D, ZR-75-1) compared to the ER- breast cancer cell lines (BT-20, MDA-
93 MB-231, Mann-Whitney test $P = 0.0008$) or the control lymphoblastoid cell line (GM06990, $P =$
94 0.002). There was, however, no difference between the number of interaction peaks per locus in the
95 ER+ breast cancer cell lines and the normal mammary epithelial cell line (Bre80, $P = 0.85$; Table 1),
96 consistent with Bre80 representing a progenitor cell population that gives rise to ER+ breast cancer
97 cells. Similarly, the median distance between interacting fragments varied across cell lines (Kruskal-
98 Wallis test $P < 1 \times 10^{-6}$; Table 1, Figure 1b) with a greater proportion of longest range interaction
99 peaks (>2Mb) in the ER+ breast cancer cell lines compared to all other cell lines. In an analysis of 51
100 breast cancer cell lines, there was no evidence that luminal (ER+) cell lines carried more genome
101 aberrations than basal (ER-) cell lines.²¹ We cannot, however, exclude the possibility that
102 rearrangements, gains and losses occur preferentially at ER+ risk loci in ER+ cell lines and this
103 contributes to the higher proportion of long range interaction peaks we observe in the T-47D and
104 ZR-75-1 breast cancer cell lines.

105 To gain further insight into the relationship between interaction peaks and cell type-specificity, we
106 looked at the number of interaction peaks that were identical between two or more cell types. Of
107 the 12,736 interaction peaks we identified 7,681 (60.3%) were present in a single cell line and 5,055
108 (39.7%) were present in multiple cell lines (FDR adjusted $P < 0.01$ Supplementary Table 3). The
109 subset of interaction peaks that were present in at least two cell lines are provided as
110 supplementary data (Supplementary Data). Excluding two loci with large outliers (i.e. 8q21.11-

111 rs2943559 and 8q24.21-rs13281615 at which there were 1,744 and 1,007 interaction peaks in T-47D
112 and ZR-75-1, respectively; Supplementary Figure 2) the numbers were 4,924 (50.6%) and 4,805
113 (49.4%; Supplementary Table 3). We found a statistically significant excess of interaction peaks that
114 were common to all four breast cancer cell lines ($N = 62$, $P = 0.0003$, Figure 1c) and all five breast cell
115 lines ($N = 53$, $P < 0.0001$ Figure 1d). We also found an excess of interaction peaks that were exclusive
116 to the lymphoblastoid cell line ($N = 304$, $P < 0.0001$) suggesting that at least a subset of interaction
117 peaks show cell type specificity. Comparing the cell lines according to receptor type, the interaction
118 peaks were marginally more similar within the two ER+ cell lines (Jaccard similarity coefficient =
119 0.18) and the two ER- cell lines (Jaccard similarity coefficient = 0.13) than between them (Figure 1e).
120 Representative examples of loci that demonstrated cell type-specific activity are shown in Figure 2.
121 At several of the ER+ risk loci, including the 10q26.13-rs2981579 (*FGFR2*) and 14q13.3-rs2236007
122 (*PAX9*) risk loci, we observed interaction peaks that were restricted to the two ER+ breast cancer cell
123 lines and the normal breast epithelial cell line (Figures 2a and c). In both these examples, the
124 transcription start site (TSS) of the target gene maps within the capture region and forms interaction
125 peaks with specific HindIII fragments that map several hundred kb from the capture region. These
126 distal fragments colocalise with DNase I hypersensitive sites, CTCF, FOXA1, GATA3 and/or ER α
127 binding sites in T-47D cells and in both of these examples the orientation of the CTCF binding sites is
128 towards the captured locus (Figures 2b, d and e)²².
129 There were, however, many exceptions to the pattern of ER+ risk loci forming interaction peaks in
130 the ER+ breast cancer cell lines and Bre80s. The 11p15.5-rs3817198 risk locus, which is associated
131 with ER+ breast cancer forms multiple interaction peaks in the ER- breast cancer cell lines, but not in
132 the ER+ breast cancer cell lines or in Bre80 (Figure 3a, Supplementary Table 2) and the 6q25.1-
133 rs2046210 (*ESR1*) locus which has been shown to be preferentially associated with ER- breast
134 cancer^{20,23} forms interaction peaks in the ER+ but not the ER- breast cancer cell lines (Figure 3b,
135 Supplementary Table 2).

136 Defining putative target genes

137 We defined putative target genes as genes that mapped within, or *in cis* ($\leq 5\text{Mb}$) to, a captured
138 region and for which the TSS mapped to an interacting fragment in at least two cell lines (Methods).
139 On this basis, we were able to assign 110 putative target genes to 33 (64.7%) of the 51 loci (Table 2,
140 Supplementary Table 4); 94 were protein coding and 16 were non-coding RNAs. The number of
141 genes per locus, for these 33 loci ranged from one (13 loci) to 19 (11q13.1-rs3903072 locus) with a
142 median of two. The distance between the published risk SNP and the TSS of the Chi-C target gene
143 ranged from 1 kb (*KCNN4*) to more than 4 Mb (3p26.1-rs6762644 with *CAV3*, *RAD18* and *SETD5*;
144 11q13.1-rs3903072 with *FADD*) with a median of 135 kb (individual distances between risk SNPs and
145 Chi-C target genes are given in Supplementary Table 5). Amongst our 51 informative risk loci there
146 were 24 (at 12 chromosomal regions) that mapped within five Mb of another locus (Supplementary
147 Table 1). We observed interaction peaks between adjacent loci at eight of these chromosomal
148 regions and were able to potentially assign target genes to three additional loci on the basis of
149 interaction peaks with the adjacent locus (Table 2). These loci were 8q21.11-rs6472903 (*HNF4G* and
150 *PEX2*), 9q31.2-rs10759243 (*KLF4*) and 14q24.1-rs999737 (*ZFP36L1*); a representative example,
151 showing interaction peaks within and between adjacent loci at 11q13, is shown in Figure 4. For the
152 interaction peaks with the longest-range gene targets (3p26.1-rs6762644: *CAV3*, *LINC00312*, *LMCD1*,
153 *c3orf32*, *RAD18*, *SETD5*, 4q24-rs9790517: *CENPE*, 8q24.21-rs13281615: *CCDC26*, 11q13.1-rs3903072:
154 *CCND1*, *FADD*) we aligned our data with topologically associated domains (TADs) generated in
155 Human Mammary Epithelial Cells (HMEC)²⁴. At each of these loci we observed interaction peaks
156 between captured fragments and target gene(s) mapping within the same TAD but also, less
157 frequently, with target gene(s) mapping to a different TAD (Supplementary Figure 3).

158 To determine whether the target genes selected using a Chi-C approach differ from those selected
159 simply on the basis of proximity to the GWAS risk SNP (“nearest gene approach”) we compared the
160 two approaches directly. At 15 of the 51 loci included in our analysis, we were unable to assign

161 target genes (i.e. there were no TSS directly or indirectly involved in statistically significant
162 interaction peaks). Of the 36 loci at which we were able to assign at least one target gene directly
163 (N=33) or indirectly (N=3) there were 24 at which the nearest gene was either the only CHI-C target
164 gene (N=9; Table 2) or one of several CHI-C target genes (N=15; Table 2). There were, however, 12
165 loci at which our data implicated genes other than the nearest gene; these loci included 13q13.1-
166 rs11571833 (CHI-C gene: *PDSB5*, nearest gene: *BRCA2*), 14q24.1-rs2588809 and rs999737 (CHI-C
167 gene: *ZFP36L1*, nearest gene: *RAD51B*) and 16q12.2-rs17817449 and rs11075995 (CHI-C genes:
168 *CRNDE*, *IRX5*, *IRX3*, *LOC100996*, nearest gene: *FTO*).

169 CHI-C target genes and eQTL analyses

170 To assess the likelihood of our putative target genes having a causal role in breast cancer aetiology,
171 we first carried out eQTL analyses using the published risk SNPs (or a close proxy, $r^2 > 0.8$) and RNA-
172 Seq data from the Cancer Genome Atlas²⁵ (TCGA) adjusted for matched DNA methylation and
173 somatic copy-number alterations. Many of risk loci we included have been shown to be associated
174 with breast cancer risk overall, albeit with evidence that the association may differ in magnitude
175 between ER+ and ER- cancers for some²⁶. Accordingly, we carried out eQTL analyses for all cancers
176 combined (N = 547) and then stratified by ER status (ER+ N = 415, ER- N = 95, ER unknown N = 37).
177 There were 9 loci (26 protein coding genes) at which there were no suitable proxies, and levels of
178 expression of 18 of our putative target genes (*KRTPA5-5*, *KRTAP5-6* and 16 non-coding RNAs) were
179 too low for analysis. eQTL analysis of the remaining 26 loci (69 protein coding genes) identified 22
180 SNP-gene combinations that were nominally significant ($P < 0.05$) in all, ER+ or ER- breast cancers
181 (Supplementary Table 6), nine of which remained significant after taking account of multiple testing
182 (FDR adjusted $P < 0.1$, Table 3). Comparing these eQTLs with “nearest genes”, three were nearest
183 genes and six were not. Including all nearest genes (regardless of whether they were also a CHI-C
184 target gene) in our eQTL analysis we found two additional SNP-gene combinations that were not
185 captured by our CHI-C analysis; rs4808801 was associated with levels of expression of *ELL* in all

186 cancers and ER+ cancers and rs8170 was associated with levels of expression of *ANKLE* in all cancers
187 (FDR adjusted $P < 0.1$).

188 Several of the CHi-C target gene eQTLs were consistent with previous reports including *IGFBP5* at
189 2q35-rs13387042^{6,27}, *COX11* at 17q22-rs6504950²⁸ and *LRRC25* at 19p13.11-rs4808801²⁹. Novel
190 eQTLs included genes that mapped within the capture region, proximal to the reported risk SNP such
191 as *CDCA7* at 2q31.1-rs1550623, *SSBP4* at 19p13.11-rs4808801 and *MRPL34* at 19p13.11-rs8170 as
192 well as genes that mapped several hundred kb from the reported SNP including *IRX3* at 16q12.2-
193 rs17817449 and *ZFP36L1* at 14q24.1-rs2588809. At 11q13.1-rs3903072, eQTL analyses support
194 multiple putative target genes of which *SNX32*, *CTSW* and *CFL1* map within the capture region but,
195 intriguingly, *FADD* and *CCND1* map at a distance of approximately 4Mb from rs39030702 (Figure 4).
196 Both *FADD* and *CCND1* map to a region of chromosome 11 that is frequently subject to
197 amplifications and copy number gains in breast cancer (*FADD* and *CCND1* map to regions of copy-
198 number gain in 20 – 30% of Metabric³⁰ and TCGA samples), raising the concern that this long range
199 eQTL association might be influenced by these samples. Excluding samples with genomic copy-
200 number gains from the analysis, however, strengthened the association between 11q13.1-rs3903072
201 and *FADD* (all samples $P_{ER+} = 0.01$, excluding 119 samples with copy-number gains $P_{ER+} = 0.004$;
202 Figure 5a and 5c), but not *CCND1* (all samples $P_{ER+} = 0.04$, excluding 130 samples $P_{ER+} = 0.05$:
203 Supplementary Figure 4).

204 CHi-C target genes and disease-specific survival (DSS)

205 To our knowledge, only one of the risk SNPs we included has been associated with disease prognosis
206 (16q12.1-rs3803662 and *TOX3*³¹); this may reflect a fundamental difference between the genetics of
207 predisposition and prognosis or a relative lack of power for observational studies of outcome in
208 which detailed information on treatment is generally lacking. As any individual regulatory variant
209 may only explain a small proportion of the total variance in gene expression, however, we looked
210 directly for an association between levels of expression of our putative target genes and patient

211 outcome in the Metabric breast cancer cohort³⁰. Given the profound effect of ER status on outcome,
212 we performed survival analyses on ER+ and ER- subpopulations separately. Of the 97 putative target
213 genes (94 protein coding, three non-coding RNAs) for which levels of expression were available, 32
214 (33%) were associated with disease-specific survival (DSS) in individuals with ER+ disease; none was
215 associated with DSS in ER- disease (FDR adjusted $P < 0.1$; Supplementary Table 7). Comparing these
216 32 genes with those for which we found eQTL associations in ER+ cancers (nominal $P < 0.05$) there
217 were six that were common to both groups (*CFL1*, *FADD*, *MRPL34*, *IGFBP5*, *IRX3*, *ZFP36L1*). In
218 addition, there was a highly significant association between levels of expression of *CDCA7* and DSS (P
219 $= 1.22 \times 10^{-8}$), which maps just 7 kb from the reported 2q31.1 risk SNP (rs1550623) but while there
220 was a robust eQTL association between rs1550623 and *CDCA7* in all cancers (nominal $P = 0.007$, FDR
221 adjusted $P = 0.09$) there was no association in ER+ cancers alone (both $P > 0.1$). We also observed
222 highly significant associations (FDR adjusted $P < 0.005$) for five genes that were excluded from eQTL
223 analysis due to a lack of suitable tag SNP (*CENPE* at 4q24, *TPCN2* and *ORAOV1* at 11q13.3, *PDS5B* at
224 13q13.1 and *SLC4A7* at 3p24.1: Supplementary Figure 5)

225 CHI-C target genes and somatic mutations in cancer genes

226 Finally, we compared our CHI-C putative target genes with the list of 727 cancer genes compiled by
227 Nik-Zainal and colleagues in their analysis of whole genome sequences of 560 breast cancers³². The
228 94 protein-coding CHI-C target genes are highly enriched for these cancer genes (14 observed,
229 Hypergeometric $P = 2.02 \times 10^{-6}$) and include well-documented cancer genes (*CCND1*, *CDKN2A*,
230 *CDKN2B*, *MYC*, *MAP3K1*, *ESR1* and *FGFR2*) as well as relatively uncharacterised examples (*TET2*,
231 *KLF4*, *MLLT10*, *FADD*, *TBX3*, *PAX9* and *ZFP36L1*).

232 Combining the somatic mutation data with the eQTL and DSS analyses, there were 48 CHI-C target
233 genes mapping to 32 loci for which there was orthogonal support from at least one additional source
234 and six genes mapping to six loci for which there was support from at least two additional sources
235 (Table 4). For four of these, *CDCA7*, *FADD*, *ZFP36L1* and *MRPL34*, levels of expression were

236 associated with both SNP genotype and DSS (Table 4) and we were able to assess whether high (or
237 low) levels of expression were similarly associated with risk and poor outcome. For *FADD* the
238 associations are inconsistent; the rare allele of 11q13.1-rs3903072 is associated with higher levels of
239 expression (Figure 5a) and lower risk²⁶, but higher levels of expression are associated with poor
240 outcome (Figure 5b). In addition, the strong influence of copy number gains on levels of expression
241 of *FADD* confounds both eQTL and DSS analyses with opposite effects; excluding 119 ER+ cancers
242 with copy-number gains strengthens the eQTL association in TCGA (Figure 5c), excluding 345 such
243 samples from the analysis of outcome in Metabric abrogates the association with DSS (Figure 5d)
244 suggesting that samples with copy-number gains at this region may have a poor outcome that is not
245 directly related to levels of expression of *FADD*. Similarly, for *CDCA7* the associations are
246 inconsistent. The risk allele of rs1550623 is the common allele²⁶; the common allele is associated
247 with lower levels of *CDCA7* expression (Figure 5e) but lower levels of expression of *CDCA7* are
248 associated with a better prognosis (Figure 5f). However, for both 14q24.1-rs2588809 (*ZFP36L1*) and
249 19p13.1-rs8170 (*MRPL34*) the rare alleles are associated with lower levels of expression (Figure 5g
250 and 5i) and higher risk²⁶; lower levels of expression are also associated with a poor outcome (Figure
251 5h and 5j) consistent with these genes acting as tumour suppressors influencing both predisposition
252 and outcome similarly.

253 **DISCUSSION**

254 The purpose of this analysis was to identify target genes at 63 breast cancer GWAS risk loci using an
255 unbiased, high resolution chromosome association method, CHi-C and evaluate this method in
256 comparison to a simple “nearest gene” approach. We were able to assign 110 putative target genes
257 to 33 loci; 94 were protein coding and 16 were non-coding RNAs. We used three publicly available
258 data sources to assess support for our CHi-C target genes as having a causal role in breast cancer
259 aetiology. In eQTL analyses we identified 22 SNP-gene combinations that were nominally significant
260 ($P < 0.05$) in all, ER+ or ER- breast cancers in TCGA. DSS analyses of ER+ breast cancers in the

261 Metabric cohort supported 32 Chi-C target genes (FDR adjusted $P < 0.1$) and 14 were listed in 727
262 cancer genes compiled by Nik-Zainal and colleagues. In all data sources combined there was support
263 for 48 Chi-C target genes mapping to 32 loci from at least one additional source and there was
264 support for six genes mapping to six loci from at least two additional sources. These data suggest
265 that a substantial proportion of the Chi-C putative target genes are likely to influence breast cancer
266 risk and warrant further investigation.

267 However, amongst the 63 risk loci that we investigated there were 12 at which we detected no
268 interaction peaks at all. This may, in part, be a consequence of our methodology as 3C-based
269 techniques are not considered reliable for detecting interactions over distances of less than 10 kb³³;
270 at three of these 12 loci (4q34.1-rs6828523 (*ADAM29*), 10q21.2-rs10995190 (*ZNF365*) and 16q12.1-
271 rs3803662 (*TOX3*) the TSS of a nominated target gene mapped within 10kb of the reported SNP.
272 Similarly, for the 15 loci at which the interaction peaks we detected did not include direct, or
273 indirect, interactions with the promoter of a RefSeq gene, there were four at which the TSS of the
274 proposed target gene mapped within 10 kb (6p25.3-rs11242675 (*FOXQ1*), 22q12.1-rs17879961
275 (*CHEK2*)) or 20kb (5p15.33-rs10069690 (*TERT*), 22q12.1-rs132390 (*EMID1*)) of the reported SNP
276 (Supplementary Table 2). In any analysis, there is a trade-off between type I and type II errors. By
277 using a rigorous threshold (FDR adjusted $P < 0.01$) for calling an interaction peak “significant” we will
278 have minimised false positives but we may also have missed potentially important low frequency
279 interactions. Finally, we may have missed important target genes by using a restricted set of cell
280 lines that will only capture interaction peaks between regulatory elements and genes that are
281 expressed in breast epithelial cells. At the other extreme, there were several loci mapping to gene-
282 rich regions (particularly 11q13 and 19p13), at which we observed interaction peaks with multiple
283 putative target genes some of which mapped to the same HindIII restriction fragment as another
284 target gene (Table 2). Reducing the size of the average restriction fragment, by using an enzyme that
285 cuts more frequently would provide greater resolution, but it is clear that Chi-C cannot resolve

286 interaction peaks at the TSS of putative target genes that map within a few hundred base pairs of
287 each other.

288 The other metrics that are frequently used for defining putative target genes are nearest gene, or
289 nearest plausible gene, and eQTL analyses. While in many cases our analyses support the nearest
290 gene or the nearest plausible gene the limitations to this approach are obvious; there are many
291 examples of long range interactions between regulatory elements and target genes that bypass
292 more proximal putative target genes³⁴⁻³⁶. Comparing CHi-C with a nearest-gene metric for assigning
293 putative target genes to risk loci, our data were informative at 36 (57%) of the 63 loci we selected
294 for analysis and at 27 (43%) our data implicated genes in addition to, or other than, the nearest
295 gene. Notably, our data implicates several protein coding genes and non-coding RNAs that map at
296 distances of more than 1Mb from the published risk SNP (ie outside the limit of many eQTL
297 analyses). While the presence of these long range interactions may inform future follow up studies,
298 they do not exclude effects that are more local to the risk loci. In their functional annotation of the
299 human genome, the ENCODE consortium estimated that the average number of TSSs that interact
300 with any given distal element is 2.5³⁷ and regulatory variants that map to such elements may
301 influence absolute or relative levels of expression of multiple genes. Of the six genes for which there
302 was support from at least two additional data sources, neither *ZFP36L1* (which maps 600 kb from
303 rs2588809) nor *FADD* (mapping 4.5 Mb from rs3903072) would have been selected by a nearest
304 gene metric supporting the use of CHi-C as a means of identifying putative target genes that map
305 several hundred kb or even Mb from the risk locus.

306 eQTL analyses provide an intuitive approach to the process of identifying putative target genes.
307 However, implicit in eQTL analyses of breast cancer or normal breast tissue is an assumption of a
308 model in which a breast cancer GWAS locus influences risk by altering steady-state expression of a
309 gene that is transcribed in normal or malignant breast tissue; this may not be true for a substantial
310 minority of loci. For this reason, a CHi-C based approach which detects “permissive” interaction

311 peaks (as well as “instructive” interaction peaks)^{33,38} may have benefits over an eQTL based
312 approach by allowing the identification of putative target genes that are poised for expression at a
313 particular stage of differentiation or in response to external stimuli such as hormones or DNA
314 damage.

315 The variants detected by GWAS are common variants with small effects (ORs are typically < 1.2) and
316 any individual risk SNP will usually only explain a small proportion of variance in levels of expression
317 of a target gene. For example, the association between 11q13.1-rs3903072 and *FADD* is weak in all
318 ER+ cancers (nominal P = 0.01); excluding ER+ cancers with copy number gains reduces the variance
319 in levels of expression of *FADD* and increases the proportion of variance explained by rs3903072
320 (nominal P = 0.004). Given the small effects of individual variants, eQTL approaches based on current
321 data sets of a few hundred samples lack power. To limit the penalty for multiple testing, most eQTL
322 analyses are restricted to genes within a 1Mb window of the risk SNP or a proposed causal variant.
323 In our eQTL analysis we used our ChI-C results to restrict our gene set to 69 protein-coding genes.
324 Despite this our eQTL analysis probably lacked power, particularly for the stratified analyses where
325 there were just four ER+ eQTLs and no ER- eQTLs that were significant after taking account of
326 multiple testing. Indeed *IGFBP5*, *KLF4*, *CFL1*, *CCND1* and *IRX3* are all fairly compelling putative target
327 genes with nominal associations for which the adjusted eQTL P-values were non-significant (all
328 nominal $P_{ER+} < 0.05$, all FDR adjusted $P_{ER+} > 0.1$).

329 For several of the risk loci that we included, functional annotation studies have been previously
330 reported on a locus-by-locus basis^{27,28,39-51} and target genes have been inferred by a combination of
331 proximity, eQTL analysis and testing for looping interactions on a candidate basis using 3C. Our ChI-C
332 targets are consistent with many of these^{27,28,43,46,47} but may implicate *CENPE* in addition to *TET2* at
333 4q24⁴⁰ and *MRPL34* in addition to *ABHD8* or *ANKLE1* at 19p13.1⁵¹. The other notable feature of our
334 data is the frequency with which we observed interaction peaks between adjacent loci several of
335 which map megabase distances apart. This feature, and our observation of an eQTL between

336 rs3903072 and both *CFL1* and *FADD* which map 4.5 Mb apart, suggests that the number of target
337 genes may be less than the number of reported risk loci albeit with, potentially, multiple co-
338 regulated target genes at some loci.

339 Overall it is difficult to evaluate our list of putative target genes when fully understanding the
340 mechanisms by which a given gene influences cancer risk are often complex and require many years'
341 work. It seems likely, however, that the first stage of this process will be short-listing candidates for
342 follow up studies. On that basis, we would argue that a high-throughput CHi-C analysis can
343 contribute to on-going efforts to functionally annotate GWAS risk loci and that CHi-C target genes
344 that are supported by additional data sources are strong candidates for in depth functional follow up
345 studies.

346 **METHODS**

347 **Target Enrichment Array Design**

348 74 SNPs mapping to 68 GWAS risk loci were selected based on all available published GWAS and
349 replication studies as of 31/01/2015. Capture regions were defined as the region that included all
350 SNPs that were correlated ($r^2 \geq 0.2$) with the published SNP based on 1000 Genomes pilot data
351 (<http://www.1000genomes.org/>; Supplementary Table 1). Biotinylated 120-mer RNA baits were
352 designed to target both ends of the HindIII restriction fragments that mapped within these capture
353 regions using Agilent eArray software (Agilent, Santa Clara, CA, USA), using 2x tiling, moderately
354 stringent repeat masking and maximum performance boosting options.

355 **Cell culture and formaldehyde crosslinking**

356 T-47D, ZR-75-1, BT-20 and MDA-MB-231 cell lines were obtained from ATCC (Middlesex, UK),
357 GM06990 cells were supplied by Coriell Cell Repositories (Coriell Institute for Medical Research, New
358 Jersey, USA). Normal Bre80 TERT-immortalised mammary epithelial cells Bre80-Q-TERT (Bre80),
359 were kindly provided by Prof Georgia Chenevix-Trench (Queensland Institute of Medical Research,
360 Brisbane, Queensland, Australia). Cell lines were authenticated using STR genotyping and were
361 regularly tested for mycoplasma contamination. Bre80 cells were grown in DMEM/F12 with phenol
362 red (Gibco, Life Technologies) supplemented with 5% horse serum, 10 μ g/ml insulin, 0.5 μ g/ml
363 hydrocortizone, 20ng/ml epidermal growth factor, 100ng/ml cholera toxin, 50U/ml penicillin and
364 50 μ g/ml streptomycin (Sigma-Aldrich, St. Louis, MO, USA). T-47D and ZR-75-1 were grown in RPMI
365 1640 (Gibco, Life Technologies) supplemented with 10% fetal bovine serum (FBS, Life Technologies),
366 50U/ml penicillin, 50 μ g/ml streptomycin (Sigma-Aldrich, St. Louis, MO, USA) and, for T-47D, 10 μ g/ml
367 insulin (Sigma-Aldrich, St. Louis, MO, USA). BT-20 cells were grown in EMEM (ATCC, Middlesex, UK)
368 supplemented with 10% FBS, 50U/ml penicillin, 50 μ g/ml streptomycin. MDA-MB-231 cells were
369 grown in DMEM supplemented with 10% FBS, 50U/ml penicillin, 50 μ g/ml streptomycin and

370 GM06990 cells were grown in RPMI 1640 supplemented with 15% FBS, 50U/ml penicillin, 50µg/ml
371 streptomycin and 2mM L-Glutamine. Formaldehyde crosslinking of 20 million cells was performed as
372 described by Belton and colleagues⁵² by substituting standard culture media with FBS-free media
373 containing 2% formaldehyde for 5 minutes at room temperature. Crosslinking was quenched by
374 addition of glycine to a final concentration of 150mM. Adherent T-47D, ZR-75-1, BT-20, MDA-MB-
375 231 and Bre80 cells were scraped off the culture flask after crosslinking, non-adherent (GM06990)
376 cells were transferred directly to a falcon tube. Cells were washed with cold PBS, snap-frozen in
377 liquid nitrogen and stored at -80°C before preparation of the Hi-C library.

378 **Hi-C library generation**

379 Each cross-linked cell aliquot (~20 million cells) was re-suspended in 50ml of permeabilisation buffer
380 (10mM Tris-HCl pH8, 10mM NaCl, 0.2% IGEPAL CA-630 (Sigma-Aldrich, St. Louis, MO, USA),
381 supplemented with complete mini EDTA-free tablets (Roche, Basel, Switzerland) and incubated on
382 ice for 30 minutes with occasional mixing. T-47D, ZR-75-1 and GM06990 cells were lysed using 10
383 strokes of a dounce homogeniser. BT-20, MDA-MB-231 and Bre80 cells were lysed by incubating
384 with trypsin (0.25%, Sigma-Aldrich, St. Louis, MO, USA) at 37°C for 5 minutes. Trypsin was
385 inactivated by addition of 500µl FBS. Permeabilised cells were centrifuged for 6 minutes at 600g and
386 washed three times in 1ml 1.3 x NEBuffer 2 (New England Biolabs, Ipswich, MA, USA). Nuclei were
387 resuspended and chromatin digestion and Hi-C library preparation were carried out as described by
388 van Berkum and colleagues⁷ with the following modifications: (i) cells were split into three
389 microcentrifuge tubes instead of five (ii) restriction fragment overhangs were filled in with
390 biotinylated dATP instead of biotinylated dCTP (iii) dGTP was not added to the reaction mixture for
391 the removal of biotinylated dATP from unligated ends (iv) an agarose gel size selection step was not
392 included, and (v) after PCR amplification (5-8 cycles) of the Hi-C library-bound streptavidin beads the
393 PCR product was pooled and subjected to target enrichment (below) before paired end sequencing.

394 **Target enrichment**

395 Target enrichment was performed based on the SureSelect protocol (Agilent, Santa Clara, CA, USA)
396 but incorporating the following modifications: (i) Biotinylated Hi-C ditags bound to streptavidin-
397 beads were amplified pre-hybridization directly from beads using 24 parallel 25µl PCR reactions with
398 five to eight cycles using Q5 High-Fidelity DNA Polymerase (New England Biolabs, Ipswich, MA, USA)
399 and pre-hybridization PCR primers: ACACTCTTCCCTACACGACGCTCTCCGATC*T and
400 CTCGGCATTCTGCTGAACCGCTCTCCGATC*T. PCR products were pooled and purified using
401 Agencourt Ampure XP beads (Beckman Coulter, Brea, CA, USA) to yield approximately 750-1300ng
402 total DNA. 750ng of library DNA was dried using a speedvac concentrator then resuspended in 3.4µl
403 of water. (ii) Enriched fragments were amplified post-hybridization again directly from the
404 streptavidin beads, using 18 parallel 25µl reactions of five to eight cycles of PCR. PCR products were
405 again pooled and purified using Agencourt Ampure XP beads (Beckman Coulter, Brea, CA, USA).
406 Post-hybridization PCR primers to the paired end adaptors were as described in Belton and
407 colleagues⁵²

408 **Paired-end next generation sequencing (NGS), mapping and filtering**

409 12 target enriched Hi-C libraries (two biological replicates for each of six cell lines) were prepared.
410 Eight of the libraries (all at concentrations >2,500 pM) were sequenced on single flow cell lanes on
411 an Illumina HiSeq2000 (Illumina, San Diego, CA, USA) generating 76bp paired-end reads. The other
412 four libraries, which were at lower concentrations (330-630 pM), were sequenced on two flow cell
413 lanes each. Casava software (v1.8, Illumina) was used to make base calls; reads failing the Illumina
414 chastity filter were removed before further analysis. Sequences were output in fastq format before
415 mapping against the human reference genome (GRCh37/hg19) generating between 86 and 153
416 million di-tags with both ends uniquely mapped to the reference genome. Filtering to remove
417 experimental artefacts was carried out using the publicly available Hi-C User Pipeline (HiCUP). Full
418 details of this pipeline are available from Babraham Bioinformatics
419 (<http://www.bioinformatics.babraham.ac.uk/>). In addition to the standard pipeline, off-target di-

420 tags (defined as di-tags where neither end mapped to one of the capture regions) were removed
421 from the final processed data sets. After excluding invalid pairs^{52,53}, PCR duplicates, and off-target
422 di-tags, the number of valid di-tags ranged from 24 to 71 million. Full details of the number and
423 proportion of excluded di-tags are given in Supplementary Table 8.

424 **Analysis of Hi-C interaction peaks**

425 The power of our analysis to detect significant interaction peaks depends on the read density, which
426 in turn depends on the size of the bin or unit of analysis. Given that our purpose was to identify
427 individual target genes, we restricted the analysis to a high resolution (single HindIII fragment)
428 analysis of valid di-tags generated by ligations between a captured fragment and (i) another
429 captured fragment in *cis* or (ii) a non-captured fragment in *cis*, mapping within five Mb⁶. We carried
430 out separate analyses for each type of ligation on the basis that the statistical properties of ligations
431 where both ends of the di-tag have been captured (type (i)) will differ from those where just one end
432 has been captured (type (ii)).

433 To assess the reproducibility of our libraries we calculated Spearman's ρ for each possible
434 combination of HindIII fragments, for each type of analysis ((i) and (ii)) using the two biological
435 replicate libraries for each of the six cell lines. We excluded combinations of HindIII fragments for
436 which there were zero read pairs in both libraries and stratified our analysis on the distance
437 between the two HindIII fragments (0 – 500 kb, 500 kb – 1 Mb, 1 Mb – 1.5 Mb, > 1.5 Mb). The
438 correlation between duplicates was strongest when both fragments were captured and mapped
439 within 500 kb of each other ($\rho = 0.78$ to $\rho = 0.92$). For fragments separated by distances of > 1Mb
440 (where most of the raw di-tags represent “noise”) there was weak or no correlation between
441 replicates (all $\rho < 0.4$); for fragments separated by 500 kb to 1 Mb correlation was moderate ($\rho =$
442 0.53 to $\rho = 0.77$ when both fragments were captured, and $\rho = 0.33$ to $\rho = 0.59$ when just one
443 fragment was captured; Supplementary Figures 6 and 7).

444 There were eight loci annotated by 10 SNPs (5p15.33-rs10069690, 5p15.33-rs7726159 and
445 rs2736108, 11q13.3-rs554219 and rs78540526, 11q13.3-rs75915166, 16q12.2-rs17817449, 16q12.2-
446 rs11075995, 19p13.1-rs8170, 19p13.1-rs2363956) where the capture regions were too close for us
447 to analyse separately. Accordingly we collapsed these eight regions into four regions. There was one
448 region at 10q23.1-rs7071985 (82909977-83064943) that failed to generate high numbers of reads in
449 any of the cell lines we assayed. After excluding this region and combining eight regions into four,
450 there were 63 separate loci for analysis (Supplementary Table 1). On our arrays, there were also
451 1,254 captured HindIII fragments that did not map to known breast cancer risk loci and were not
452 considered further in this study. Three of these fragments comprising the *GSTP1* promoter, mapped
453 within 5Mb of the 11q13.1-rs3903072 capture region and formed interaction peaks with this capture
454 region. For clarity these interaction peaks are excluded from Figure 4. For the 63 risk loci we
455 generated data sets that comprised all di-tags in both categories (type (i) and (ii)) using the SeqMonk
456 mapped sequence analysis tool (www.bioinformatics.babraham.ac.uk/projects/seqmonk/). Where a
457 captured region mapped within five Mb of another captured region we considered HindIII fragments
458 mapping to these two regions as part of a “within capture” (type (i)) analysis.

459 In common with other “C”-based techniques, our Capture Hi-C methodology includes several steps
460 that will show local differences in efficiency thereby introducing biases in the detection of
461 interaction peaks³⁵. To correct for these biases, we used a modification of the procedure described
462 by Sanyal and colleagues^{6,35}. Briefly, our method assumes that some of our captured fragments
463 “fail” and we exclude these; we then use a truncated negative binomial model, which takes account
464 of both the large number of zero counts in the data and allows for overdispersion, to model all
465 ligations for which one end maps an unexcluded captured fragment. R-scripts are available on
466 request. In detail, on the assumption that the majority of *trans* ligations represent random events,
467 we calculated the total number of *trans* ligations (N_{τ}) made by each of the captured HindIII
468 fragments as a measure of the fragment’s “interactability”, its propensity to interact with other
469 fragments. The interactability had a bimodal distribution which we assumed to arise from two

470 components corresponding to low numbers of counts, which we regarded as stochastic noise, and
471 higher numbers of counts, which we regarded as genuine signal. For each cell line and biological
472 replicate a truncated negative binomial distribution, based on the number of di-tags, was fitted to
473 the higher component. By visually inspecting the histogram, it was apparent that the truncation
474 point varied between each cell line and biological replicate. Both the histograms and individual
475 truncation points were used to define an individual threshold, this being the 5% quantile point of the
476 corresponding non-truncated distribution, for each cell line and replicate. All fragments with a total
477 number of trans di-tags below the corresponding threshold value were regarded as noise and were
478 filtered out. This resulted in excluding between 8.6% and 30.0% of the fragments with the lowest
479 number of *trans* ligations (di-tags). We fitted negative binomial regression models to the filtered
480 data sets, combining data from the two biological replicates for each cell line. We corrected for
481 experimental biases due to differing interactability of fragments by including as a covariate the \log_e
482 of the total number of *trans* ligations ($\ln(N_T)$) for each captured fragment from each biological
483 replicate; for *cis* ligations within the capture regions we also included a term for interaction products
484 of $\ln(N_T)$ for each of the two ligated fragments in each biological replicate. We corrected for distance
485 between the ligated fragments by including as a covariate the \log_e of the distance between the mid-
486 points of the two fragments ($\ln(D)$); to approximate local smoothing we fitted the data in bins each
487 of which contained 1 percentile of the distance range. P-values were obtained by comparing the
488 observed counts to the fitted distributions. For each capture region in each cell line, we controlled
489 the false discovery rate using the method of Benjamini & Hochberg⁵⁴. Supplementary Figures 8 and
490 9 show raw read counts aligned to the reported interaction peaks in two libraries ((i) T-47D and (ii)
491 MDA-MB-231 at the 10q26.13-rs2981579 locus (see Figure 2a) and the 11p15.5-rs3817198 locus
492 (see Figure 3a).

493 **Comparison of interaction peaks between cell lines**

494 Non-parametric equality tests (Mann-Whitney for two samples, Kruskal-Wallis for multiple samples)
495 were used to test for a difference in the median number of interaction peaks per locus and the
496 median distance between interacting fragments, across cell lines. We tested the probability of an
497 excess of shared interaction peaks among breast cancer cell lines and all breast-specific cell lines
498 using a random sampling (10,000 permutations) and we estimated the similarity according to
499 receptor status (ER+/ER-) for the breast cancer cell lines using the Jaccard similarity coefficient.

500 **Allocating putative target genes and nearest genes**

501 To define a set of putative target genes, we identified all catalogued RefSeq genes (GRCh37/hg19),
502 mapping within, or *in cis* ($\leq 5\text{Mb}$) to a captured region. From these we selected the subset for which
503 the transcription start-site (TSS) mapped to one end of an interaction peak (an interacting fragment).
504 Given that cancer cell lines are aneuploid, with multiple rearrangements and regions of loss or gain,
505 we further required that the TSS mapped to an interacting fragment in at least two cell lines. For
506 SNPs that mapped to a RefSeq gene (UTR, exon or intron), this gene was considered to be the
507 nearest gene (Table 2). For intergenic SNPs, the nearest gene was determined on the basis of the
508 nearest RefSeq catalogued TSS. Where the nearest catalogued TSS was for a non-coding RNA, this
509 non-coding RNA is listed along with the nearest protein-coding gene (Table 2).

510 **Aligning CHi-C data with TADs.**

511 In order to align CHi-C data with TADs, we accessed Hi-C data generated in HMECs²⁴ through the 3D
512 genome browser (<http://promoter.bx.psu.edu/>).

513 **eQTL Analysis**

514 TCGA breast cancer (BRCA) dataset was used to test for an association between genotype and mRNA
515 abundance further adjusted for DNA methylation and somatic copy number profiles. Pre-processed
516 controlled access germline genotype calls (birdseed algorithm) were downloaded from the TCGA
517 data portal. Putative target genes at each of the risk loci were assigned as described above. For SNPs

518 missing from the Affymetrix SNP6 platform, proxy SNPs were identified using phase 3 data from the
519 1000 Genomes project ($r^2 > 0.8$, distance limit = 500KB). TCGA BRCA mRNA, DNA methylation and
520 DNA copy number data were downloaded from GDAC (version: 2016_01_28). After excluding data
521 from women of Asian (N=37), African (N=159) or American Indian/Alaska Native (N=1) ethnicity,
522 matched data (including germ-line genotype data) were available for 547 samples; 415 ER+ samples
523 and 95 ER- samples (ER status was unknown for 37 samples). mRNA data was \log_2 transformed for
524 eQTL analysis. Statistical association between mRNA abundance levels and genotype groups (AA, AB,
525 BB) was estimated using multivariate linear regression models with one degree of freedom for
526 genotype groups, adjusted for DNA methylation and copy number data. For DNA methylation arrays,
527 methylation levels of the probe with strongest inverse correlation (otherwise minimum correlation
528 coefficient) with its target gene's expression were used as representative methylation levels of the
529 target gene. Analyses for ER+ and ER- subsets were performed separately. P values were adjusted
530 for multiple comparisons using the Benjamini-Hochberg method⁵⁴. eQTL analyses in ER+ samples of
531 genes at chromosome 11q13 were further stratified by copy-number gains using the threshold
532 defined in TCGA²⁵ (\log_2 copy number > 0.3). The variation in copy number within strata was greatly
533 reduced and the eQTL regression models for these additional stratified analyses were as described
534 above, but adjusted for DNA methylation only.

535 **Survival Analysis**

536 The Metabric³⁰ breast cancer cohort (EGA Study ID: EGAS00000000083) was used for disease-
537 specific survival analysis (DSS). Data were summarized and quantile-normalized from the raw
538 expression files generated by Illumina BeadStudio (R packages: beadarray v2.4.2 and illuminaHuman
539 v3.db_1.12.2). Raw data files of one Metabric sample were not available at the time of our analysis,
540 and were therefore excluded. The most variable probe was used as a representative for the
541 corresponding gene's mRNA abundance levels. A Cox proportional hazards model was used to
542 estimate pair-wise hazard ratios with the lowest expression group treated as baseline. P-values for

543 pairwise comparison of survival curves were estimated using Wald tests. The overall test of the Null
544 hypothesis that the expression-derived survival curves show no association with patient outcome
545 was tested with Wald tests (1 degree of freedom, P-trend). P values were further adjusted for
546 multiple comparisons using the Benjamini-Hochberg method⁵⁴. Survival analysis was carried out for
547 ER+ and ER- subsets separately.

548 **Data availability**

549 All Chi-C data sets generated as part of this analysis are publicly available at
550 (<https://www.ebi.ac.uk/ena>) under the accession code PRJEB23968

551 Processed data can be visualised at bit.ly/CHiC-BC/. Publicly available data sets that were accessed
552 for this analysis are detailed in Supplementary Table 9.

554 REFERENCES

- 555 1. DeSantis, C., Ma, J., Bryan, L. & Jemal, A. Breast cancer statistics, 2013. *CA Cancer J Clin* **64**,
556 52-62 (2014).
- 557 2. Michailidou, K. *et al.* Genome-wide association analysis of more than 120,000 individuals
558 identifies 15 new susceptibility loci for breast cancer. *Nat Genet* **47**, 373-80 (2015).
- 559 3. Hindorff, L.A. *et al.* Potential etiologic and functional implications of genome-wide
560 association loci for human diseases and traits. *Proc Natl Acad Sci U S A* **106**, 9362-7 (2009).
- 561 4. Freedman, M.L. *et al.* Principles for the post-GWAS functional characterization of cancer risk
562 loci. *Nat Genet* **43**, 513-8 (2011).
- 563 5. Edwards, S.L., Beesley, J., French, J.D. & Dunning, A.M. Beyond GWASs: illuminating the dark
564 road from association to function. *Am J Hum Genet* **93**, 779-97 (2013).
- 565 6. Dryden, N.H. *et al.* Unbiased analysis of potential targets of breast cancer susceptibility loci
566 by Capture Hi-C. *Genome Res* **24**, 1854-68 (2014).
- 567 7. van Berkum, N.L. *et al.* Hi-C: a method to study the three-dimensional architecture of
568 genomes. *J Vis Exp* (2010).
- 569 8. Ahmed, S. *et al.* Newly discovered breast cancer susceptibility loci on 3p24 and 17q23.2. *Nat*
570 *Genet* **41**, 585-90 (2009).
- 571 9. Antoniou, A.C. *et al.* A locus on 19p13 modifies risk of breast cancer in BRCA1 mutation
572 carriers and is associated with hormone receptor-negative breast cancer in the general
573 population. *Nat Genet* **42**, 885-92 (2010).
- 574 10. Easton, D.F. *et al.* Genome-wide association study identifies novel breast cancer
575 susceptibility loci. *Nature* **447**, 1087-93 (2007).
- 576 11. Fletcher, O. *et al.* Novel breast cancer susceptibility locus at 9q31.2: results of a genome-
577 wide association study. *J Natl Cancer Inst* **103**, 425-35 (2011).
- 578 12. Garcia-Closas, M. *et al.* Genome-wide association studies identify four ER negative-specific
579 breast cancer risk loci. *Nat Genet* **45**, 392-8, 398e1-2 (2013).
- 580 13. Ghoussaini, M. *et al.* Genome-wide association analysis identifies three new breast cancer
581 susceptibility loci. *Nat Genet* **44**, 312-8 (2012).
- 582 14. Haiman, C.A. *et al.* A common variant at the TERT-CLPTM1L locus is associated with estrogen
583 receptor-negative breast cancer. *Nat Genet* **43**, 1210-4 (2011).
- 584 15. Hunter, D.J. *et al.* A genome-wide association study identifies alleles in FGFR2 associated
585 with risk of sporadic postmenopausal breast cancer. *Nat Genet* **39**, 870-4 (2007).
- 586 16. Stacey, S.N. *et al.* Common variants on chromosomes 2q35 and 16q12 confer susceptibility
587 to estrogen receptor-positive breast cancer. *Nat Genet* **39**, 865-9 (2007).
- 588 17. Stacey, S.N. *et al.* Common variants on chromosome 5p12 confer susceptibility to estrogen
589 receptor-positive breast cancer. *Nat Genet* **40**, 703-6 (2008).
- 590 18. Thomas, G. *et al.* A multistage genome-wide association study in breast cancer identifies two
591 new risk alleles at 1p11.2 and 14q24.1 (RAD51L1). *Nat Genet* **41**, 579-84 (2009).
- 592 19. Turnbull, C. *et al.* Genome-wide association study identifies five new breast cancer
593 susceptibility loci. *Nat Genet* **42**, 504-7 (2010).
- 594 20. Zheng, W. *et al.* Genome-wide association study identifies a new breast cancer susceptibility
595 locus at 6q25.1. *Nat Genet* **41**, 324-8 (2009).
- 596 21. Neve, R.M. *et al.* A collection of breast cancer cell lines for the study of functionally distinct
597 cancer subtypes. *Cancer Cell* **10**, 515-27 (2006).
- 598 22. Khan, A. *et al.* JASPAR 2018: update of the open-access database of transcription factor
599 binding profiles and its web framework. *Nucleic Acids Res* (2017).

- 600 23. Hein, R. *et al.* Comparison of 6q25 breast cancer hits from Asian and European Genome
601 Wide Association Studies in the Breast Cancer Association Consortium (BCAC). *PLoS One* **7**,
602 e42380 (2012).
- 603 24. Rao, S.S. *et al.* A 3D map of the human genome at kilobase resolution reveals principles of
604 chromatin looping. *Cell* **159**, 1665-80 (2014).
- 605 25. Cancer Genome Atlas, N. Comprehensive molecular portraits of human breast tumours.
606 *Nature* **490**, 61-70 (2012).
- 607 26. Michailidou, K. *et al.* Large-scale genotyping identifies 41 new loci associated with breast
608 cancer risk. *Nat Genet* **45**, 353-61, 361e1-2 (2013).
- 609 27. Ghousaini, M. *et al.* Evidence that breast cancer risk at the 2q35 locus is mediated through
610 IGFBP5 regulation. *Nat Commun* **4**, 4999 (2014).
- 611 28. Darabi, H. *et al.* Fine scale mapping of the 17q22 breast cancer locus using dense SNPs,
612 genotyped within the Collaborative Oncological Gene-Environment Study (COGs). *Sci Rep* **6**,
613 32512 (2016).
- 614 29. Li, Q. *et al.* Expression QTL-based analyses reveal candidate causal genes and loci across five
615 tumor types. *Hum Mol Genet* **23**, 5294-302 (2014).
- 616 30. Curtis, C. *et al.* The genomic and transcriptomic architecture of 2,000 breast tumours reveals
617 novel subgroups. *Nature* **486**, 346-52 (2012).
- 618 31. Fasching, P.A. *et al.* The role of genetic breast cancer susceptibility variants as prognostic
619 factors. *Hum Mol Genet* **21**, 3926-39 (2012).
- 620 32. Nik-Zainal, S. *et al.* Landscape of somatic mutations in 560 breast cancer whole-genome
621 sequences. *Nature* **534**, 47-54 (2016).
- 622 33. de Laat, W. & Duboule, D. Topology of mammalian developmental enhancers and their
623 regulatory landscapes. *Nature* **502**, 499-506 (2013).
- 624 34. Mifsud, B. *et al.* Mapping long-range promoter contacts in human cells with high-resolution
625 capture Hi-C. *Nat Genet* **47**, 598-606 (2015).
- 626 35. Sanyal, A., Lajoie, B.R., Jain, G. & Dekker, J. The long-range interaction landscape of gene
627 promoters. *Nature* **489**, 109-13 (2012).
- 628 36. Schoenfelder, S. *et al.* The pluripotent regulatory circuitry connecting promoters to their
629 long-range interacting elements. *Genome Res* **25**, 582-97 (2015).
- 630 37. Consortium, E.P. An integrated encyclopedia of DNA elements in the human genome. *Nature*
631 **489**, 57-74 (2012).
- 632 38. Javierre, B.M. *et al.* Lineage-Specific Genome Architecture Links Enhancers and Non-coding
633 Disease Variants to Target Gene Promoters. *Cell* **167**, 1369-1384 e19 (2016).
- 634 39. Horne, H.N. *et al.* Fine-Mapping of the 1p11.2 Breast Cancer Susceptibility Locus. *PLoS One*
635 **11**, e0160316 (2016).
- 636 40. Guo, X. *et al.* Fine-scale mapping of the 4q24 locus identifies two independent loci
637 associated with breast cancer risk. *Cancer Epidemiol Biomarkers Prev* **24**, 1680-91 (2015).
- 638 41. Ghousaini, M. *et al.* Evidence that the 5p12 Variant rs10941679 Confers Susceptibility to
639 Estrogen-Receptor-Positive Breast Cancer through FGF10 and MRPS30 Regulation. *Am J Hum*
640 *Genet* **99**, 903-911 (2016).
- 641 42. Bojesen, S.E. *et al.* Multiple independent variants at the TERT locus are associated with
642 telomere length and risks of breast and ovarian cancer. *Nat Genet* **45**, 371-84, 384e1-2
643 (2013).
- 644 43. Glubb, D.M. *et al.* Fine-scale mapping of the 5q11.2 breast cancer locus reveals at least three
645 independent risk variants regulating MAP3K1. *Am J Hum Genet* **96**, 5-20 (2015).
- 646 44. Dunning, A.M. *et al.* Breast cancer risk variants at 6q25 display different phenotype
647 associations and regulate ESR1, RMND1 and CCDC170. *Nat Genet* **48**, 374-86 (2016).
- 648 45. Shi, J. *et al.* Fine-scale mapping of 8q24 locus identifies multiple independent risk variants
649 for breast cancer. *Int J Cancer* **139**, 1303-17 (2016).

- 650 46. Orr, N. *et al.* Fine-mapping identifies two additional breast cancer susceptibility loci at
651 9q31.2. *Hum Mol Genet* **24**, 2966-84 (2015).
- 652 47. Meyer, K.B. *et al.* Fine-scale mapping of the FGFR2 breast cancer risk locus: putative
653 functional variants differentially bind FOXA1 and E2F1. *Am J Hum Genet* **93**, 1046-60 (2013).
- 654 48. French, J.D. *et al.* Functional variants at the 11q13 risk locus for breast cancer regulate cyclin
655 D1 expression through long-range enhancers. *Am J Hum Genet* **92**, 489-503 (2013).
- 656 49. Zeng, C. *et al.* Identification of independent association signals and putative functional
657 variants for breast cancer risk through fine-scale mapping of the 12p11 locus. *Breast Cancer*
658 *Res* **18**, 64 (2016).
- 659 50. Udler, M.S. *et al.* Fine scale mapping of the breast cancer 16q12 locus. *Hum Mol Genet* **19**,
660 2507-15 (2010).
- 661 51. Lawrenson, K. *et al.* Functional mechanisms underlying pleiotropic risk alleles at the 19p13.1
662 breast-ovarian cancer susceptibility locus. *Nat Commun* **7**, 12675 (2016).
- 663 52. Belton, J.M. *et al.* Hi-C: a comprehensive technique to capture the conformation of
664 genomes. *Methods* **58**, 268-76 (2012).
- 665 53. Wingett, S. *et al.* HiCUP: pipeline for mapping and processing Hi-C data. *F1000Res* **4**, 1310
666 (2015).
- 667 54. Benjamini, Y., Drai, D., Elmer, G., Kafkafi, N. & Golani, I. Controlling the false discovery rate in
668 behavior genetics research. *Behav Brain Res* **125**, 279-84 (2001).

669

670 **URLs**

671 WashU Epigenome Browser: <http://epigenomegateway.wustl.edu/browser/>

672 ENCODE: <https://www.encodeproject.org/>

673 **ACKNOWLEDGMENTS**

674 We thank Prof Georgia Chenevix-Trench for providing Bre80-Q-TERT cells. This work was supported
675 by Breast Cancer Now and Cancer Research UK. We acknowledge National Health Service funding to
676 the NIHR Royal Marsden Biomedical Research Centre. FD and OCL are additionally supported by the
677 MRC (K006215) and the European Union's Horizon 2020 research and innovation programme under
678 grant agreement No 634570 (FORECEE).

679

680 **AUTHOR CONTRIBUTIONS**

681 OF, JSB, NHD, NO, FD and SH were responsible for the overall design of the study. JSB, NHD, NJ, VF,
682 NS, LAM, IA and KF were responsible for data generation. Bioinformatics analyses were carried out

683 by SM, SA, SWW, RC, AGR and SH. OCL, FD, SH and OF were responsible for the statistical analyses.

684 JSB, OF and SH wrote the manuscript with input from all authors.

Table 1: Characteristics of 51 informative risk loci in six cell lines

Cell line	T-47D	ZR-75-1	Bre80 ¹	BT-20	MDA-MB-231	GM06990 ²
Origin	Breast	Breast	Breast	Breast	Breast	Lymphoblastoid
Cancer/normal	Cancer	Cancer	Normal	Cancer	Cancer	Normal
Receptor status	ER+	ER+	ER-	ER-	ER-	ER-
Breast cancer subtype	Luminal	Luminal	Normal	Basal A	Basal B	N/A
Informative loci (%)	34 (66.7)	38 (74.5)	41 (80.4)	21 (41.2)	27 (52.9)	23 (45.1)
Median peaks per locus (range)	7 (0 - 1,107)	9 (0 – 1,744)	10 (0 – 181)	0 (0 – 246)	1 (0 – 466)	0 (0 – 155)
Median distance between interacting fragments ³ (kb)	1392	1647	349	338	388	534
Number (%) of peaks > 2000 kb	1453 (35.4)	1417 (36.4)	94 (7.3)	128 (9.7)	108 (8.0)	102 (12.5)

¹Bre80-Q-TERT (Bre80) are normal Bre80 TERT-immortalised mammary epithelial cells, kindly provided by Prof Georgia Chenevix-Trench (Queensland Institute of Medical Research, Brisbane, Queensland, Australia). ²GM06990 are Epstein-Barr virus transformed B-lymphocytes from the Coriell Cell Repositories (Coriell Institute for Medical Research, New Jersey, USA). ³Range is not given as it was pre-defined to be 10kb to 5Mb.

Table 2: Risk loci which formed interaction peaks directly (N=33), or via an adjacent risk locus (N=3), with 110 target genes.

Locus	SNP(s)	CHi-C target genes	Nearest gene	Agrees
1p36.22	rs616488	<i>APITD1; DFFA, PEX14; PGD</i>	<i>PEX14</i>	v+
1p13.2	rs11552449	<i>OLFML3; HIPK1</i>	<i>DCLREB1</i>	X
2q31.1	rs2016394	<i>CDCA7; DLX2; DYNC1I2</i>	<i>DLX2</i>	v+
2q31.1	rs1550623	<i>CDCA7; DLX2</i>	<i>CDCA7</i>	v+
2q35	rs13387042	<i>IGFBP5</i>	<i>LINC01921, TNP1</i>	X
2q35	rs16857609	<i>IGFBP5; RPL37A</i>	<i>DIRC3, TNS1</i>	X
3p26.1	rs6762644	<i>BHLHE40; CAV3; LINC00312; LMCD1; c3orf32; RAD18; SETD5</i>	<i>ITPR1</i>	X
3p24.1	rs4973768	<i>NGLY1, OXSM; SLC4A7</i>	<i>SLC4A7</i>	v+
4q24	rs9790517	<i>CENPE; PPA2; TET2</i>	<i>TET2</i>	v+
5q11.2	rs889312	<i>MAP3K1</i>	<i>MAP3K1</i>	v
6p23	rs204247	<i>RANBP9</i>	<i>RANBP9</i>	v
6q25.1	rs12662670, rs2046210	<i>ESR1</i>	<i>CCDC170</i>	X
8q21.11	rs6472903	<i>HNF4G/PEX2</i> (adj)	<i>CASC9, HNF4G</i>	v+
8q21.11	rs2943559	<i>HNF4G; PEX2;</i>	<i>HNF4G</i>	v+
8q24.21	rs13281615	<i>CCDC26; CASC11, MYC</i>	<i>CASC8, CASC21, POU5F1B, MYC</i>	v+
8q24.21	rs11780156	<i>CCDC26; CASC11, MYC</i>	<i>PVT1, MYC</i>	v+
9p21.3	rs1011970	<i>CDKN2A; CDKN2B; MTAP</i>	<i>CDKN2B</i>	v+
9q31.2	rs10759243	<i>KLF4</i> (adj)	<i>KLF4</i>	v

9q31.2	rs865686	KLF4	KLF4	√
10p12.31	rs7072776, rs11814448	<i>BMI1; COMMD3; LOC100499489; MIR1915, c10orf114; MLLT10; c10orf140</i>	DNAJC1, MLLT10	√+
10q22.3	rs704010	ZMIZ1	ZMIZ1	√
10q26.13	rs2981579	FGFR2	FGFR2	√
11p15.5	rs3817198	<i>FAM99B, KRTAP5-6; IGF2; KRTAP5-5; MRPL23, SNORD131; TNNT3; LSP1, LINC01150.</i>	LSP1	√+
11q13.1	rs3903072	<i>DKFZp761E198, MIR1234, OVOL1; C11orf68, DRAP1; CCDC85B; CFL1; CTSW, FIBP; FOSL1; KAT5; MUS81, EFEMP2; RNASEH2C; SART1; SNX32; TSGA10IP; CCND1, FADD</i>	SNX32	√+
11q13.3	rs554219, rs78540526, rs75915166	CCND1; LINC01488; ORAOV1; FADD; LOC338694; MIR3164; MRGPRF; MRPL21, IGHMBP2; MYEOV; TPCN2	<i>LINC01488, CCND1</i>	√+
12q24.21	rs1292011	TBX3	TBX3	√
13q13.1	rs11571833	<i>PDS5B</i>	BRCA2	X
14q13.3	rs2236007	PAX9	PAX9	√
14q24.1	rs2588809	<i>ZFP36L1</i>	RAD51B	X
14q24.1	rs999737	<i>ZFP36L1(adj)</i>	RAD51B	X
16q12.2	rs17817449, rs11075995	<i>CRNDE, IRX5; IRX3; LINC02169</i>	FTO	X
17q22	rs6504950	STXBP4, COX11; TOM1L1	STXBP4	√+
19p13.1	rs8170, rs2363956	<i>ANO8, GTPBP3; DDA1; MRPL34; NR2F6 ; USE1, OCEL1</i>	BABAM1, ANKLE1	X
19p13.11	rs4808801	<i>LRRC25; SSBP4, ISYNA1; LSM4; MRPL34; PGPEP1, GDF15, MIR3189; UPF1</i>	ELL	X
19q13.31	rs3760982	KCNN4	KCNN4	√
22q13.1	rs6001930	<i>LOC101927257</i>	MLK1	X

Where TSS for two or more target genes map to a single HindIII fragment, the genes are separated by a comma. Non-coding RNAs (long non-coding RNAs, microRNAs and small nucleolar RNAs) are indicated in green. There were three loci at which the target gene is assigned indirectly on the basis of interaction peaks with an adjacent locus; these are indicated by (adj). Defining nearest gene; for SNPs that map within a gene (UTR, exons or introns) this gene is considered to be the nearest gene, for SNPs that do not map within a gene, nearest gene is assigned based on the location of the nearest TSS according to RefSeq genes (GRCh37/hg19). Where the nearest gene is a non-coding RNA, the nearest protein-coding gene is also given. Chi-C target genes that are also the nearest gene are indicated in bold. Chi-C targets and the nearest gene are compared in the “Agrees” column; V = Chi-C data were consistent with the nearest gene being the sole target gene, V+ = Chi-C data were consistent with the nearest gene being one of several target genes, X = Chi-C data support a gene other than the nearest gene as a target.

Table 3: Nine Chi-C putative target genes that were statistically significant eQTLs (FDR adjusted P < 0.1)

Cytoband	SNP	Proxy	Gene		All cancers		ER+ cancers		ER- cancers	
			Nearest	Chi-C target	P	P _{adj}	P	P _{adj}	P	P _{adj}
2q31.1	rs1550623		<i>CDCA7</i>	<i>CDCA7</i>	0.007	0.087	0.511	0.666	0.330	0.892
11q13.1	rs3903072		<i>SNX32</i>	<i>CTSW</i>	0.006	0.087	0.064	0.326	0.001	0.101
11q13.1	rs3903072		<i>SNX32</i>	<i>SNX32</i>	0.007	0.087	0.032	0.268	0.036	0.506
14q13.3	rs2236007	rs1018464	<i>PAX9</i>	<i>PAX9</i>	0.003	0.066	0.054	0.317	0.248	0.854
14q24.1	rs2588809		<i>RAD51B</i>	<i>ZFP36L1</i>	0.079	0.380	0.004	0.091	0.256	0.854
17q22	rs6504950	rs9902718	<i>STXBP4</i>	<i>COX11</i>	0.002	0.059	0.001	0.032	0.403	0.892
19p13.11	rs8170	rs34084277	<i>BABAM1</i> , <i>ANKLE1*</i>	<i>MRPL34</i>	0.001	0.059	0.011	0.173	0.131	0.829
19p13.11	rs4808801		<i>ELL*</i>	<i>LRRC25</i>	0.009	0.092	0.004	0.091	0.768	0.954
19p13.11	rs4808801		<i>ELL*</i>	<i>SSBP4</i>	0.002	0.059	0.0002	0.016	0.475	0.892

* in an analysis that included all genes that are nearest genes, regardless of whether they were also a Chi-C target gene, rs4808801 was also associated with expression of *ELL* (FDR adjusted P = 0.05 for all cancers and P = 0.04 for ER+ cancers) and rs8170 was also associated with expression of *ANKLE1* (FDR corrected P = 0.05 for all cancers). P = P value (1df t-test) per allele association with gene expression, adjusted for methylation and copy number; P_{adj} = P value further adjusted for multiple testing

Table 4: Six Chi-C putative target genes for which there was orthogonal support for at least two additional data sources

Locus	SNP	Gene	eQTL P _{all}	eQTL P _{ER+}	DSS P _{ER+}	727 cancer genes source
2q31.1	rs1550623	<i>CDCA7</i>	0.007 (0.09)	0.51 (0.67)	4 x 10 ⁻⁷	
11q13.1	rs3903072	<i>FADD</i>	0.04 (0.28)	0.01* (0.17)	0.0009	cancer related genes panel
12q24.21	rs1292011	<i>TBX3</i>	0.28 (0.50)	0.21 (0.52)	0.012	cancer gene census
14q13.3	rs2236007	<i>PAX9</i>	0.003 (0.07)	0.05 (0.32)	0.20	cancer related genes panel
14q24.1	rs2588809	<i>ZFP36L1</i>	0.08 (0.38)	0.004 (0.09)	0.09	identified in Nik-Zainal et al 2016
19p13.11	rs8170	<i>MRPL34</i>	0.001 (0.06)	0.01 (0.17)	0.004	

eQTL P values in parenthesis are FDR adjusted; *excluding 119 ER+ cancers with copy-number gains at *FADD*, P = 0.004